

Answers to Test 2

1. (i) (a) $p_\theta = \partial L / \partial \dot{\theta} = \dot{\theta}$, $p_\phi = \partial L / \partial \dot{\phi} = \sin^2 \theta \dot{\phi}$ so that

$$\begin{aligned} H = p_\theta \dot{\theta} + p_\phi \dot{\phi} - L &= p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} - \frac{1}{2} \left[p_\theta^2 + \sin^2 \theta \left(\frac{p_\phi}{\sin^2 \theta} \right)^2 \right] - g \cos \theta \\ &= \frac{p_\theta^2}{2} + \frac{p_\phi^2}{2 \sin^2 \theta} - g \cos \theta. \end{aligned}$$

[4 marks]

- (b) Hamilton's equations:

$$\begin{aligned} \dot{\theta} &= \frac{\partial H}{\partial p_\theta} = p_\theta, & \dot{\phi} &= \frac{\partial H}{\partial p_\phi} = \frac{p_\phi}{\sin^2 \theta}, \\ \dot{p}_\theta &= -\frac{\partial H}{\partial \theta} = \frac{\cos \theta}{\sin^3 \theta} p_\phi^2 - g \sin \theta, & \dot{p}_\phi &= -\frac{\partial H}{\partial \phi} = 0. \end{aligned}$$

[5 marks]

- (c) $p_\theta = 0$ requires that θ is constant. p_ϕ is constant for all solutions. If θ is constant then so is $\dot{\phi} = p_\phi / \sin^2 \theta$. These solutions represent horizontal circular orbits. We have

$$0 = \dot{p}_\theta = \frac{\cos \theta}{\sin^3 \theta} p_\phi^2 - g \sin \theta,$$

so that $p_\phi^2 = g \sin^4 \theta / \cos \theta$ giving

$$p_\phi = \pm \sin^2 \theta \sqrt{\frac{g}{\cos \theta}} \quad \text{and} \quad \dot{\phi} = \pm \sqrt{\frac{g}{\cos \theta}}.$$

[5 marks]

- (ii) (a)

$$\{Q, P\} = \frac{\partial Q}{\partial q} \frac{\partial P}{\partial p} - \frac{\partial Q}{\partial p} \frac{\partial P}{\partial q} = \cos t \cdot \cos t - \sin t \cdot (-\sin t) = 1.$$

[4 marks]

- (b) From the first equation

$$p = \frac{Q}{\sin t} - q \cot t.$$

Inserting this into the second equation gives

$$P = -q \sin t + \left(\frac{Q}{\sin t} - q \cot t \right) \cos t = -\frac{q}{\sin t} + Q \cot t.$$

Accordingly

$$F(q, Q, t) = \frac{qQ}{\sin t} - \frac{q^2 + Q^2}{2} \cot t.$$

[5 marks]

(c) It is not defined if t is a multiple of π ; here the canonical transformation is the identity transformation or $Q = -q$, $P = -p$ (the identity plus a sign flip).

[2 marks]

[Total: 25 marks]