

## 3 Sequences

A sequence  $(a_n)_{n \geq 1}$  of real (or complex, etc.) numbers is an infinite list of numbers  $a_1, a_2, a_3, \dots$  all in  $\mathbb{R}$  (or  $\mathbb{C}$ , etc.) Formally:

**Definition.** A *sequence* is a function  $a : \mathbb{N}_{>0} \rightarrow \mathbb{R}$ .

**Notation:** We let  $a_n \in \mathbb{R}$  denote  $a(n)$  for  $n \in \mathbb{N}_{>0}$ . The data  $(a_n)_{n=1,2,\dots}$  is equivalent to the function  $a : \mathbb{N}_{>0} \rightarrow \mathbb{R}$  because a function  $a$  is determined by its values  $a_n$  over all  $n \in \mathbb{N}_{>0}$ .

We will denote  $a$  by  $a_1, a_2, a_3, \dots$  or  $(a_n)_{n \in \mathbb{N}_{>0}}$  or  $(a_n)_{n \geq 1}$  or even just  $(a_n)$ .

*Remark 3.1.*  $a_i$ s could be repeated, so  $(a_n)$  is *not* equivalent to the set  $\{a_n : n \in \mathbb{N}_{>0}\} \subset \mathbb{R}$ . E.g.  $(a_n) = 1, 0, 1, 0, \dots$  is different from  $(b_n) = 1, 0, 0, 1, 0, 0, 1, \dots$ . This is why we use round brackets  $( )$  instead of  $\{ \}$ .

We can describe a sequence in many ways,

- As a **list**  $1, 0, 1, 0, \dots$ ,
- Via a **closed formula**, like  $a_n = \frac{1-(-1)^n}{2}$  for the sequence above,
- By a **recursion**, e.g. the Fibonacci sequence  $F_1 = 1 = F_2$ ,  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 3$  (so  $(F_n)$  is  $1, 1, 2, 3, 5, 8, 13, \dots$ )
- By a summation, e.g.  $a_n = \sum_{i=1}^n \frac{1}{i} = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$ . Such a sequence  $a_n = \sum_{i=1}^n b_i$  is called a **series** and will be studied later in the course.


**Exercise 3.2.** Show any sequence  $(a_n)$  can be written as a series  $a_n = \sum_{i=1}^n b_i$  for an appropriate choice of sequence  $(b_n)$ .

### 3.1 Convergence of Sequences

We want to *rigorously* define  $a_n \rightarrow a \in \mathbb{R}$ , or “ $a_n$  converges to  $a$  as  $n \rightarrow \infty$ ” or “ $a$  is the limit of  $(a_n)$ ”. We will spend a while exploring various formulations before we choose our definitive definition.

**Idea 1:**  $a_n$  should get closer and closer to  $a$ . Not necessarily monotonically, e.g. for:

$$a_n = \begin{cases} \frac{1}{n} & n \text{ odd} \\ \frac{1}{2n} & n \text{ even} \end{cases} \quad \text{we want } a_n \rightarrow 0.$$



**Idea 2:** But notice that  $\frac{1}{n}$  also gets closer and closer to  $-73.6$ ! So we want to say instead that  $a_n$  gets “as close as we like to  $a$ ” or “arbitrarily close to  $a$ ”. We will measure this with  $\epsilon > 0$ : we say  $a_n$  gets to within  $\epsilon$  of  $a$  by

$$|a_n - a| < \epsilon \quad \text{or} \quad a_n \in (a - \epsilon, a + \epsilon).$$

We phrase “ $a_n$  gets arbitrarily close to  $a$ ” by “ $a_n$  gets to within  $\epsilon$  of  $a$  for **any**  $\epsilon > 0$ ”. This suggests the following definition.


**Exercise 3.3.** Dedekind tries to define  $a_n \rightarrow a$  if and only if  $\forall n$  sufficiently large,  $|a_n - a|$  is *arbitrarily small*. When pushed they define a real number  $b \in \mathbb{R}$  to be arbitrarily small if it is smaller than any  $\epsilon > 0$  i.e.  $\forall \epsilon > 0, |b| < \epsilon$ .

Leaving aside what he means by “sufficiently large” for now, which of these sequences converges (to some  $a \in \mathbb{R}$ ) according to their definition?

1.  $0, 1, 0, 1, \dots$
2.  $1, 1, 1, 1, \dots$
3.  $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$
4.  $a_n = 2^{-n}$
5. More than one of these
6. None of these

**Idea 3:** Prof B said that once  $n$  is large enough,  $|a_n - a|$  is less than every  $\epsilon > 0$ , but that means it’s zero, i.e.  $a_n = a$ . The problem he missed is that if we take smaller  $\epsilon$  we will usually have to take bigger  $n$  to make  $|a_n - a| < \epsilon$ .

So we want to say that to get *arbitrarily close to the limit  $a$*  (i.e.  $|a_n - a| < \epsilon$ ), we need to go sufficiently far down the sequence. If I change  $\epsilon > 0$  to be smaller, I simply go further down the sequence to get within  $\epsilon$  of  $a$ .



Don't fall for the same trap as Dedekind - there will not be a “ $n$  sufficiently large” that works for all  $\epsilon$  at once! (Unless  $a_n \equiv a$  eventually.)

That is, we want to *reverse* the order of specifying  $n$  and  $\epsilon$ : only once we've seen how small  $\epsilon$  is do we know how big to take  $n$ . If we chose a smaller  $\epsilon$  we can then choose a larger  $n$ .

For *any* (fixed)  $\epsilon > 0$  we want there to be an  $n$  sufficiently large such that  $|a_n - a| < \epsilon$ . So we change “ $\exists n$  such that  $\forall \epsilon > 0$ ” to “ $\forall \epsilon > 0, \exists n$ ”. *This allows  $n$  to depend on  $\epsilon$ .*

**Exercise 3.4.** Professor Buzzard takes your point, and modifies his definition of  $a_n \rightarrow a$  to

$$\forall \epsilon > 0 \exists n \in \mathbb{N}_{>0} \text{ such that } |a_n - a| < \epsilon.$$

Which of these sequences converges to  $a = 0$  according to his new definition?

1.  $0, 1, 0, 1, \dots$
2.  $1, 1, 1, 1, \dots$
3.  $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$
4.  $a_n = 2^{-n}$
5. More than one of these
6. None of these

**Idea 4:** So we measure “*eventually*” (or “sufficiently large”) by a point  $N \in \mathbb{N}_{>0}$  beyond which (“ $\forall n \geq N$ ”)  $a_n$  **stays** within  $\epsilon$  of  $a$ . That is

**Definition** (Convergence)

We say that  $a_n \rightarrow a$  as  $n \rightarrow \infty$  if and only if

Read this as follows:

*However close* ( $\forall \epsilon > 0$ ) I want to get to the limit  $a$ , there's a point in the sequence ( $\exists N \in \mathbb{N}_{>0}$ ) beyond which ( $n \geq N$ ) *all*  $a_n$  are indeed that close to the limit  $a$  ( $|a_n - a| < \epsilon$ ).


*Remark 3.5.*  $N$  depends on  $\epsilon$ ! For a while we will sometimes denote it  $N_\epsilon$ , as a reminder. We often write ( $a_n \rightarrow a$  as  $n \rightarrow \infty$ ) as just ( $a_n \rightarrow a$ ) or ( $\lim_{n \rightarrow \infty} a_n = a$ ).

Equivalently:

or equivalently

**Example 3.6.** Prove  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ .

*Rough working:* Fix  $\epsilon > 0$ . I want to find  $N_\epsilon \in \mathbb{N}_{>0}$  such that  $|a_n - a| = |\frac{1}{n} - 0| = \frac{1}{n} < \epsilon$  for all  $n \geq N_\epsilon$ .



**How to prove**  $a_n \rightarrow a$ 

- (I) Fix  $\epsilon > 0$ .
- (II) Calculate  $|a_n - a|$ .
- (II') Find a good estimate  $|a_n - a| \leq b_n$ .
- (III) Try to solve  $b_n < \epsilon$ .      (\*)
- (IV) Find  $N_\epsilon \in \mathbb{N}_{>0}$  such that (\*) holds whenever  $n \geq N_\epsilon$ .
- (V) Put everything together into a logical proof (usually involves rewriting everything in reverse order - see examples below).

Notice you only have to do this for **one**  $\epsilon > 0$ , so long as it is arbitrary; that way you've done it for **any**  $\epsilon > 0$ .

The key point is to choose  $b_n$  so that solving  $b_n < \epsilon$  is easier than solving  $|a_n - a| < \epsilon$ .

**Example 3.7.** Prove that  $a_n = \frac{n+5}{n+1} \rightarrow 1$ .

*Rough working:*

*Proof.*

**Example 3.8.** Prove that  $a_n = \frac{n+2}{|n-2|} \rightarrow 1$ .

*Rough working:*

(Notice using  $2 < n$  here would ruined everything.)

*Proof.*

**Definition.** We say that  $a_n$  *converges* if and only if  $\exists a \in \mathbb{R}$  such that  $a_n \rightarrow a$ , i.e.

Negating the above statement gives the following

**Definition.** We say  $a_n$  *diverges* if and only if it does not converge (to any  $a \in \mathbb{R}$ ), i.e.

*Remark 3.9.* Notice *diverge* does not mean  $\rightarrow \pm\infty$ , for instance we will prove later that  $a_n = (-1)^n$  diverges.

**Exercise 3.10.** Fix a sequence of real numbers  $(a_n)_{n \geq 1}$ . Consider

$$\forall n \geq 1 \exists \epsilon > 0 \text{ such that } |a_n| < \epsilon$$

This means?

1.  $a_n \rightarrow 0$
2.  $(a_n)_{n \geq 1}$  is bounded
3. Precisely nothing
4. More than one of these
5. None of these

**Exercise 3.11.** What about

$$\exists \epsilon > 0 \text{ such that } \forall n \geq 1, |a_n| < \epsilon?$$

1.  $a_n \rightarrow 0$
2.  $(a_n)_{n \geq 1}$  is bounded
3. Precisely nothing
4. More than one of these
5. None of these



We can also define limits for *complex sequences*. Let  $|z| := \sqrt{(\operatorname{Re} z)^2 + (\operatorname{Im} z)^2}$ .

**Definition.**  $a_n \in \mathbb{C}$ ,  $\forall n \geq 1$ . We say  $a_n \rightarrow a \in \mathbb{C}$  if and only if

$$\forall \epsilon > 0, \exists N \in \mathbb{N}_{>0} \text{ such that } n \geq N \implies |a_n - a| < \epsilon.$$

This definition is equivalent to  $(\operatorname{Re} a_n) \rightarrow \operatorname{Re} a$  and  $(\operatorname{Im} a_n) \rightarrow \operatorname{Im} a$  (see problem sheet 4!).

**Example 3.12.** Prove  $a_n = \frac{e^{in}}{n^3 - n^2 - 6} \rightarrow 0$  as  $n \rightarrow \infty$ .

*Rough working:*

*Proof.*


**Example 3.13.** Set  $\delta = 10^{-1000000}$ ,  $a_n = (-1)^n \delta$ . Prove that  $a_n$  diverges, that is it does not converge (to any  $a \in \mathbb{R}$ ).



Assume for contradiction that  $a_n \rightarrow a$ , i.e.

$$\forall \epsilon > 0 \exists N \in \mathbb{N}_{>0} \text{ such that } n \geq N \implies |a_n - a| < \epsilon.$$

*Rough working:* Draw a picture! But don't make  $\delta$  small in your picture, as then you won't see the contradiction. Magnify it to be big.




For small enough  $\epsilon > 0$  (the picture shows that any  $\epsilon \leq \delta$  will do), the fact that  $a$  is within  $\epsilon$  of  $\delta$  ( $a_{2n}$ ) and  $-\delta$  ( $a_{2n+1}$ ) will be a contradiction.

*Proof 1.* Fix  $a \in \mathbb{R}$ . Take  $\epsilon = \delta$ .

Then if  $\exists N$  such that  $\forall n \geq N$ ,  $|a_n - a| < \epsilon$  this implies

1.  $|a_{2N} - a| < \epsilon \iff a \in (\delta - \epsilon, \delta + \epsilon) \implies a > \delta - \epsilon = 0$ , and
2.  $|a_{2N+1} - a| < \epsilon \iff a \in (-\delta - \epsilon, -\delta + \epsilon) \implies a < -\delta + \epsilon = 0 \quad \times$

So  $a_n \not\rightarrow a$ , but this holds  $\forall a \in \mathbb{R}$ , so  $a_n$  does not converge.



Or, *Proof 2:* Both  $\pm\delta$  close to the limit  $a$  so must be close to each other by the triangle inequality:

$$|\delta - (-\delta)| \leq |\delta - a| + |a - (-\delta)| < \epsilon + \epsilon \implies 2\delta < 2\epsilon = 2\delta \quad \times$$

So  $a_n \not\rightarrow a$ , but this holds  $\forall a \in \mathbb{R}$ , so  $a_n$  does not converge.  $\square$

An alternative approach to that question is provided by the following.

**Theorem 3.14: Uniqueness of Limits**

Limits are unique. If  $a_n \rightarrow a$  and  $a_n \rightarrow b$ , then  $a = b$ .


*Idea:* For  $n$  large,  $a_n$  is arbitrarily close to both  $a$  and  $b$ . So  $a$  arbitrarily close to  $b \implies a = b$ .

*Proof 1.*

Proof of this last claim:

$\square$

*Proof 2.* By contradiction. Assume  $a \neq b$  and again draw a *magnified* picture.



Eventually  $a_n$  is in *both* corridors. So if we choose  $\epsilon$  sufficiently small so that the corridors don't overlap then we get a contradiction.

**Exercise 3.15.** Let  $a_n$  be defined by  $a_1 = a_2 = 0$  and  $a_n = \frac{1}{n-2}$  for  $n > 2$ . Show  $a_n \rightarrow 0$ .

Which step is incorrect in this student's strategy?


Fix  $\epsilon > 0$ . We assume  $n > 2$ . Then

1. We want  $|\frac{1}{n-2} - 0| = \frac{1}{n-2} < \epsilon$
2.  $\implies n - 2 > 1/\epsilon$
3.  $\implies n > 2 + 1/\epsilon$
4.  $\implies n > 1/\epsilon \quad (*)$
5. So take  $N > \max(1/\epsilon, 2)$ , then
6.  $\forall n \geq N, n > 1/\epsilon$  which is  $(*)$
7. So  $\frac{1}{n-2} \rightarrow 0$
8. More than one mistake
9. All correct

**Proposition 3.16.** *If  $(a_n)$  is convergent, then it is bounded.*

[I.e.  $a_n \rightarrow a \implies \exists A \in \mathbb{R}$  such that  $|a_n| \leq A \forall n$ .]

*Proof.* Fix  $\epsilon = 1$ . Then  $\exists N \in \mathbb{N}$  such that  $\forall n \geq N, |a_n - a| < 1 \implies |a_n| < 1 + |a|$ .



Then  $|a_n|$  is bounded  $\forall n$  by  $\max\{|a_1|, |a_2|, \dots, |a_{N-1}|, |a| + 1\}$ .  $\square$

Notice  $a_n = \frac{1}{n-7}$  is not a counterexample! It is not a well defined sequence of real numbers because  $a_7$  is either not defined or not real. Instead we could take

$$a_n = \begin{cases} \frac{1}{n-7} & n \neq 7, \\ 0 & n = 7. \end{cases}$$

This is then indeed bounded as  $\forall n \in \mathbb{N}_{>0}$  we have

$$-1 = a_6 \leq a_n \leq a_8 = 1.$$

**Exercise 3.17.** Give an example of a bounded sequence that is divergent.

**Exercise 3.18.** Let  $(a_n)$  be a bounded sequence. Let  $(b_n)$  be a sequence with  $b_n = a_n$  for all  $n \geq 100$ . Prove that  $b_n$  is bounded.