

# Introduction to Quantum Mechanics – Problem sheet 2

## 1. A quantum wave function

Our system is a quantum particle in one dimension described by the wave function

$$\psi(x, t_0) = Nx e^{-\frac{(x-x_0)^2}{2}},$$

with real constants  $N > 0$ , and  $x_0 > 0$ .




Figure 1: A plot of  $|\psi(x)|^2$  for  $x_0 = 0.5$  and  $N = 1$ .

- (a)
- (b) The most probable location is the absolute maximum of the probability density. We first look for extrema of the probability density, where it holds

$$\frac{d}{dx}|\psi(x)|^2 = 0.$$

This gives

$$N^2(-2(x-x_0)x^2 + 2x)e^{-(x-x_0)^2} = 0,$$

which we solve to obtain

$$x = 0, \frac{1}{2}(x_0 \pm \sqrt{4 + x_0^2}).$$

For the value  $x_0 = 0.5$  the maximum at  $x = \frac{1+\sqrt{17}}{16}$  is the absolute maximum, consistent with the plot in part (a).

## 2. A quantum wave function, probabilities and energies




Figure 2: Plot of  $\psi(x)$  for  $a = b = 1$

(a)

(b) We have

$$\int_{-\infty}^{\infty} |\psi(x)|^2 dx = 1$$

$$\begin{aligned} \int_{-\infty}^{\infty} |\psi(x)|^2 dx &= \int_{-a}^a b^2 (a^4 - 2a^2 x^2 + x^4) dx \\ &= b^2 \left[ a^4 x - \frac{2}{3} a^2 x^3 + \frac{1}{5} x^5 \right]_{-a}^a \\ &= b^2 \left[ 2a^5 - \frac{4}{3} a^5 + \frac{2}{5} a^5 \right] \\ &= \frac{16}{15} a^5 b^2 \end{aligned}$$

Therefore

$$b = \frac{\sqrt{15}}{4} a^{-\frac{5}{2}}.$$

(c) Calculate the probability of finding the particle between  $-a/2$  and  $a/4$ .

$$\begin{aligned} \int_{-a/2}^{a/4} |\psi(x)|^2 dx &= \frac{15}{16} a^{-5} \left[ a^4 x - \frac{2}{3} a^2 x^3 + \frac{1}{5} x^5 \right]_{-a/2}^{a/4} \\ &= \frac{2813}{16384} \end{aligned}$$

(d) We have

$$\frac{\partial^2}{\partial x^2} \psi(x) = -2b$$

The expectation of the Hamiltonian is given by

$$\begin{aligned}
\langle H \rangle &= -\frac{\hbar^2}{2m} \int_{-\infty}^{+\infty} \psi^*(x) \psi''(x) dx \\
&= \frac{\hbar^2}{2m} \int_{-a}^a 2b^2(a^2 - x^2) dx \\
&= \frac{\hbar^2 b^2}{m} \left[ a^2 x - \frac{1}{3} x^3 \right]_{-a}^a \\
&= \frac{\hbar^2}{m} \frac{5}{4} a^{-2}
\end{aligned}$$

#### 4. Stationary states and complex energies

Consider the two stationary states

$$\begin{aligned}
\psi_A(x, t) &= e^{-iE_A t} \psi_A(x, 0) \\
\psi_B(x, t) &= e^{-iE_B t} \psi_B(x, 0),
\end{aligned}$$

where  $E_A \neq E_B$ .

- (a) Calculate the probability densities of  $\psi_A$ ,  $\psi_B$ ,  $\psi_A + \psi_B$  and  $\psi_A - \psi_B$ .

$$|\psi_A(x, t)|^2 = e^{-iE_A t} e^{iE_A t} |\psi_A(x, 0)|^2 = |\psi_A(x, 0)|^2$$

$$|\psi_B(x, t)|^2 = |\psi_B(x, 0)|^2$$

$$\begin{aligned}
|\psi_A(x, t) + \psi_B(x, t)|^2 &= (\psi_A(x, t) + \psi_B(x, t))(\psi_A(x, t)^* + \psi_B(x, t)^*) \\
&= |\psi_A(x, 0)|^2 + |\psi_B(x, 0)|^2 + 2\text{Re}(e^{i(E_A - E_B)t} \psi_A(x, 0) \psi_B(x, 0)^*) \\
&= |\psi_A(x, 0)|^2 + |\psi_B(x, 0)|^2 + 2\cos((E_A - E_B)t) \text{Re}(\psi_A(x, 0) \psi_B(x, 0)^*)
\end{aligned}$$

$$\begin{aligned}
|\psi_A(x, t) - \psi_B(x, t)|^2 &= (\psi_A(x, t) - \psi_B(x, t))(\psi_A(x, t)^* - \psi_B(x, t)^*) \\
&= |\psi_A(x, 0)|^2 + |\psi_B(x, 0)|^2 - 2\text{Re}(e^{i(E_A - E_B)t} \psi_A(x, 0) \psi_B(x, 0)^*) \\
&= |\psi_A(x, 0)|^2 + |\psi_B(x, 0)|^2 - 2\cos((E_A - E_B)t) \text{Re}(\psi_A(x, 0) \psi_B(x, 0)^*)
\end{aligned}$$

- (b) We have

$$\psi_A(x, t) = e^{(-iE_R + E_I)t} \psi_A(x, 0).$$

The Probability density of  $\psi_A(x, t)$  is therefore given by

$$|\psi_A(x, t)|^2 = e^{2E_I t} |\psi_A(x, 0)|^2.$$

We see that the norm of the state, that is the overall probability is time dependent. In particular the probability decays exponentially with time for  $E_I < 0$  and growth exponentially with time for  $E_I > 0$ . While a growth of the overall probability is a little strange and it is difficult to imagine physical situations corresponding to this, an exponential decay of the overall probability arises in some familiar situations, such as nuclear decay.