

MVC: Hints, tips, answers Sheet 4

1.

1/ Consider Jacobian $\begin{pmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{pmatrix}$
& show that $\det(J) \neq 0$ when $(u,v) = (1,0)$.

To find u_x, v_x diff. implicitly wrt x
and then rewrite as

$$\begin{pmatrix} \quad & \quad \end{pmatrix} \begin{pmatrix} u_x \\ v_x \end{pmatrix} = \begin{pmatrix} \quad \end{pmatrix}$$

Then invert to get $\begin{pmatrix} u_x \\ v_x \end{pmatrix}$ after substituting $(u,v) = (1,0)$.

To find u_y, v_y diff expressions for x & y wrt y
and proceed as above.

2/ (i) Diff implicitly wrt x . Same method as Q1
leads to

$$\begin{pmatrix} 2u & 4v \\ v_y & uy - xy \end{pmatrix} \begin{pmatrix} u_x \\ v_x \end{pmatrix} = \begin{pmatrix} 2x \\ v_y \end{pmatrix}$$

Soln exists provided \det of 2×2 matrix $\neq 0$
when evaluated at (x_0, y_0, u_0, v_0) .

Diff wrt y to get similar matrix equation
for $\begin{pmatrix} u_y \\ v_y \end{pmatrix}$.

Then show that $u^2 + 2v^2 + y^2 - x^2 = 6$ & $uvy - vxy = 1$
are satisfied for both given choices of (x_0, y_0, u_0, v_0) .
For $(1, 1, 2, 1)$ The determinant is $\begin{vmatrix} 4 & 4 \\ 1 & 1 \end{vmatrix} = 0$

For $(1+\sqrt{2}, -1-\sqrt{2}, 2, 1)$ we have $\begin{vmatrix} 4 & 4 \\ -1-\sqrt{2} & 1 \end{vmatrix} \neq 0$

(ii) $\begin{pmatrix} \partial F_1/\partial u & \partial F_1/\partial v \\ \partial F_2/\partial u & \partial F_2/\partial v \end{pmatrix} = \begin{pmatrix} 2u & 4v \\ v_y & uy - xy \end{pmatrix}$ and so is
the same as matrix in (i)

(iii) Think about whether map is 1-1.

MVC: Hints & tips
& answers
for Sheet 4

(full
solutions
to follow
later)

20

3/ Find $\frac{\partial \underline{r}}{\partial \xi}$, $\frac{\partial \underline{r}}{\partial \eta}$ & show dot product is zero.

$$\text{Then } h_1 = \left| \frac{\partial \underline{r}}{\partial \xi} \right| = \dots = \frac{c}{\cosh \xi - \cos \eta}$$

$$h_2 = h_1$$

$$h_3 = 1$$

4/ Can proceed as in Q3
to find $h_1 = h_2 = (u^2 + v^2)^{1/2}$, $h_3 = 1$.

$$\text{Then } \hat{e}_1 = \frac{\partial \underline{r} / \partial u}{|\partial \underline{r} / \partial u|} = \frac{u \hat{i} + v \hat{j}}{(u^2 + v^2)^{1/2}}$$

$$\text{Similarly } \hat{e}_2 = (-v \hat{i} + u \hat{j}) / ()^{1/2}$$

$$\& \hat{e}_3 = \hat{k}.$$

5/ (i), (ii)
use expressions for div, curl in curvilinear coordinates

(iii) use unit vectors from Q4

$$\text{In Cartesians, } \text{div } \underline{F} = 8x, \text{Curl } \underline{F} = -8y \hat{k}.$$

$$6/ \hat{r} = (\cos \varphi) \hat{i} + (\sin \varphi) \hat{j}$$

$$\hat{\varphi} = -\hat{i} \sin \varphi + \hat{j} \cos \varphi, \hat{z} = \hat{k}$$

$$\& \text{ then } \hat{i} = \hat{r} \cos \varphi - \hat{\varphi} \sin \varphi$$

$$\hat{j} = \hat{r} \sin \varphi + \hat{\varphi} \cos \varphi$$

$$\& F_r = r \sin \varphi \cos \varphi + z \sin \varphi, F_\varphi = -r \sin^2 \varphi + z \cos \varphi, F_z = r \cos \varphi$$

MVC: Hints, tips, answers for Sheet 4

13.

7. Find $|\det J| = 4(x^2 + y^2)$. Integral $\rightarrow \frac{1}{4} \int_{v=2}^{v=4} \int_{u=-1}^1 (u^2 + v^2) du dv$
 $= \dots = 29/3$.

8. $\det J = r$. Integral $\rightarrow \int_0^{2\pi} \int_0^1 (r^4 \cos^4 \theta + r^4 \sin^4 \theta) r dr d\theta$
 Then write in terms of $\cos 4\theta$.

9. $\det J = 2$. Integral $\rightarrow \int_{v=0}^1 \int_{u=-\frac{v}{2}}^v \frac{v^2}{2} \cos(uv) du dv$
 $= \dots = \frac{1}{2}(1 - \cos(1))$

10. Calculate $\underline{J} = \frac{\partial \underline{r}}{\partial \lambda} \times \frac{\partial \underline{r}}{\partial s} = (\sin s, -\cos s, \lambda)$

Integral $\rightarrow \int_{s=0}^{2\pi} \int_{\lambda=0}^1 (1 + \lambda^2)^{1/2} d\lambda ds$ Solve using $\lambda = \sinh t$

11. $|\underline{J}| = \left| \frac{\partial \underline{r}}{\partial t} \times \frac{\partial \underline{r}}{\partial \theta} \right| = \dots = b(a + b \cos t)$

Then $\int_S z^2 dS = \int_{\theta=0}^{2\pi} \int_{t=0}^{2\pi} (a + b \cos t) b^3 \sin^2 t dt d\theta$.