



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Eric Li
Assignment title: Coursework 2 Drop Box Spring 25
Submission title: Coursework_2.pdf
File name: Coursework_2.pdf
File size: 146.79K
Page count: 5
Word count: 1,570
Character count: 5,417
Submission date: 08-Mar-2025 10:39PM (UTC+0000)
Submission ID: 252098493

Quantum Mechanics II, Coursework 2

Jiaru (Eric) Li (CID: 02216531)

8 March 2025

1. Recall that the Pauli Y operator corresponds to the Pauli matrix $\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$. This matrix has eigenvalues $1, -1$ which respectively correspond to the (normalised) eigenvectors

$$v_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}, \quad v_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix}.$$

The eigenstates of the Pauli Y operator are therefore

$$|y_+\rangle = \frac{|0\rangle + i|1\rangle}{\sqrt{2}}, \quad |y_-\rangle = \frac{|0\rangle - i|1\rangle}{\sqrt{2}}.$$

We can then express the state $|\psi\rangle = |0\rangle$ in terms of $|y_+\rangle, |y_-\rangle$ as

$$|0\rangle = \frac{1}{\sqrt{2}} (|y_+\rangle + |y_-\rangle),$$

so the probability of measuring $+1$ is the square of the modulus of the coefficient of $|y_+\rangle$, i.e.,

$$P(+1) = \left(\frac{1}{\sqrt{2}} \right)^2 = \frac{1}{2}.$$

2. Recall again that the Pauli matrices for X and Z are $\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Their $+1$ and -1 eigenstates can be easily calculated respectively as

$$|x_+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |x_-\rangle = |0\rangle, \quad |z_+\rangle = \frac{|0\rangle - |1\rangle}{\sqrt{2}}, \quad |z_-\rangle = |1\rangle.$$

We are given the operator $\hat{O} = \cos(\theta)\hat{Z} + \sin(\theta)\hat{X}$. By the properties of \hat{Z} and \hat{X} , we have

$$\hat{Z}|x_+\rangle = |x_+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad \hat{Z}|x_-\rangle = |x_-\rangle = |0\rangle, \quad \hat{X}|x_+\rangle = |x_+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad \hat{X}|x_-\rangle = |z_-\rangle = |1\rangle,$$

so that the expectations are

$$\langle x_+|\hat{Z}|x_+\rangle = 0, \quad \langle z_+|\hat{Z}|z_+\rangle = 1, \quad \langle x_+|\hat{X}|x_+\rangle = 1, \quad \langle z_+|\hat{X}|z_+\rangle = 0.$$

Since \hat{O} is a linear combination of \hat{Z} and \hat{X} , we have

$$\langle x_+|\hat{O}|x_+\rangle = \sin\theta, \quad \langle z_+|\hat{O}|z_+\rangle = \cos\theta.$$

To compute the probability of measuring $+1$, recall that

$$\langle \hat{O} \rangle = (+1)P(+1) + (-1)P(-1) = P(+1) - P(-1), \quad P(+1) + P(-1) = 1$$

from the definition of expectation and probability, so we have $P(+1) = \frac{1 + \langle \hat{O} \rangle}{2}$

1