

Analysis 1A

Lecture 3 - Countability

Ajay Chandra

Countability

Definition

A set S is *countably infinite* if and only if there exists a bijection $f: \mathbb{N}_{>0} \rightarrow S$

A set that is finite, or countably infinite, is said to be countable.

Countability

Definition

A set S is *countably infinite* if and only if there exists a bijection $f: \mathbb{N}_{>0} \rightarrow S$

A set that is finite, or countably infinite, is said to be countable.

Intuition: f lets us write all of S as an infinite list:

$$S = \{s_1, s_2, s_3, s_4, s_5, \dots\}$$

with no repeats (all s_i distinct). Here $s_n := f(n)$.

Countability

Definition

A set S is *countably infinite* if and only if there exists a bijection $f: \mathbb{N}_{>0} \rightarrow S$

A set that is finite, or countably infinite, is said to be countable.

Intuition: f lets us write all of S as an infinite list:

$$S = \{s_1, s_2, s_3, s_4, s_5, \dots\}$$

with no repeats (all s_i distinct). Here $s_n := f(n)$.

Intuition: Given sets A and B , the existence of a bijection $f: A \rightarrow B$ means A and B have the same “number of elements”.

Since the even positive numbers $2\mathbb{N}_{>0} \subset \mathbb{N}_{>0}$, you might think there are less of them, but..

Since the even positive numbers $2\mathbb{N}_{>0} \subset \mathbb{N}_{>0}$, you might think there are less of them, but..

Proposition 2.16

Suppose $S \subset \mathbb{N}$ is infinite. Then S is countably infinite.

Since the even positive numbers $2\mathbb{N}_{>0} \subset \mathbb{N}_{>0}$, you might think there are less of them, but..

Proposition 2.16

Suppose $S \subset \mathbb{N}$ is infinite. Then S is countably infinite.

Proof: We list the elements of S in order of size.

Since the even positive numbers $2\mathbb{N}_{>0} \subset \mathbb{N}_{>0}$, you might think there are less of them, but..

Proposition 2.16

Suppose $S \subset \mathbb{N}$ is infinite. Then S is countably infinite.

Proof: We list the elements of S in order of size.

More precisely, we define $f: \mathbb{N}_{>0} \rightarrow S$ inductively as follows:

- $f(1) = \min S$,

Since the even positive numbers $2\mathbb{N}_{>0} \subset \mathbb{N}_{>0}$, you might think there are less of them, but..

Proposition 2.16

Suppose $S \subset \mathbb{N}$ is infinite. Then S is countably infinite.

Proof: We list the elements of S in order of size.

More precisely, we define $f: \mathbb{N}_{>0} \rightarrow S$ inductively as follows:

- $f(1) = \min S$,
- Assume $f(1), \dots, f(n-1)$ are defined already. Since S is infinite the set $S \setminus \{f(1), \dots, f(n-1)\}$ is nonempty and so we may define

$$f(n) := \min \left(S \setminus \{f(1), \dots, f(n-1)\} \right).$$

Finishing the proof of Prop. 2.16

- Need to check f is injective and surjective.

Finishing the proof of Prop. 2.16

- Need to check f is injective and surjective.
- To show f is injective, note that for distinct $i, j \in \mathbb{N}_{>0}$ we have either $i < j$ or $j < i$ and therefore either $f(i) < f(j)$ or $f(j) < f(i)$.

Finishing the proof of Prop. 2.16

- Need to check f is injective and surjective.
- To show f is injective, note that for distinct $i, j \in \mathbb{N}_{>0}$ we have either $i < j$ or $j < i$ and therefore either $f(i) < f(j)$ or $f(j) < f(i)$.
- To show f is surjective, we argue by contradiction.

Finishing the proof of Prop. 2.16

- Need to check f is injective and surjective.
- To show f is injective, note that for distinct $i, j \in \mathbb{N}_{>0}$ we have either $i < j$ or $j < i$ and therefore either $f(i) < f(j)$ or $f(j) < f(i)$.
- To show f is surjective, we argue by contradiction. If f isn't surjective, then \exists smallest $s \in S \setminus \text{im}(f)$.

Finishing the proof of Prop. 2.16

- Need to check f is injective and surjective.
- To show f is injective, note that for distinct $i, j \in \mathbb{N}_{>0}$ we have either $i < j$ or $j < i$ and therefore either $f(i) < f(j)$ or $f(j) < f(i)$.
- To show f is surjective, we argue by contradiction. If f isn't surjective, then \exists smallest $s \in S \setminus \text{im}(f)$. Since $s \neq \min S$ (because $f(1) = \min S$) we know $\exists s' \in S$ such that $s' < s$

Finishing the proof of Prop. 2.16

- Need to check f is injective and surjective.
- To show f is injective, note that for distinct $i, j \in \mathbb{N}_{>0}$ we have either $i < j$ or $j < i$ and therefore either $f(i) < f(j)$ or $f(j) < f(i)$.
- To show f is surjective, we argue by contradiction. If f isn't surjective, then \exists smallest $s \in S \setminus \text{im}(f)$. Since $s \neq \min S$ (because $f(1) = \min S$) we know $\exists s' \in S$ such that $s' < s$ – picking the largest such s' , then $s' = f(n)$ for some $n \in \mathbb{N}_{>0}$ and by our rule we must have $s = f(n+1)$.



Proposition 2.17

\mathbb{Z} is countably infinite.

Proposition 2.17


\mathbb{Z} is countably infinite.

Proof We list them as $\mathbb{Z} = \{0, 1, -1, 2, -2, 3, -3, \dots\}$.

Proposition 2.17

\mathbb{Z} is countably infinite.


Proof We list them as $\mathbb{Z} = \{0, 1, -1, 2, -2, 3, -3, \dots\}$.



Proposition 2.17

\mathbb{Z} is countably infinite.

Proof We list them as $\mathbb{Z} = \{0, 1, -1, 2, -2, 3, -3, \dots\}$.




Formally, define a bijection $f : \mathbb{N}_{>0} \rightarrow \mathbb{Z}$ by declaring, for $k \geq 1$,

$$\begin{cases} f(2k-1) &:= -(k-1), \\ f(2k) &:= k. \end{cases} \quad \square$$

Proposition 2.17

\mathbb{Z} is countably infinite.

Proof We list them as $\mathbb{Z} = \{0, 1, -1, 2, -2, 3, -3, \dots\}$.



Formally, define a bijection $f : \mathbb{N}_{>0} \rightarrow \mathbb{Z}$ by declaring, for $k \geq 1$,

$$\begin{cases} f(2k-1) &:= -(k-1), \\ f(2k) &:= k. \end{cases} \quad \square$$


Exercise 2.18

Carefully show that f as above is indeed bijective.

Proposition 2.17

\mathbb{Z} is countably infinite.

Proof We list them as $\mathbb{Z} = \{0, 1, -1, 2, -2, 3, -3, \dots\}$.



Formally, define a bijection $f : \mathbb{N}_{>0} \rightarrow \mathbb{Z}$ by declaring, for $k \geq 1$,

$$\begin{cases} f(2k-1) &:= -(k-1), \\ f(2k) &:= k. \end{cases} \quad \square$$

Exercise 2.18

Carefully show that f as above is indeed bijective.

Show similarly that A, B countable $\Rightarrow A \cup B$ countable.

Theorem 2.19

\mathbb{Q} is countably infinite.

Theorem 2.19

\mathbb{Q} is countably infinite.

Proof: First let's show $\mathbb{Q}_{>0} = \{x \in \mathbb{Q} : x > 0\}$ is countably infinite.

Theorem 2.19


\mathbb{Q} is countably infinite.

Proof: First let's show $\mathbb{Q}_{>0} = \{x \in \mathbb{Q} : x > 0\}$ is countably infinite.

Here is one proof “by picture” where we show one way to “list” the elements of $\mathbb{Q}_{>0}$


For this proof by picture we arrange the pairs $(p, q) \in \mathbb{N}_{>0}^2$ in a square:

For this proof by picture we arrange the pairs $(p, q) \in \mathbb{N}_{\geq 0}^2$ in a square:



Now list the pairs according to the path shown, *missing out pairs which aren't in lowest terms*.

For this proof by picture we arrange the pairs $(p, q) \in \mathbb{N}_{\geq 0}^2$ in a square:



Now list the pairs according to the path shown, *missing out pairs which aren't in lowest terms*. It is not clear how to write an explicit formula for this list....

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite.

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite. Define

$$f: \mathbb{Q}_{>0} \rightarrow \mathbb{N}_{>0}, \quad f(m/n) := 2^m 3^n$$

where $m, n \geq 1$ and m/n is in lowest terms. Then we can show

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite. Define

$$f: \mathbb{Q}_{>0} \rightarrow \mathbb{N}_{>0}, \quad f(m/n) := 2^m 3^n$$

where $m, n \geq 1$ and m/n is in lowest terms. Then we can show

- f is injective.

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite. Define

$$f: \mathbb{Q}_{>0} \rightarrow \mathbb{N}_{>0}, \quad f(m/n) := 2^m 3^n$$

where $m, n \geq 1$ and m/n is in lowest terms. Then we can show

- f is injective.
- $\text{im}(f)$ is an infinite subset of \mathbb{N} and therefore a countably infinite set by Proposition 2.16.

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite. Define

$$f: \mathbb{Q}_{>0} \rightarrow \mathbb{N}_{>0}, \quad f(m/n) := 2^m 3^n$$

where $m, n \geq 1$ and m/n is in lowest terms. Then we can show

- f is injective.
- $\text{im}(f)$ is an infinite subset of \mathbb{N} and therefore a countably infinite set by Proposition 2.16.
- Therefore we have a bijection $F: \mathbb{N}_{>0} \rightarrow \mathbb{Q}_{>0}$.

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite. Define

$$f: \mathbb{Q}_{>0} \rightarrow \mathbb{N}_{>0}, \quad f(m/n) := 2^m 3^n$$

where $m, n \geq 1$ and m/n is in lowest terms. Then we can show

- f is injective.
- $\text{im}(f)$ is an infinite subset of \mathbb{N} and therefore a countably infinite set by Proposition 2.16.
- Therefore we have a bijection $F: \mathbb{N}_{>0} \rightarrow \mathbb{Q}_{>0}$.

Now we show the countability of \mathbb{Q} - define a bijection $g: \mathbb{N}_{>0} \rightarrow \mathbb{Q}$ we then set

$$g(1) := 0 \quad \text{and} \quad \begin{cases} g(2k) & := F(k), \\ g(2k+1) & := -F(k). \end{cases}$$

Here is a more explicit argument to show that $\mathbb{Q}_{>0}$ is countably infinite. Define

$$f: \mathbb{Q}_{>0} \rightarrow \mathbb{N}_{>0}, \quad f(m/n) := 2^m 3^n$$

where $m, n \geq 1$ and m/n is in lowest terms. Then we can show

- f is injective.
- $\text{im}(f)$ is an infinite subset of \mathbb{N} and therefore a countably infinite set by Proposition 2.16.
- Therefore we have a bijection $F: \mathbb{N}_{>0} \rightarrow \mathbb{Q}_{>0}$.

Now we show the countability of \mathbb{Q} - define a bijection $g: \mathbb{N}_{>0} \rightarrow \mathbb{Q}$ we then set

$$g(1) := 0 \quad \text{and} \quad \begin{cases} g(2k) &:= F(k), \\ g(2k+1) &:= -F(k). \end{cases}$$

That is, if q_1, q_2, \dots ($q_i := F(i)$) is our list of elements of $\mathbb{Q}_{>0}$ then our new list is $0, q_1, -q_1, q_2, -q_2, \dots$



Theorem 2.21

\mathbb{R} is uncountable.

Proof: (Cantor's Diagonal Argument) We argue by contradiction, suppose that you can “list” all the real numbers.

We write this list as follows, using decimal expansions with no $\overline{9}$ s:

$$x_1 = a_1. a_{11} a_{12} a_{13} a_{14} \dots$$

$$x_2 = a_2. a_{21} a_{22} a_{23} a_{24} \dots$$

$$x_3 = a_3. a_{31} a_{32} a_{33} a_{34} \dots$$

$$x_4 = a_4. a_{41} a_{42} a_{43} a_{44} \dots$$

$$\vdots$$

$$x_m = a_m. a_{m1} a_{m2} a_{m3} a_{m4} \dots$$

$$\vdots$$

We write this list as follows, using decimal expansions with no $\overline{9}$ s:

$$x_1 = a_1. a_{11} a_{12} a_{13} a_{14} \dots$$

$$x_2 = a_2. a_{21} a_{22} a_{23} a_{24} \dots$$

$$x_3 = a_3. a_{31} a_{32} a_{33} a_{34} \dots$$

$$x_4 = a_4. a_{41} a_{42} a_{43} a_{44} \dots$$

$$\vdots$$

$$x_m = a_m. a_{m1} a_{m2} a_{m3} a_{m4} \dots$$

$$\vdots$$

As usual $a_1, a_2, a_3, \dots \in \mathbb{Z}$ and $a_{11}, a_{12}, \dots \in \{0, 1, 2, \dots, 9\}$.

Now we produce a real number $x := 0.b_1 b_2 \dots b_n \dots$ not on the list:

Now we produce a real number $x := 0.b_1 b_2 \dots b_n \dots$ not on the list:

- 1 Pick $b_1 \in \{0, 1, \dots, 8\}$ such that $b_1 \neq a_{11}$,
- 2 Pick $b_2 \in \{0, 1, \dots, 8\}$ such that $b_2 \neq a_{22}$,
- \vdots
- n . Pick $b_n \in \{0, 1, \dots, 8\}$ such that $b_n \neq a_{nn}$
- \vdots

Now we produce a real number $x := 0. b_1 b_2 \dots b_n \dots$ not on the list:

1 Pick $b_1 \in \{0, 1, \dots, 8\}$ such that $b_1 \neq a_{11}$,

2 Pick $b_2 \in \{0, 1, \dots, 8\}$ such that $b_2 \neq a_{22}$,

\vdots

n . Pick $b_n \in \{0, 1, \dots, 8\}$ such that $b_n \neq a_{nn}$

\vdots

- Since we don't allow 9 we don't end up with a decimal ending in $\overline{9}$ and we really have $x \in \mathbb{R}$

Now we produce a real number $x := 0. b_1 b_2 \dots b_n \dots$ not on the list:

- 1 Pick $b_1 \in \{0, 1, \dots, 8\}$ such that $b_1 \neq a_{11}$,
- 2 Pick $b_2 \in \{0, 1, \dots, 8\}$ such that $b_2 \neq a_{22}$,
- \vdots
- n . Pick $b_n \in \{0, 1, \dots, 8\}$ such that $b_n \neq a_{nn}$
- \vdots

- Since we don't allow 9 we don't end up with a decimal ending in $\bar{9}$ and we really have $x \in \mathbb{R}$
- Then $\forall i \geq 1$, we see $x \neq x_i$ because it differs in the i th decimal place a_{ii} .

Now we produce a real number $x := 0. b_1 b_2 \dots b_n \dots$ not on the list:

- 1 Pick $b_1 \in \{0, 1, \dots, 8\}$ such that $b_1 \neq a_{11}$,
- 2 Pick $b_2 \in \{0, 1, \dots, 8\}$ such that $b_2 \neq a_{22}$,
- \vdots
- n . Pick $b_n \in \{0, 1, \dots, 8\}$ such that $b_n \neq a_{nn}$
- \vdots

- Since we don't allow 9 we don't end up with a decimal ending in $\overline{9}$ and we really have $x \in \mathbb{R}$
- Then $\forall i \geq 1$, we see $x \neq x_i$ because it differs in the i th decimal place a_{ii} .

Therefore we have found an $x \in \mathbb{R}$ not on the list.



- There's a set \mathbb{A} with $\mathbb{Q} \subset \mathbb{A} \subset \mathbb{R}$ called the set of *algebraic numbers*: \mathbb{A} is the collection of $x \in \mathbb{R}$ which satisfy a polynomial equation $p(x) = 0$, where p has integer coefficients.

- There's a set \mathbb{A} with $\mathbb{Q} \subset \mathbb{A} \subset \mathbb{R}$ called the set of *algebraic numbers*: \mathbb{A} is the collection of $x \in \mathbb{R}$ which satisfy a polynomial equation $p(x) = 0$, where p has integer coefficients.
- Any rational number p/q satisfies an equation $p(x) := qx - p = 0$, so we indeed have $\mathbb{Q} \subset \mathbb{A}$
- $\sqrt[n]{m}$ satisfies $p(x) := x^n - m = 0$, so $\sqrt[n]{m} \in \mathbb{A}$.

- There's a set \mathbb{A} with $\mathbb{Q} \subset \mathbb{A} \subset \mathbb{R}$ called the set of *algebraic numbers*: \mathbb{A} is the collection of $x \in \mathbb{R}$ which satisfy a polynomial equation $p(x) = 0$, where p has integer coefficients.
- Any rational number p/q satisfies an equation $p(x) := qx - p = 0$, so we indeed have $\mathbb{Q} \subset \mathbb{A}$
- $\sqrt[n]{m}$ satisfies $p(x) := x^n - m = 0$, so $\sqrt[n]{m} \in \mathbb{A}$.
- On Problem Sheet 1 you'll prove that the set of algebraic numbers is also *countable*.

- There's a set \mathbb{A} with $\mathbb{Q} \subset \mathbb{A} \subset \mathbb{R}$ called the set of *algebraic numbers*: \mathbb{A} is the collection of $x \in \mathbb{R}$ which satisfy a polynomial equation $p(x) = 0$, where p has integer coefficients.
- Any rational number p/q satisfies an equation $p(x) := qx - p = 0$, so we indeed have $\mathbb{Q} \subset \mathbb{A}$
- $\sqrt[n]{m}$ satisfies $p(x) := x^n - m = 0$, so $\sqrt[n]{m} \in \mathbb{A}$.
- On Problem Sheet 1 you'll prove that the set of algebraic numbers is also *countable*.
- Therefore (by Exercise 2.18) the set of *transcendental numbers* – real numbers which are not algebraic – is uncountable.

- There's a set \mathbb{A} with $\mathbb{Q} \subset \mathbb{A} \subset \mathbb{R}$ called the set of *algebraic numbers*: \mathbb{A} is the collection of $x \in \mathbb{R}$ which satisfy a polynomial equation $p(x) = 0$, where p has integer coefficients.
- Any rational number p/q satisfies an equation $p(x) := qx - p = 0$, so we indeed have $\mathbb{Q} \subset \mathbb{A}$
- $\sqrt[n]{m}$ satisfies $p(x) := x^n - m = 0$, so $\sqrt[n]{m} \in \mathbb{A}$.
- On Problem Sheet 1 you'll prove that the set of algebraic numbers is also *countable*.
- Therefore (by Exercise 2.18) the set of *transcendental numbers* – real numbers which are not algebraic – is uncountable.
- Examples of transcendental numbers are e and π

- There's a set \mathbb{A} with $\mathbb{Q} \subset \mathbb{A} \subset \mathbb{R}$ called the set of *algebraic numbers*: \mathbb{A} is the collection of $x \in \mathbb{R}$ which satisfy a polynomial equation $p(x) = 0$, where p has integer coefficients.
- Any rational number p/q satisfies an equation $p(x) := qx - p = 0$, so we indeed have $\mathbb{Q} \subset \mathbb{A}$
- $\sqrt[n]{m}$ satisfies $p(x) := x^n - m = 0$, so $\sqrt[n]{m} \in \mathbb{A}$.
- On Problem Sheet 1 you'll prove that the set of algebraic numbers is also *countable*.
- Therefore (by Exercise 2.18) the set of *transcendental numbers* – real numbers which are not algebraic – is uncountable.
- Examples of transcendental numbers are e and π (proving e or π is transcendental is harder than proving $\sqrt{3}$ is irrational....).