

# Quantum Mechanics II, Coursework 1

Jiaru (Eric) Li (CID: 02216531)

17 February 2025

## Question 1

(a) We are given that  $\phi(x) = Axe^{-x^2/a^2}$ , which must satisfy the normalisation condition

$$\int_{-\infty}^{\infty} |\phi(x)|^2 dx = 1.$$

We have  $|\phi(x)|^2 = A^2 x^2 e^{-2x^2/a^2}$ , and by setting  $\alpha = 2/a^2$ , we get

$$\begin{aligned} \int_{-\infty}^{\infty} A^2 x^2 e^{-2x^2/a^2} dx &= A^2 \int_{-\infty}^{\infty} x^2 e^{-\alpha x^2} dx \\ &= -A^2 \int_{-\infty}^{\infty} \frac{d}{d\alpha} (e^{-\alpha x^2}) dx \\ &= -A^2 \frac{d}{d\alpha} \int_{-\infty}^{\infty} e^{-\alpha x^2} dx, \end{aligned}$$

where we recall the Gaussian integral

$$\int_{-\infty}^{\infty} e^{-\alpha x^2} dx = \sqrt{\frac{\pi}{\alpha}}.$$

This gives

$$\frac{d}{d\alpha} \int_{-\infty}^{\infty} e^{-\alpha x^2} dx = -\frac{\sqrt{\pi}}{2} \alpha^{-3/2} = -\frac{a^3 \sqrt{\pi}}{2^{5/2}}.$$

Substituting into the normalisation condition gives

$$\begin{aligned} (-A^2) \left( -\frac{a^3 \sqrt{\pi}}{2^{5/2}} \right) &= 1 \\ A^2 &= \frac{2^{5/2}}{a^3 \sqrt{\pi}}, \end{aligned}$$

so we take  $A$  to be

$$A = \frac{2^{5/4}}{a^{3/2} \pi^{1/4}}.$$

(b) To compute  $\tilde{\phi}(p)$  in the momentum basis, we use the Fourier transform relation, namely

$$\tilde{\phi}(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(x) e^{-ixp/\hbar} dx.$$

Recall the standard definition of Fourier transform in the form of

$$\mathcal{F}\{f(x)\}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ikx} dx.$$

Therefore, by setting  $k = p/\hbar$ , we get

$$\tilde{\phi}(p) = \frac{1}{\sqrt{\hbar}} \mathcal{F}\{\phi(x)\}(k).$$

To compute the Fourier transform of  $\phi(x) = Axe^{-x^2/a^2}$ , we recall a property of Fourier transform

$$\mathcal{F}\{xf(x)\}(k) = i \frac{d}{dk} (\mathcal{F}\{f(x)\}(k)),$$

and we already know the Fourier transform of a Gaussian function  $f(x) = e^{-\alpha x^2}$  as

$$\mathcal{F}\{f(x)\}(k) = \sqrt{\frac{1}{2\alpha}} e^{-k^2/4\alpha}.$$

We then have

$$\begin{aligned} \mathcal{F}\{\phi(x)\}(k) &= iA \frac{d}{dk} \left( \mathcal{F}\left\{e^{-x^2/a^2}\right\}(k) \right) \\ &= iA \frac{d}{dk} \left( \frac{a}{\sqrt{2}} e^{-a^2 k^2/4} \right) \\ &= iA \frac{a}{\sqrt{2}} e^{-a^2 k^2/4} \left( -\frac{a^2 k}{2} \right), \end{aligned}$$

so by plugging  $A$  from part (a) in, we get

$$\begin{aligned} \mathcal{F}\{\phi(x)\}(k) &= -i \frac{2^{5/4}}{a^{3/2} \pi^{1/4}} \frac{a}{\sqrt{2}} \frac{a^2}{2} \cdot k e^{-a^2 k^2/4} \\ &= -i \frac{a^{3/2}}{(2\pi)^{1/4}} k e^{-a^2 k^2/4} \\ &= -i \frac{a^{3/2}}{\hbar (2\pi)^{1/4}} p e^{-a^2 p^2/4\hbar^2}, \end{aligned}$$

which gives the final answer

$$\boxed{\tilde{\phi}(p) = -i \frac{(a/\hbar)^{3/2}}{(2\pi)^{1/4}} p e^{-a^2 p^2/4\hbar^2}}.$$

(c) Again recall the Fourier transform relation between  $\phi(x)$  and  $\tilde{\phi}(p)$  given as

$$\tilde{\phi}(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(x) e^{-ixp/\hbar} dx.$$

Therefore, we have

$$\tilde{\phi}(-p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(x) e^{ixp/\hbar} dx,$$

so if we substitute  $x = -u$  into the integral and use the identity  $\phi(x) = \phi(-x)$  given, we get

$$\begin{aligned} \tilde{\phi}(-p) &= \frac{1}{\sqrt{2\pi\hbar}} \int_{\infty}^{-\infty} \phi(-u) e^{-iup/\hbar} (-du) \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(u) e^{-iup/\hbar} du \\ &= \tilde{\phi}(p), \end{aligned}$$

which implies that  $\boxed{\tilde{\phi}(p)}$  is also even.

- (d) We are now given the condition  $\phi(x) = [\phi(-x)]^*$  where  $a^*$  denotes the complex conjugate of  $a$ . Recall again that

$$\tilde{\phi}(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(x) e^{-ixp/\hbar} dx.$$

The complex conjugate of  $\tilde{\phi}(p)$  is given by

$$\begin{aligned} [\tilde{\phi}(p)]^* &= \left[ \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(x) e^{-ixp/\hbar} dx \right]^* \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} [\phi(x) e^{-ixp/\hbar}]^* dx \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} [\phi(x)]^* e^{ixp/\hbar} dx, \end{aligned}$$

where we have used the identity  $(e^z)^* = e^{z^*}$ .

We can then substitute  $x = -u$  into the integral, giving

$$\begin{aligned} [\tilde{\phi}(p)]^* &= \frac{1}{\sqrt{2\pi\hbar}} \int_{\infty}^{-\infty} [\phi(-u)]^* e^{-iup/\hbar} (-du) \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \phi(u) e^{-iup/\hbar} du, \end{aligned}$$

which is precisely the definition of  $\tilde{\phi}(p)$ .

This implies that the complex conjugate of  $\tilde{\phi}(p)$  equals to itself, i.e.,  $\tilde{\phi}(p)$  is real.

## Question 2

- (a) The rotation operator  $\hat{R}$  is given by  $\hat{R} = |3\rangle\langle 1| + |1\rangle\langle 2| + |2\rangle\langle 3|$ , so its Hermitian conjugate  $\hat{R}^\dagger$  is  $\hat{R}^\dagger = |1\rangle\langle 3| + |2\rangle\langle 1| + |3\rangle\langle 2|$ .

By using the basic fact that  $\langle a|b\rangle = 1$  when  $a = b$  and 0 otherwise, we can then verify that

$$\begin{aligned} \hat{R}^\dagger \hat{R} &= (|3\rangle\langle 1| + |1\rangle\langle 2| + |2\rangle\langle 3|) \cdot (|1\rangle\langle 3| + |2\rangle\langle 1| + |3\rangle\langle 2|) \\ &= |3\rangle\langle 1|1\rangle\langle 3| + |3\rangle\langle 1|2\rangle\langle 1| + |3\rangle\langle 1|3\rangle\langle 2| + \\ &\quad |1\rangle\langle 2|1\rangle\langle 3| + |1\rangle\langle 2|2\rangle\langle 1| + |1\rangle\langle 2|3\rangle\langle 2| + \\ &\quad |2\rangle\langle 3|1\rangle\langle 3| + |2\rangle\langle 3|2\rangle\langle 1| + |2\rangle\langle 3|3\rangle\langle 2| \\ &= |3\rangle\langle 3| + |1\rangle\langle 1| + |2\rangle\langle 2| = I. \end{aligned}$$

Similarly, the reflection operator  $\hat{\sigma} = |1\rangle\langle 2| + |2\rangle\langle 1| + |3\rangle\langle 3|$  has Hermitian conjugate  $\hat{\sigma}^\dagger = |2\rangle\langle 1| + |1\rangle\langle 2| + |3\rangle\langle 3|$  and satisfies

$$\begin{aligned} \hat{\sigma}^\dagger \hat{\sigma} &= (|1\rangle\langle 2| + |2\rangle\langle 1| + |3\rangle\langle 3|)(|2\rangle\langle 1| + |1\rangle\langle 2| + |3\rangle\langle 3|) \\ &= |1\rangle\langle 2|2\rangle\langle 1| + |1\rangle\langle 2|1\rangle\langle 2| + |1\rangle\langle 2|3\rangle\langle 3| + \\ &\quad |2\rangle\langle 1|2\rangle\langle 1| + |2\rangle\langle 1|1\rangle\langle 2| + |2\rangle\langle 1|3\rangle\langle 3| + \\ &\quad |3\rangle\langle 3|2\rangle\langle 1| + |3\rangle\langle 3|1\rangle\langle 2| + |3\rangle\langle 3|3\rangle\langle 3| \\ &= |1\rangle\langle 1| + |2\rangle\langle 2| + |3\rangle\langle 3| = I. \end{aligned}$$

We have therefore shown that  $\hat{R}, \hat{\sigma}$  are unitary.

To prove commutativity with the Hamiltonian  $\hat{\mathcal{H}}$  given, recall that

$$\hat{\mathcal{H}} = -w[|1\rangle\langle 2| + |2\rangle\langle 3| + |3\rangle\langle 1| + \text{h.c.}].$$

Note that we have

$$\begin{aligned}\hat{R}|1\rangle &= (|3\rangle\langle 1| + |1\rangle\langle 2| + |2\rangle\langle 3|)|1\rangle = |3\rangle, \\ \hat{R}|2\rangle &= (|3\rangle\langle 1| + |1\rangle\langle 2| + |2\rangle\langle 3|)|2\rangle = |1\rangle, \\ \hat{R}|3\rangle &= (|3\rangle\langle 1| + |1\rangle\langle 2| + |2\rangle\langle 3|)|3\rangle = |2\rangle,\end{aligned}$$

so  $\hat{R}$  is essentially a cyclic permutation that maps  $\{1, 2, 3\}$  to  $\{3, 1, 2\}$ , leaving  $\hat{\mathcal{H}}$  unchanged.

Intuitively, if we think of  $\hat{\mathcal{H}}$  as describing a particle that can jump between vertices of an equilateral triangle, as given in the problem, then  $\hat{R}$  is a  $120^\circ$  rotation.

Similarly, we can also write

$$\begin{aligned}\hat{\sigma}|1\rangle &= (|1\rangle\langle 2| + |2\rangle\langle 1| + |3\rangle\langle 3|)|1\rangle = |2\rangle, \\ \hat{\sigma}|2\rangle &= (|1\rangle\langle 2| + |2\rangle\langle 1| + |3\rangle\langle 3|)|2\rangle = |1\rangle, \\ \hat{\sigma}|3\rangle &= (|1\rangle\langle 2| + |2\rangle\langle 1| + |3\rangle\langle 3|)|3\rangle = |3\rangle,\end{aligned}$$

so  $\hat{\sigma}$  only swaps 1 and 2, which again preserves the symmetry of  $\hat{\mathcal{H}}$ .

Intuitively,  $\hat{\sigma}$  is a reflection with the axis being the median from the vertex with label 3.

We have now proven that  $\boxed{\hat{R}, \hat{\sigma} \text{ commutes with } \hat{\mathcal{H}}}$ .

From the geometric picture given, and by looking at the structure of the Hamiltonian, we could find another two distinct unitary symmetry operators, which are basically the other two reflections,

$$\begin{aligned}\hat{\sigma}_1 &= |2\rangle\langle 3| + |3\rangle\langle 2| + |1\rangle\langle 1|, \\ \hat{\sigma}_2 &= |1\rangle\langle 3| + |3\rangle\langle 1| + |2\rangle\langle 2|,\end{aligned}$$

in addition to the (trivial) identity operator.

*(In fact, in the language of group theory, these symmetries form a dihedral group of order 6,  $D_6$ . This group has 6 elements: one identity; two  $120^\circ$  rotations, clockwise and anticlockwise (corresponding to  $\hat{R}$  and  $\hat{R}^\dagger$ ); and three reflections. The fact that  $D_6$  is isomorphic to the symmetric group of order 3,  $S_3$ , tells us about the symmetry of the Hamiltonian in an elegant way.)*

- (b) From part (a), we can write  $\hat{R}$  and  $\hat{\sigma}$  in matrix form with respect to the given basis  $\{|1\rangle, |2\rangle, |3\rangle\}$  as permutation matrices,

$$\hat{R} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad \hat{\sigma} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

To find the eigenvalues of  $\hat{R}$ , we compute

$$\begin{aligned}\det(\hat{R} - \lambda I) &= \begin{vmatrix} -\lambda & 1 & 0 \\ 0 & -\lambda & 1 \\ 1 & 0 & -\lambda \end{vmatrix} \\ &= -\lambda \begin{vmatrix} -\lambda & 1 \\ 0 & -\lambda \end{vmatrix} - \begin{vmatrix} 0 & 1 \\ 1 & -\lambda \end{vmatrix} \\ &= -\lambda^3 + 1,\end{aligned}$$

whose roots are the third roots of unity  $\boxed{1, \omega, \omega^2}$  where  $\omega = e^{2\pi i/3}$ . These are the eigenvalues required.

The corresponding eigenvectors  $v_i$  for  $i = 1, 2, 3$  must satisfy  $\hat{R}v_i = \lambda v_i$ , and so by letting  $v_i = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$ ,

we get  $\hat{R}v_i = \begin{pmatrix} c \\ a \\ b \end{pmatrix}$  and  $\lambda v_i = \begin{pmatrix} \lambda a \\ \lambda b \\ \lambda c \end{pmatrix}$ , and we have

$$\begin{cases} c = \lambda a, \\ a = \lambda b, \\ b = \lambda c. \end{cases}$$

For  $\lambda = 1$ , one possible solution is  $(a, b, c) = (1, 1, 1)$ . For  $\lambda = \omega$ , one possible solution is  $(a, b, c) = (1, \omega, \omega^2)$ . For  $\lambda = \omega^2$ , one possible solution is  $(a, b, c) = (1, \omega^2, \omega)$ .

The eigenstates of  $\hat{R}$  can therefore be expressed as

$$\begin{aligned} |\phi_1\rangle &= |1\rangle + |2\rangle + |3\rangle, \\ |\phi_2\rangle &= |1\rangle + \omega |2\rangle + \omega^2 |3\rangle, \\ |\phi_3\rangle &= |1\rangle + \omega^2 |2\rangle + \omega |3\rangle, \end{aligned}$$

so an eigenbasis is given by  $\{|\phi_1\rangle, |\phi_2\rangle, |\phi_3\rangle\}$ .

Now we return to the Hamiltonian given,

$$\hat{\mathcal{H}} = -w[|1\rangle\langle 2| + |2\rangle\langle 3| + |3\rangle\langle 1| + \text{h.c.}].$$

We observe that this can alternatively be written as  $\hat{\mathcal{H}} = -w(\hat{R} + \hat{R}^\dagger) = -w(\hat{R} + \hat{R}^2)$ .

Therefore, we have

$$\begin{aligned} \hat{\mathcal{H}}|\phi_i\rangle &= -w(\hat{R} + \hat{R}^2)|\phi_i\rangle \\ &= -w(\lambda + \lambda^2)|\phi_i\rangle. \end{aligned}$$

Since  $\lambda + \lambda^2$  can take values of  $1 + 1^2 = 2, \omega + \omega^2 = -1$  and  $\omega^2 + (\omega^2)^2 = -1$ , the three eigenvalues of  $\hat{\mathcal{H}}$  are given by  $-2w, w, w$  under the same eigenbasis.

We then return our attention to  $\hat{\sigma} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . This matrix has eigenvalues  $\mu$  such that

$$\begin{aligned} \det(\hat{\sigma} - \mu I) &= \begin{vmatrix} -\mu & 1 & 0 \\ 1 & -\mu & 0 \\ 0 & 0 & 1 - \mu \end{vmatrix} \\ &= -\mu \begin{vmatrix} -\mu & 0 \\ 0 & 1 - \mu \end{vmatrix} - \begin{vmatrix} 1 & 0 \\ 0 & 1 - \mu \end{vmatrix} \\ &= \mu^2(1 - \mu) - (1 - \mu) \\ &= -(1 - \mu)^2(1 + \mu), \end{aligned}$$

which has roots  $\mu = 1, 1, -1$ . These correspond to eigenvectors  $u_i = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$ , so  $\hat{\sigma}u_i = \begin{pmatrix} b \\ a \\ c \end{pmatrix}$  and

$\mu u_i = \begin{pmatrix} \mu a \\ \mu b \\ \mu c \end{pmatrix}$ , and we have

$$\begin{cases} b = \mu a, \\ a = \mu b, \\ c = \mu c. \end{cases}$$

For  $\mu = 1$ , we have  $a = b$  and  $c$  is arbitrary. For  $\mu = -1$ , we need  $a = -b$  and  $c = 0$ . Therefore, the eigenstates of  $\hat{\sigma}$  takes the form of  $|\psi_1\rangle = |1\rangle - |2\rangle$  and  $|\psi_{2,3}\rangle = a(|1\rangle + |2\rangle) + b|3\rangle$ , where  $a$  and  $b$  are constants.

We now compute

$$\begin{aligned} \hat{\mathcal{H}}|\psi_{2,3}\rangle &= -w[|1\rangle\langle 2| + |2\rangle\langle 3| + |3\rangle\langle 1| + \text{h.c.}][a(|1\rangle + |2\rangle) + b|3\rangle] \\ &= -w(a|1\rangle\langle 2|1\rangle + a|1\rangle\langle 2|2\rangle + b|1\rangle\langle 2|3\rangle + a|2\rangle\langle 3|1\rangle + a|2\rangle\langle 3|2\rangle + b|2\rangle\langle 3|3\rangle + \\ &\quad a|3\rangle\langle 1|1\rangle + a|3\rangle\langle 1|2\rangle + b|3\rangle\langle 1|3\rangle + a|2\rangle\langle 1|1\rangle + a|2\rangle\langle 1|2\rangle + b|2\rangle\langle 1|3\rangle + \\ &\quad a|3\rangle\langle 2|1\rangle + a|3\rangle\langle 2|2\rangle + b|3\rangle\langle 2|3\rangle + a|1\rangle\langle 3|1\rangle + a|1\rangle\langle 3|2\rangle + b|1\rangle\langle 3|3\rangle) \\ &= -w((a+b)|1\rangle + (a+b)|2\rangle + 2a|3\rangle). \end{aligned}$$

Within the same eigenbasis, we have  $\hat{\mathcal{H}}|\psi_{2,3}\rangle = \lambda|\psi_{2,3}\rangle$ . By comparison, we get  $\lambda a = -w(a+b)$  and  $\lambda b = -2aw$ . Recall that the three eigenvalues of  $\hat{\mathcal{H}}$  are given by  $-2w, w, w$ . For  $\lambda = -2w$ , the solutions are  $b = a$ ; for  $\lambda = w$ , the solutions are  $b = -2a$ .

The eigenstates of  $\hat{\sigma}$  can therefore be expressed as

$$\begin{aligned} |\psi_1\rangle &= |1\rangle - |2\rangle, \\ |\psi_2\rangle &= |1\rangle + |2\rangle + |3\rangle, \\ |\psi_3\rangle &= |1\rangle + |2\rangle - 2|3\rangle, \end{aligned}$$

so an eigenbasis is given by  $\{|\psi_1\rangle, |\psi_2\rangle, |\psi_3\rangle\}$ .

(c) We have already calculated that

$$\hat{R} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad \hat{\sigma} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

so we could compute

$$\hat{R}\hat{\sigma} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \hat{\sigma}\hat{R} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

and therefore

$$[\hat{R}, \hat{\sigma}] = \hat{R}\hat{\sigma} - \hat{\sigma}\hat{R} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{pmatrix}.$$

Note that  $|\phi\rangle = |1\rangle + |2\rangle + |3\rangle$  is a common eigenstate for both  $\hat{R}, \hat{\mathcal{H}}$  and  $\hat{\sigma}, \hat{\mathcal{H}}$ . We have

$$\begin{pmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix},$$

So we have  $[\hat{R}, \hat{\sigma}]|\phi\rangle = 0$ . This is consistent with Section 3.2 of the notes: while  $[\hat{R}, \hat{\sigma}] \neq 0$ , we do have  $[\hat{R}, \hat{\sigma}]|\phi\rangle = 0$  for the non-degenerate eigenstate  $|\phi\rangle$ , verifying the relation between symmetry and degeneracy.

### Question 3

(a) We start with the 1d harmonic oscillator Hamiltonian

$$\hat{\mathcal{H}} = \hbar\omega \left( \hat{a}^\dagger \hat{a} + \frac{1}{2} \right),$$

where the ladder operators  $\hat{a}^\dagger$  and  $\hat{a}$  satisfy  $[\hat{a}, \hat{a}^\dagger] = 1$ .

Therefore, we have

$$[\hat{a}, \hat{\mathcal{H}}] = \hbar\omega [\hat{a}, \hat{a}^\dagger \hat{a}] = \hbar\omega ([\hat{a}, \hat{a}^\dagger] \hat{a} + \hat{a}^\dagger [\hat{a}, \hat{a}]) = \hbar\omega \hat{a}.$$

In the Heisenberg picture, the equation of motion for the ladder operator  $\hat{a}_H$  is given by

$$i\hbar \frac{d}{dt} \hat{a}_H(t) = \hat{\mathcal{U}}^\dagger [\hat{a}, \hat{\mathcal{H}}] \hat{\mathcal{U}} = \hbar\omega \hat{\mathcal{U}}^\dagger \hat{a} \hat{\mathcal{U}} = \hbar\omega \hat{a}_H(t), \quad \hat{a}_H(0) = \hat{a},$$

which can be solved to give  $\hat{a}_H(t) = e^{-i\omega t} \hat{a}$ . Thus, we also have  $\hat{a}_H^\dagger(t) = e^{i\omega t} \hat{a}^\dagger$ .

Recall that we have

$$\hat{a}|\phi_0\rangle = 0, \quad \hat{a}|\phi_1\rangle = |\phi_0\rangle, \quad \hat{a}^\dagger|\phi_0\rangle = |\phi_1\rangle, \quad \hat{a}^\dagger|\phi_1\rangle = \sqrt{2}|\phi_2\rangle,$$

where  $|\phi_0\rangle$  and  $|\phi_1\rangle$  are the ground and first excited states of the Hamiltonian.

Therefore, we have

$$\begin{aligned} \langle\phi_0|\hat{a}|\phi_0\rangle &= \langle\phi_1|\hat{a}|\phi_1\rangle = \langle\phi_1|\hat{a}|\phi_0\rangle = 0, & \langle\phi_0|\hat{a}|\phi_1\rangle &= \langle\phi_0|\phi_0\rangle = 1, \\ \langle\phi_0|\hat{a}^\dagger|\phi_0\rangle &= \langle\phi_1|\hat{a}^\dagger|\phi_1\rangle = \langle\phi_0|\hat{a}^\dagger|\phi_1\rangle = 0, & \langle\phi_1|\hat{a}^\dagger|\phi_0\rangle &= \langle\phi_1|\phi_1\rangle = 1. \end{aligned}$$

Since the initial state is given as  $|\psi(0)\rangle = (|\phi_0\rangle + |\phi_1\rangle)/\sqrt{2}$ , the expectation value of  $\hat{a}$  and  $\hat{a}^\dagger$  at time  $t$  can be found as

$$\begin{aligned} \langle\hat{a}\rangle(t) &= \langle\psi(0)|\hat{a}_H(t)|\psi(0)\rangle = e^{-i\omega t} \langle\psi(0)|\hat{a}|\psi(0)\rangle = \frac{e^{-i\omega t}}{2} \langle(|\phi_0\rangle + |\phi_1\rangle)|\hat{a}(|\phi_0\rangle + |\phi_1\rangle)\rangle \\ &= \frac{e^{-i\omega t}}{2} (\langle\phi_0|\hat{a}|\phi_0\rangle + \langle\phi_1|\hat{a}|\phi_1\rangle + \langle\phi_0|\hat{a}|\phi_1\rangle + \langle\phi_1|\hat{a}|\phi_0\rangle) \\ &= \frac{e^{-i\omega t}}{2} (0 + 0 + 1 + 0) = \frac{e^{-i\omega t}}{2}, \\ \langle\hat{a}^\dagger\rangle(t) &= \langle\psi(0)|\hat{a}_H^\dagger(t)|\psi(0)\rangle = e^{i\omega t} \langle\psi(0)|\hat{a}^\dagger|\psi(0)\rangle = \frac{e^{i\omega t}}{2} \langle(|\phi_0\rangle + |\phi_1\rangle)|\hat{a}^\dagger(|\phi_0\rangle + |\phi_1\rangle)\rangle \\ &= \frac{e^{i\omega t}}{2} (\langle\phi_0|\hat{a}^\dagger|\phi_0\rangle + \langle\phi_1|\hat{a}^\dagger|\phi_1\rangle + \langle\phi_0|\hat{a}^\dagger|\phi_1\rangle + \langle\phi_1|\hat{a}^\dagger|\phi_0\rangle) \\ &= \frac{e^{i\omega t}}{2} (0 + 0 + 0 + 1) = \frac{e^{i\omega t}}{2}. \end{aligned}$$

Recall that the position operator  $\hat{x}$  and the momentum operator  $\hat{p}$  can be represented as

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}}(\hat{a} + \hat{a}^\dagger), \quad \hat{p} = -i\sqrt{\frac{m\hbar\omega}{2}}(\hat{a} - \hat{a}^\dagger).$$

By linearity of expectation, we could therefore determine the expectation of  $\hat{x}$  as

$$\langle\hat{x}\rangle(t) = \sqrt{\frac{\hbar}{2m\omega}} (\langle\hat{a}\rangle(t) + \langle\hat{a}^\dagger\rangle(t)) = \sqrt{\frac{\hbar}{2m\omega}} \left( \frac{e^{-i\omega t}}{2} + \frac{e^{i\omega t}}{2} \right) = \boxed{\sqrt{\frac{\hbar}{2m\omega}} \cos \omega t},$$

where we have used the identities  $e^{i\omega t} = \cos \omega t + i \sin \omega t$  and  $e^{-i\omega t} = \cos \omega t - i \sin \omega t$ .

(b) Similarly, we have

$$\langle\hat{p}\rangle(t) = -i\sqrt{\frac{m\hbar\omega}{2}} (\langle\hat{a}\rangle(t) - \langle\hat{a}^\dagger\rangle(t)) = -i\sqrt{\frac{m\hbar\omega}{2}} \left( \frac{e^{-i\omega t}}{2} - \frac{e^{i\omega t}}{2} \right) = -\sqrt{\frac{m\hbar\omega}{2}} \sin \omega t.$$

To determine the variance, we also need to calculate the expectation of  $\hat{x}^2$  and  $\hat{p}^2$ .

We have

$$\begin{aligned} \hat{x}^2 &= \frac{\hbar}{2m\omega} (\hat{a} + \hat{a}^\dagger)^2 = \frac{\hbar}{2m\omega} (\hat{a}^2 + \hat{a}\hat{a}^\dagger + \hat{a}^\dagger\hat{a} + \hat{a}^{\dagger 2}), \\ \hat{p}^2 &= -\frac{m\hbar\omega}{2} (\hat{a} - \hat{a}^\dagger)^2 = -\frac{m\hbar\omega}{2} (\hat{a}^2 - \hat{a}\hat{a}^\dagger - \hat{a}^\dagger\hat{a} + \hat{a}^{\dagger 2}). \end{aligned}$$

Note that, by definition, we have

$$\hat{a}^2|\phi_0\rangle = 0, \quad \hat{a}^2|\phi_1\rangle = \hat{a}|\phi_0\rangle = 0, \quad \hat{a}^{\dagger 2}|\phi_0\rangle = \hat{a}^\dagger|\phi_1\rangle = |\phi_2\rangle, \quad \hat{a}^{\dagger 2}|\phi_1\rangle = \sqrt{2}\hat{a}^\dagger|\phi_2\rangle = \sqrt{6}|\phi_3\rangle,$$

so we must have  $\langle\hat{a}^2\rangle(t) = 0$  since both states result in 0 and  $\langle\hat{a}^{\dagger 2}\rangle(t) = 0$  for any time  $t$  since both states are outside of (orthogonal to) the original states.

For the number operator  $\hat{a}^\dagger \hat{a}$ , we have

$$\hat{a}^\dagger \hat{a} |\phi_0\rangle = 0, \quad \hat{a}^\dagger \hat{a} |\phi_1\rangle = |\phi_1\rangle, \quad (\hat{a}^\dagger \hat{a})_{\text{H}} = \hat{a}_{\text{H}}^\dagger \hat{a}_{\text{H}}(t) = e^{i\omega t} \hat{a}^\dagger e^{-i\omega t} \hat{a} = \hat{a}^\dagger \hat{a},$$

so its expectation is

$$\begin{aligned} \langle \hat{a}^\dagger \hat{a} \rangle(t) &= \langle \psi(0) | (\hat{a}^\dagger \hat{a})_{\text{H}} | \psi(0) \rangle = \langle \psi(0) | \hat{a}^\dagger \hat{a} | \psi(0) \rangle = \frac{1}{2} \langle (|\phi_0\rangle + |\phi_1\rangle) | \hat{a}^\dagger \hat{a} (|\phi_0\rangle + |\phi_1\rangle) \rangle \\ &= \frac{1}{2} (\langle \phi_0 | \hat{a}^\dagger \hat{a} | \phi_0 \rangle + \langle \phi_1 | \hat{a}^\dagger \hat{a} | \phi_1 \rangle + \langle \phi_0 | \hat{a}^\dagger \hat{a} | \phi_1 \rangle + \langle \phi_1 | \hat{a}^\dagger \hat{a} | \phi_0 \rangle) \\ &= \frac{1}{2} (0 + 1 + 0 + 0) = \frac{1}{2}. \end{aligned}$$

By the identity  $[\hat{a}, \hat{a}^\dagger] = \hat{a}\hat{a}^\dagger - \hat{a}^\dagger\hat{a} = 1$ , we have

$$\langle \hat{a}\hat{a}^\dagger \rangle(t) = 1 + \langle \hat{a}^\dagger \hat{a} \rangle(t) = 1 + \frac{1}{2} = \frac{3}{2}.$$

Finally, we are able to compute

$$\begin{aligned} \langle \hat{x}^2 \rangle(t) &= \frac{\hbar}{2m\omega} (\langle \hat{a}^2 \rangle(t) + \langle \hat{a}\hat{a}^\dagger \rangle(t) + \langle \hat{a}^\dagger \hat{a} \rangle(t) + \langle \hat{a}^{\dagger 2} \rangle(t)) = \frac{\hbar}{2m\omega} \left(0 + \frac{3}{2} + \frac{1}{2} + 0\right) = \frac{\hbar}{m\omega}, \\ \langle \hat{p}^2 \rangle(t) &= -\frac{m\hbar\omega}{2} (\langle \hat{a}^2 \rangle(t) - \langle \hat{a}\hat{a}^\dagger \rangle(t) - \langle \hat{a}^\dagger \hat{a} \rangle(t) + \langle \hat{a}^{\dagger 2} \rangle(t)) = -\frac{m\hbar\omega}{2} \left(0 - \frac{3}{2} - \frac{1}{2} + 0\right) = m\hbar\omega. \end{aligned}$$

The variance the position and momentum at time  $t$  are, respectively,

$$\begin{aligned} \sigma_x^2(t) &= \langle \hat{x}^2 \rangle(t) - (\langle \hat{x} \rangle(t))^2 = \frac{\hbar}{m\omega} - \left( \sqrt{\frac{\hbar}{2m\omega}} \cos \omega t \right)^2 = \boxed{\frac{\hbar}{m\omega} \left(1 - \frac{1}{2} \cos^2 \omega t\right)}, \\ \sigma_p^2(t) &= \langle \hat{p}^2 \rangle(t) - (\langle \hat{p} \rangle(t))^2 = m\hbar\omega - \left( -\sqrt{\frac{m\hbar\omega}{2}} \sin \omega t \right)^2 = \boxed{m\hbar\omega \left(1 - \frac{1}{2} \sin^2 \omega t\right)}. \end{aligned}$$

The product between these variances is

$$\begin{aligned} \sigma_x^2(t) \sigma_p^2(t) &= \frac{\hbar}{m\omega} \left(1 - \frac{1}{2} \cos^2 \omega t\right) \cdot m\hbar\omega \left(1 - \frac{1}{2} \sin^2 \omega t\right) \\ &= \frac{\hbar^2}{4} (2 - \cos^2 \omega t) (2 - \sin^2 \omega t), \end{aligned}$$

and because  $\cos^2 \omega t, \sin^2 \omega t \leq 1$  for  $t \in \mathbb{R}$ , we have

$$\sigma_x^2(t) \sigma_p^2(t) \geq \frac{\hbar^2}{4} (2 - 1) (2 - 1) = \frac{\hbar^2}{4},$$

which is indeed consistent with the Heisenberg uncertainty principle.

In fact, for this case, we can calculate the lower bound of the variance product  $\sigma_x^2(t) \sigma_p^2(t)$  by using  $\cos^2 x + \sin^2 x = 1$  where  $x = \omega t$ , so that

$$\begin{aligned} (2 - \cos^2 x) (2 - \sin^2 x) &= (2 - \cos^2 x)(1 + \cos^2 x) \\ &= 2 + \cos^2 x - \cos^4 x = \frac{9}{4} - \left( \cos^2 x - \frac{1}{2} \right)^2, \end{aligned}$$

which is minimised when  $\cos^2 x = 0$  or  $\cos^2 x = 1$  since  $0 \leq \cos^2 x \leq 1$ , with minimum  $9/4 - 1/4 = 2$ .

This corresponds to the cases where  $x = k\pi$  or  $x = k\pi + \pi/2$  where  $k \in \mathbb{Z}$ , i.e.,

$$t = \frac{k\pi}{\omega} \text{ or } \frac{k\pi + \pi/2}{\omega}.$$

At these times, the variance product is minimised at  $(\sigma_x^2(t) \sigma_p^2(t))_{\min} = \hbar^2/2$ .