

B.1. Use Gaussian elimination to find every solution to the following equation:

$$\begin{pmatrix} 2 & 4 & 1 \\ 2 & 6 & 1 \\ 3 & 9 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

We know how to solve  $Ax = 0$ , but this is of the form  $Ax = x$ . We need to transform the given equation into one we know how to handle.

$$\begin{aligned} \begin{pmatrix} 2 & 4 & 1 \\ 2 & 6 & 1 \\ 3 & 9 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \\ \left( \begin{pmatrix} 2 & 4 & 1 \\ 2 & 6 & 1 \\ 3 & 9 & 3 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \\ \begin{pmatrix} 1 & 4 & 1 \\ 2 & 5 & 1 \\ 3 & 9 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \end{aligned}$$

Then we can apply Gaussian elimination to find the solutions.

$$\begin{pmatrix} 1 & 4 & 1 \\ 2 & 5 & 1 \\ 3 & 9 & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 4 & 1 \\ 0 & -3 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

Giving a solution set

$$\left\{ \alpha \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} : \alpha \in \mathbb{R} \right\}.$$

B.2. Let  $A$  be an  $n \times m$  matrix, and  $b \in \mathbb{R}^n$ . Suppose  $Ax = b$  has at least one solution  $x_0 \in \mathbb{R}^m$ . Show all solutions are of the form  $x = x_0 + h$ , where  $h$  solves  $Ah = 0$ .

Suppose that  $y$  is a different solution to  $Ax = b$  than  $x_0$ . Let  $h = (y - x_0)$ . Then  $y = x_0 + h$ . Additionally

$$\begin{aligned} Ah &= A(y - x_0) \\ &= Ay - Ax_0 \\ &= b - b \\ &= 0 \end{aligned}$$

Therefore every solution of  $Ax = b$  is of the form  $x = x_0 + h$ , where  $h$  solves  $Ah = 0$ .

B.3. Let  $A$  and  $B$  be square  $n \times n$  matrices with real entries. For each of the following statements, either give a **proof**, or find a **counterexample with**  $n = 2$ .

(i) If  $AB = 0$  then  $A$  and  $B$  cannot both be invertible.

Suppose that  $AB = 0$  and that  $A$  is invertible. Then

$$\begin{aligned} B &= A^{-1}AB \\ &= A^{-1}0 \\ &= 0 \end{aligned}$$

and therefore  $B$  is not invertible. Very similarly, if  $AB = 0$  and  $B$  is invertible, then  $A = 0$ , and is therefore non-invertible.

Thus, if  $AB = 0$  then  $A$  and  $B$  cannot both be invertible.

- (ii) If  $A$  and  $B$  are invertible then  $A + B$  is invertible.

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad B = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

Then  $A^{-1} = A$  and  $B^{-1} = B$ , but  $A + B$  is not invertible.

- (iii) If  $A$  and  $B$  are invertible then  $AB$  is invertible.

Let  $A$  and  $B$  be invertible matrices. Then  $AB(B^{-1}A^{-1}) = I$ , so  $AB$  has an inverse, and is hence invertible.

- (iv) If  $A$  and  $B$  are invertible and  $(AB)^2 = A^2B^2$ , then  $AB = BA$ . Suppose that  $A$  and  $B$  are invertible, and that  $(AB)^2 = A^2B^2$ . Then

$$\begin{aligned} ABAB &= AAB B \\ A^{-1}ABAB B^{-1} &= A^{-1}AAB B B^{-1} \\ BA &= AB \end{aligned}$$

- (v) If  $ABA = 0$  and  $B$  is invertible then  $A^2 = 0$ .

Whenever you look for counterexamples involving matrices, especially if the property in question uses multiplication, I would advise you to consider two types of matrix. First, the elementary matrices. These are all invertible, and multiplying by them is very easy to calculate. Second, those with lots and lots of zeros in them, as they are easy to calculate.

In this case, these two types of matrices give a very nice counterexample.

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

- (vi) If  $ABA = I$  then  $A$  is invertible and  $B = (A^{-1})^2$ .

The uniqueness of inverses shows that if  $ABA = I$  then  $A^{-1} = BA$ . This means that if  $ABA = I$  then  $A$  is invertible. Furthermore

$$\begin{aligned} A^{-1} &= BA \\ (A^{-1})^2 &= BAA^{-1} \\ (A^{-1})^2 &= B \end{aligned}$$

as desired.

- (vii) If  $A$  has a left inverse  $B$  and a right inverse  $C$  then  $B = C$ .

The key to proving this is to consider  $BAC$ . If we multiply  $BA$  first, we get  $C$ , and if we multiply  $AC$  first, we get  $B$ . We know that matrix multiplication is associative, and that is doesn't matter what order we do the multiplication in, so we conclude that  $B = C$ .

B.4. Let  $n \geq 2$  and let  $A_n = (a_{ij})$  be the  $n \times n$  matrix such that

$$\begin{aligned} a_{i-1,i} &= 1 \text{ for } i = 2, \dots, n, \\ a_{i+1,i} &= 1 \text{ for } i = 1, \dots, n-1, \end{aligned}$$

and  $a_{ij} = 0$  for all other  $i, j$ . Write down  $A_2, A_3$  and  $A_4$ . Prove that  $A_n$  is invertible for all even values of  $n$ , and is not invertible for all odd values of  $n$ . Find  $A_2^{-1}$  and  $A_4^{-1}$ .

For  $n$  even you can reduce  $A_n$  to the identity by subtracting row  $n$  from row  $n-2$ , then row  $n-2$  from row  $n-4$  and so on, and then swapping rows; hence  $A_n$  is invertible. For  $n$  odd you can reduce to a matrix with a bottom row of zeros by subtracting row 1 from row 3, then row 3 from row 5 and so on; hence  $A_n$  is not invertible.