

MATH50001 Problems Sheet 5

Solutions

1)

a)

$$\frac{1}{2} \cdot \frac{1}{z-2i} + \sum_{n=0}^{\infty} \frac{i^{n-1}}{2^{2n+3}} (z-2i)^n, \quad 0 < |z-2i| < 4.$$

b)

$$\sum_{n=-1}^{\infty} \frac{1}{e(n+1)!} (z+1)^n, \quad |z+1| > 0.$$

2)

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2^n(z-1)^n} - \sum_{n=0}^{\infty} (z-1)^n.$$

3) Obviously

$$\frac{9}{(z-4)(z+5)} = \frac{1}{z-4} - \frac{1}{z+5}.$$

3a) If $|z| < 4$ we have

$$\frac{1}{z-4} = -\frac{1}{4-z} = -\frac{1}{4} \frac{1}{1-z/4} = -\frac{1}{4} \sum_{n=0}^{\infty} \frac{1}{4^n} z^n$$

and

$$-\frac{1}{z+5} = -\frac{1}{5} \frac{1}{1-(-z/5)} = -\frac{1}{5} \sum_{n=0}^{\infty} \frac{(-1)^n}{5^n} z^n.$$

Therefore

$$\frac{9}{(z-4)(z+5)} = -\sum_{n=0}^{\infty} \left(\frac{1}{4^{n+1}} + \frac{(-1)^n}{5^{n+1}} \right) z^n.$$

3b) If $2 < |z| < 5$, then

$$\frac{1}{z-4} = \frac{1}{z} \frac{1}{1-4/z} = \frac{1}{z} \sum_{n=0}^{\infty} \frac{4^n}{z^n}$$

2

and thus

$$\begin{aligned}\frac{9}{(z-4)(z+5)} &= \sum_{n=0}^{\infty} \frac{4^n}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{(-1)^n}{5^{n+1}} z^n \\ &= \sum_{n=-\infty}^{-1} 4^{-n-1} z^n - \sum_{n=0}^{\infty} \frac{(-1)^n}{5^{n+1}} z^n.\end{aligned}$$

3c) If $5 < |z|$, then

$$-\frac{1}{z+5} = -\frac{1}{z} \frac{1}{1 - (-5/z)} = -\frac{1}{z} \sum_{n=0}^{\infty} \left(\frac{-5}{z}\right)^n$$

which implies

$$\begin{aligned}\frac{9}{(z-4)(z+5)} &= \sum_{n=0}^{\infty} \frac{4^n}{z^{n+1}} - \frac{1}{z} \sum_{n=0}^{\infty} \left(\frac{-5}{z}\right)^n \\ &= \sum_{n=-\infty}^{-1} (4^{-n-1} - (-5)^{-n-1}) z^n.\end{aligned}$$

4) Let $f(z) = z e^z$ and $z_0 = 2$. Then

$$\begin{aligned}f(z) &= (z-2) e^{z-2} e^2 + 2e^{z-2} e^2 = \sum_{n=0}^{\infty} \frac{e^2}{n!} (z-2)^{n+1} + \sum_{n=0}^{\infty} \frac{2e^2}{n!} (z-2)^n \\ &= 2e^2 + \sum_{n=1}^{\infty} \left(\frac{e^2}{(n-1)!} + \frac{2e^2}{n!} \right) (z-2)^n.\end{aligned}$$

5)

a) If f is holomorphic at z_0 and has a zero of order m at z_0 , then there is $g(z)$ holomorphic at z_0 , $g(z_0) \neq 0$ such that $f(z) = (z - z_0)^m g(z)$. Therefore

$$\frac{1}{f(z)} = \frac{1}{(z - z_0)^m} \cdot \frac{1}{g(z)}.$$

Thus $1/f$ has a pole of order m at z_0 .

b)

$$\begin{aligned}(2 \cos z - 2 - z^2)^2 &= \left(2 \left(1 - \frac{1}{2} z^2 + \frac{1}{4!} z^4 - \frac{1}{6!} z^6 + \dots\right) - 2 + z^2\right)^2 \\ &= 4 \left(\frac{1}{4!} z^4 - \frac{1}{6!} z^6 + \dots\right)^2 = z^8 \cdot g(z),\end{aligned}$$

where $g(z)$ is holomorphic at 0 and $g(0) \neq 0$. Therefore $\frac{1}{(2 \cos z - 2 + z^2)^2}$ has a pole at 0 of order 8.

6)

- a) $z = 0$, essential singularity;
- b) $z = 0$, pole of order 4;
- c) $z = n\pi, (6n \pm 1)\pi/3$, poles of order 1.

7) The function $f(z) = \frac{e^z}{z(z-2)^3}$ has two poles inside $\gamma = \{|z| = 3\}$. One of them is at $z_1 = 0$ of order one and the other one is at $z_2 = 2$ of order three. Therefore

$$\begin{aligned} \oint_{\gamma} \frac{e^z}{z(z-2)^3} dz &= 2\pi i (\text{Res}[f, z_1] + \text{Res}[f, z_2]) \\ &= 2\pi i \left(\frac{e^0}{(-2)^3} + \lim_{z \rightarrow 2} \frac{1}{2} \frac{d^2}{dz^2} \frac{(z-2)^3 e^z}{z(z-2)^3} \right) \\ &= -\frac{\pi i}{4} + \pi i \lim_{z \rightarrow 2} \left(\frac{e^z}{z} - 2 \frac{e^z}{z^2} + 2 \frac{e^z}{z^3} \right) \\ &= -\frac{\pi i}{4} + \pi i e^2 \left(\frac{1}{2} - \frac{1}{2} + \frac{1}{4} \right) = \frac{\pi(e^2 - 1)i}{4}. \end{aligned}$$

8) Substituting $z = e^{i\theta}$ gives

$$\begin{aligned} I &:= \int_0^{2\pi} \frac{d\theta}{1 - 2a \cos \theta + a^2} = \frac{1}{i} \oint_{|z|=1} \frac{1}{-az^2 + (1+a^2)z - a} dz \\ &= \frac{1}{i} \oint_{|z|=1} \frac{1}{(z-a)(1-az)} dz. \end{aligned}$$

If $|a| < 1$ there is a pole inside $|z| = 1$ at a with the residue $1/(1-a^2)$. Therefore $I = 2\pi/(1-a^2)$.

If $|a| > 1$ the pole inside $|z| = 1$ is at $1/a$ and the residue is

$$\lim_{z \rightarrow 1/a} \frac{z - 1/a}{(z-a)(az-1)} = \frac{1}{a^2 - 1}.$$

Hence $I = 2\pi/(a^2 - 1)$.

9)

$$\begin{aligned}
\oint_{\gamma} \frac{e^z - 1}{z^2(z-1)} &= 2\pi i \operatorname{Res} \left[\frac{e^z - 1}{z^2(z-1)}, 1 \right] - 2\pi i \operatorname{Res} \left[\frac{e^z - 1}{z^2(z-1)}, 0 \right] \\
&= 4\pi i \left\{ \frac{e^z - 1}{z^2} \Big|_{z=1} - \frac{z}{dz} \frac{e^z - 1}{z-1} \Big|_{z=0} \right\} \\
&= 4\pi i \left\{ e - 1 - \frac{e^z(z-1) - (e^z - 1)}{(z-1)^2} \Big|_{z=0} \right\} \\
&= 4\pi i \left\{ e - 1 + 1 \Big|_{z=0} \right\} = 4\pi i e.
\end{aligned}$$

10) Indeed,

$$\begin{aligned}
\frac{1}{2\pi i} \oint_{|z|=r} z^{n-1} |f(z)|^2 dz &= \frac{1}{2\pi i} \int_0^{2\pi} r^{n-1} e^{i(n-1)\theta} |f(re^{i\theta})|^2 i e^{i\theta} r d\theta \\
&= \frac{r^n}{2\pi} \int_0^{2\pi} e^{i(n-1)\theta} \left(\sum_{k=0}^n a_k r^k e^{ik\theta} \right) \overline{\left(\sum_{m=0}^n a_m r^m e^{im\theta} \right)} e^{i\theta} d\theta.
\end{aligned}$$

The only term that survive if $n - 1 + k - m + 1 = 0$. The only possibility for that is $k = 0$ and $m = n$ and thus

$$\frac{1}{2\pi i} \oint_{|z|=r} z^{n-1} |f(z)|^2 dz = a_0 \bar{a}_n r^{2n}.$$