

# A Simple L<sup>A</sup>T<sub>E</sub>X Template

Cunxin Huang \*

April 2, 2023 8:35am

## Abstract

This is an example.

**Keywords:** Keyword1, Keyword2, Keyword3, Keyword4, Keyword5

## 1 Example section 1

**Definition 1.** This is an example definition.

**Theorem 1.** *This is an example theorem.*

**Proof.** This is the proof for Theorem 1.

Recall Definition 1.

The proof is complete. □

**Proposition 1.** *This is an example proposition.*

**Proof.** According to [?, Theorem 3.14], we have

$$e^{i\pi} + 1 = 0. \tag{1.1} \boxed{\text{eq:euler}}$$

This completes the proof. □

**Corollary 1.** *This is an example corollary.*

**Proof.** Due to equation (1.1),

$$e^{i\pi} = -1,$$

which finishes the proof. □

---

\*Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong, China  
(huangcunxin@outlook.com).

## 2 Example section 2

`<sec:example2>?` To continue the discussion in Section 1, we first recall Algorithm 6.18 in [?].

---

**Algorithm 2.1** An example algorithm

---

`<alg:example>?` Input  $x_0 \in \mathbb{R}^n$ ,  $\rho_0 > 0$ ,  $\eta_1 \geq 1 > \eta_2 > 0$ . For  $k = 0, 1, 2, \dots$ , do the following.

1. Choose a set of nonzero vectors  $D_k \subset \mathbb{R}^n$  (deterministically or stochastically).
  2. Check whether there exists a  $y_k \in \{x_k + \rho_k d : d \in D_k\}$  such that  $f(y_k) < f(x_k)$ .
  3. If  $y_k$  exists, set  $x_{k+1} = y_k$ ,  $\rho_{k+1} = \eta_1 \rho_k$ ; otherwise, set  $x_{k+1} = x_k$ ,  $\rho_{k+1} = \eta_2 \rho_k$ .
  4. If a certain stopping criterion is met, exit and output  $x_{k+1}$ .
-