

# Non-convergence Analysis of Probabilistic Direct Search

---

Cunxin Huang

Co-supervised by Dr. Zaikun Zhang and Prof. Xiaojun Chen

Graduate Student Forum of ORSC Mathematical Programming Branch  
Chongqing, China, 2023

Department of Applied Mathematics  
The Hong Kong Polytechnic University

# Table of contents

## 1. Preliminaries

What is derivative-free optimization and why

Probabilistic direct search

## 2. Non-convergence analysis

Motivation and basic idea

Main results

## 3. Tightness of analysis

Almost zero gap

Tightness of assumptions

## 4. Conclusions

# Preliminaries

---

# What is derivative-free optimization and why

## Derivative-free optimization (DFO)

- A branch of optimization
- Do not use derivatives (only use function evaluations)

# What is derivative-free optimization and why

## Derivative-free optimization (DFO)

- A branch of optimization
- Do not use derivatives (only use function evaluations)

## Why DFO?

- Derivatives are not available
- Problems are often noisy (finite difference?)
- Function evaluations are expensive (e.g.: PDE simulation)

# What is derivative-free optimization and why


## Derivative-free optimization (DFO)

- A branch of optimization
- Do not use derivatives (only use function evaluations)

## Why DFO?

- Derivatives are not available
- Problems are often noisy (finite difference?)
- Function evaluations are expensive (e.g.: PDE simulation)

## Typical situation: black box




# What is derivative-free optimization and why

## Derivative-free optimization (DFO)

- A branch of optimization
- **Do not use derivatives** (only use function evaluations)

## Why DFO?

- Derivatives are **not available**
- Problems are often **noisy** (finite difference?)
- Function evaluations are **expensive** (e.g.: PDE simulation)



Nuclear Physics



Machine Learning



Cosmology

In this talk, we consider the **unconstrained** problem

$$\min_{x \in \mathbb{R}^n} f(x),$$

where

- $\nabla f$  is **Lipschitz continuous**, although it cannot be evaluated,
- $f$  is bounded below.



---

**Algorithm 1:** Probabilistic direct search based on sufficient decrease

---

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on sufficient decrease

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on [sufficient decrease](#)

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on sufficient decrease

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on [sufficient decrease](#)

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on [sufficient decrease](#)

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on [sufficient decrease](#)

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

        Expand step size, and move to that point

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on [sufficient decrease](#)

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

        Expand step size, and move to that point

        Set  $\alpha_{k+1} = \gamma \alpha_k$  and  $x_{k+1} = x_k + \alpha_k d$ .

**else**



# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on [sufficient decrease](#)

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

        Expand step size, and move to that point

        Set  $\alpha_{k+1} = \gamma \alpha_k$  and  $x_{k+1} = x_k + \alpha_k d$ .

**else**

        Shrink step size, and stand still

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on sufficient decrease

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly.

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

        Expand step size, and move to that point

        Set  $\alpha_{k+1} = \gamma \alpha_k$  and  $x_{k+1} = x_k + \alpha_k d$ .

**else**

        Shrink step size, and stand still

        Set  $\alpha_{k+1} = \theta \alpha_k$  and  $x_{k+1} = x_k$ .

---

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on sufficient decrease

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly. **How to select?**

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

        Expand step size, and move to that point

        Set  $\alpha_{k+1} = \gamma \alpha_k$  and  $x_{k+1} = x_k + \alpha_k d$ .

**else**

        Shrink step size, and stand still

        Set  $\alpha_{k+1} = \theta \alpha_k$  and  $x_{k+1} = x_k$ .

---

# Probabilistic direct search (PDS)

---

**Algorithm 1:** Probabilistic direct search based on sufficient decrease

---

**Input:**  $x_0 \in \mathbb{R}^n$ ,  $\alpha_0 \in (0, \infty)$ ,  $0 < \theta < 1 < \gamma$ .

**for**  $k = 0, 1, \dots$  **do**

    Select a finite set  $D_k \subset \mathbb{R}^n$  randomly. **How to select?**

    (In this talk, assume  $D_k$  is a set of unit vectors for simplicity.)

**if**  $f(x_k + \alpha_k d) < f(x_k) - \alpha_k^2$  for some  $d \in D_k$  **then**

        Expand step size, and move to that point

        Set  $\alpha_{k+1} = \gamma \alpha_k$  and  $x_{k+1} = x_k + \alpha_k d$ .

**else**


        Shrink step size, and stand still

        Set  $\alpha_{k+1} = \theta \alpha_k$  and  $x_{k+1} = x_k$ .


---

Typical choice:  $D_k = \{d_1, \dots, d_m\}$ , where  $d_i \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(\mathcal{S}^{n-1})$


# Illustration of how PDS works




# Illustration of how PDS works



## Illustration of how PDS works




## Illustration of how PDS works







## Illustration of how PDS works




## Illustration of how PDS works




# Illustration of how PDS works




# Illustration of how PDS works




# Illustration of how PDS works




# Illustration of how PDS works



# Illustration of how PDS works




# Illustration of how PDS works





# Illustration of how PDS works



## Definition (Cosine measure)

Cosine measure for a finite set of nonzero vectors  $D \subseteq \mathbb{R}^n$  w.r.t. a given vector  $v \in \mathbb{R}^n$ :

$$\text{cm}(D, v) = \max_{d \in D} \frac{d^\top v}{\|d\| \|v\|}.$$

# Convergence theory


## Definition (Cosine measure)

Cosine measure for a finite set of nonzero vectors  $D \subseteq \mathbb{R}^n$  w.r.t. a given vector  $v \in \mathbb{R}^n$ :

$$\text{cm}(D, v) = \max_{d \in D} \frac{d^\top v}{\|d\| \|v\|}.$$

## Example

$$\text{cm}(D, v) = \cos \theta$$



# Convergence theory


## Definition (Cosine measure)

Cosine measure for a finite set of nonzero vectors  $D \subseteq \mathbb{R}^n$  w.r.t. a given vector  $v \in \mathbb{R}^n$ :

$$\text{cm}(D, v) = \max_{d \in D} \frac{d^\top v}{\|d\| \|v\|}.$$

## Example

$$\text{cm}(D, v) = \cos \theta$$



Measure the ability that “ $D$  approximates  $v$ ”

## Definition ( $p$ -probabilistically $\kappa$ -descent)

$(D_k)_{k \geq 0}$  is said to be  $p$ -probabilistically  $\kappa$ -descent, if

$$\mathbb{P}(\text{cm}(D_k, -g_k) \geq \kappa \mid D_0, \dots, D_{k-1}) \geq p \quad \text{for each } k \geq 0,$$

where  $g_k = \nabla f(x_k)$ .

# Convergence theory

## Definition ( $p$ -probabilistically $\kappa$ -descent)

$(D_k)_{k \geq 0}$  is said to be  $p$ -probabilistically  $\kappa$ -descent, if

$$\mathbb{P}(\text{cm}(D_k, -g_k) \geq \kappa \mid D_0, \dots, D_{k-1}) \geq p \quad \text{for each } k \geq 0,$$

where  $g_k = \nabla f(x_k)$ .

Intuition:

each  $D_k$  is “good enough with lower-bounded probability”,  
no matter what happened before

# Convergence theory

## Definition ( $p$ -probabilistically $\kappa$ -descent)

$(D_k)_{k \geq 0}$  is said to be  $p$ -probabilistically  $\kappa$ -descent, if

$$\mathbb{P}(\text{cm}(D_k, -g_k) \geq \kappa \mid D_0, \dots, D_{k-1}) \geq p \quad \text{for each } k \geq 0,$$

where  $g_k = \nabla f(x_k)$ .

Intuition:

each  $D_k$  is “good enough with lower-bounded probability”,  
no matter what happened before

## Theorem (Gratton et al. 2015)

If  $(D_k)_{k \geq 0}$  is  $p$ -probabilistically  $\kappa$ -descent with  $\kappa > 0$  and

$$p = \log \theta / \log(\gamma^{-1} \theta),$$

then PDS converges with probability 1.

Corollary (Gratton et al. 2015)

If  $D_k = \{d_1, \dots, d_m\}$ , where  $d_i \stackrel{i.i.d.}{\sim} \text{Unif}(\mathcal{S}^{n-1})$ , then PDS is convergent if

$$m > \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right).$$



Corollary (Gratton et al. 2015)

If  $D_k = \{d_1, \dots, d_m\}$ , where  $d_i \stackrel{i.i.d.}{\sim} \text{Unif}(\mathcal{S}^{n-1})$ , then PDS is convergent if

$$m > \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right).$$


A nature question: what if

$$m \leq \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right)?$$

# Practical choice and natural question

A nature question: what if

$$m \leq \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right) ?$$



## Non-convergence analysis

---

# Motivation: non-convergence analysis matters

Many well-known algorithms have non-convergence analysis.

- S. Reddi, S. Kale, and S. Kumar. On the convergence of Adam and beyond. In Y. Bengio, Y. LeCun, T. Sainath, I. Murray, M. Ranzato, and O. Vinyals, editors, *International Conference on Learning Representations (ICLR 2018)*. Curran Associates, Inc., 2018.
- C. Chen, B. He, Y. Ye, and X. Yuan. The direct extension of ADMM for multi-block convex minimization problems is not necessarily convergent. *Math. Program.*, 155:57-79, 2016.
- W. Mascarenhas. The divergence of the BFGS and Gauss Newton methods. *Math. Program.*, 147:253-276, 2014.
- ...

# Naive idea of non-convergence

Recall  $p$ -probabilistically  $\kappa$ -descent:

$$\mathbb{P}(\text{cm}(D_k, -g_k) \geq \kappa \mid D_0, \dots, D_{k-1}) \geq p \quad \text{for each } k \geq 0.$$

# Naive idea of non-convergence

Recall  $p$ -probabilistically  $\kappa$ -descent:

$$\mathbb{P}(\text{cm}(D_k, -g_k) \geq \kappa \mid D_0, \dots, D_{k-1}) \geq p \quad \text{for each } k \geq 0.$$


$q$ -probabilistically ascent

$$\mathbb{P}(\text{cm}(D_k, -g_k) > 0 \mid D_0, \dots, D_{k-1}) \leq q \quad \text{for each } k \geq 0.$$

# Naive idea of non-convergence

$q$ -probabilistically ascent


$$\mathbb{P}(\text{cm}(D_k, -g_k) > 0 \mid D_0, \dots, D_{k-1}) \leq q \quad \text{for each } k \geq 0.$$



# Naive idea of non-convergence

$q$ -probabilistically ascent

$$\mathbb{P}(\text{cm}(D_k, -g_k) > 0 \mid D_0, \dots, D_{k-1}) \leq q \quad \text{for each } k \geq 0.$$



non-convergence for convex functions



non-convergence in general



## Establishment of non-convergence

$(D_k)_{k \in \mathbb{N}}$  is  
 $q$ -probabilistically ascent

## Establishment of non-convergence

$(D_k)_{k \in \mathbb{N}}$  is  
 $q$ -probabilistically ascent



$\alpha_k$  shrinks with high probability

## Establishment of non-convergence

$(D_k)_{k \in \mathbb{N}}$  is  
 $q$ -probabilistically ascent



$\alpha_k$  shrinks with high probability



$$\sum_{k=0}^{\infty} \alpha_k < \infty \quad \text{a.s. ?}$$

# Establishment of non-convergence

$(D_k)_{k \in \mathbb{N}}$  is  
 $q$ -probabilistically ascent



$\alpha_k$  shrinks with high probability



$$\sum_{k=0}^{\infty} \alpha_k < \infty \quad \text{a.s. ?}$$



$$x_0 \notin \bar{B}(x^*, \sum_{k=0}^{\infty} \alpha_k)$$

# Establishment of non-convergence

$(D_k)_{k \in \mathbb{N}}$  is  
 $q$ -probabilistically ascent



$\alpha_k$  shrinks with high probability



$$\sum_{k=0}^{\infty} \alpha_k < \infty \quad \text{a.s. ?}$$



$$x_0 \notin \bar{\mathcal{B}}(x^*, \sum_{k=0}^{\infty} \alpha_k)$$



$$\mathbb{P}(\text{Convergence}) < 1$$

- Define indicator function  $Y_k = \mathbb{1}_{\{\text{cm}(D_k, -g_k) > 0\}}$   
Indicator for “good” event

- Define indicator function  $Y_k = \mathbb{1}_{\{\text{cm}(D_k, -g_k) > 0\}}$   
Indicator for “good” event
- $\alpha_{k+1} \leq \gamma^{Y_k} \theta^{1-Y_k} \alpha_k$ , when  $f$  is convex

- Define indicator function  $Y_k = \mathbb{1}_{\{\text{cm}(D_k, -g_k) > 0\}}$   
Indicator for “good” event
- $\alpha_{k+1} \leq \gamma^{Y_k} \theta^{1-Y_k} \alpha_k$ , when  $f$  is convex
- $\alpha_k \leq \alpha_0 \prod_{i=0}^{k-1} \gamma^{Y_i} \theta^{1-Y_i} =: \alpha_0 U_k$



- Define indicator function  $Y_k = \mathbb{1}_{\{\text{cm}(D_k, -g_k) > 0\}}$   
Indicator for “good” event
- $\alpha_{k+1} \leq \gamma^{Y_k} \theta^{1-Y_k} \alpha_k$ , when  $f$  is convex
- $\alpha_k \leq \alpha_0 \prod_{i=0}^{k-1} \gamma^{Y_i} \theta^{1-Y_i} =: \alpha_0 U_k$
- $\sum_{k=1}^{\infty} \alpha_k \leq \alpha_0 \sum_{k=1}^{\infty} U_k$

- Define indicator function  $Y_k = \mathbb{1}_{\{\text{cm}(D_k, -g_k) > 0\}}$   
Indicator for “good” event
- $\alpha_{k+1} \leq \gamma^{Y_k} \theta^{1-Y_k} \alpha_k$ , when  $f$  is convex
- $\alpha_k \leq \alpha_0 \prod_{i=0}^{k-1} \gamma^{Y_i} \theta^{1-Y_i} =: \alpha_0 U_k$
- $\sum_{k=1}^{\infty} \alpha_k \leq \alpha_0 \sum_{k=1}^{\infty} U_k < \infty$  a.s. ?

# Key to analysis

- Define indicator function  $Y_k = \mathbb{1}_{\{\text{cm}(D_k, -g_k) > 0\}}$   
Indicator for “good” event
- $\alpha_{k+1} \leq \gamma^{Y_k} \theta^{1-Y_k} \alpha_k$ , when  $f$  is convex
- $\alpha_k \leq \alpha_0 \prod_{i=0}^{k-1} \gamma^{Y_i} \theta^{1-Y_i} =: \alpha_0 U_k$
- $\sum_{k=1}^{\infty} \alpha_k \leq \alpha_0 \sum_{k=1}^{\infty} U_k < \infty$  a.s. ?

Under  $q$ -probabilistically ascent assumption, can we find a constant  $\zeta$  such that

$$\mathbb{P} \left( \sum_{k=1}^{\infty} U_k < \zeta \right) > 0?$$

## Assumption

$\mathbb{P}(\text{cm}(D_k, -g_k) \leq 0 \mid D_0, \dots, D_{k-1}) \geq q > q_0$  for each  $k \geq 0$ ,  
where  $q_0 = 1 - p_0 = \log \gamma / \log(\theta^{-1} \gamma)$ .

## Assumption

$\mathbb{P}(Y_k = 0 \mid Y_0, \dots, Y_{k-1}) \geq q > q_0$  for each  $k \geq 0$ ,  
where  $q_0 = 1 - p_0 = \log \gamma / \log(\theta^{-1}\gamma)$ .

# Main results

## Assumption

$$\mathbb{P}(Y_k = 0 \mid Y_0, \dots, Y_{k-1}) \geq q > q_0 \quad \text{for each } k \geq 0,$$

where  $q_0 = 1 - p_0 = \log \gamma / \log(\theta^{-1} \gamma)$ .

## Result

1.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \infty\right) = 1$$

2.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \zeta\right) > 0 \iff \zeta > \frac{\theta}{1 - \theta}$$

# Main results

## Assumption

$$\mathbb{P}(Y_k = 0 \mid Y_0, \dots, Y_{k-1}) \geq q > q_0 \quad \text{for each } k \geq 0,$$

where  $q_0 = 1 - p_0 = \log \gamma / \log(\theta^{-1} \gamma)$ .

## Result

1.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \infty\right) = 1$$

2.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \zeta\right) > 0 \iff \zeta > \frac{\theta}{1 - \theta}$$

Note that  $\sum_{k=1}^{\infty} U_k = \sum_{k=1}^{\infty} \prod_{i=0}^{k-1} \gamma^{Y_i} \theta^{1-Y_i} \geq \theta / (1 - \theta)$

# Main results

## Assumption

$$\mathbb{P}(Y_k = 0 \mid Y_0, \dots, Y_{k-1}) \geq q > q_0 \quad \text{for each } k \geq 0,$$

where  $q_0 = 1 - p_0 = \log \gamma / \log(\theta^{-1} \gamma)$ .

## Result

1.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \infty\right) = 1$$

2.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \zeta\right) > 0 \iff \zeta > \frac{\theta}{1 - \theta}$$



# Main results

## Assumption

$$\liminf_{k \rightarrow \infty} \mathbb{P}(Y_k = 0 \mid Y_0, \dots, Y_{k-1}) > q_0,$$

where  $q_0 = 1 - p_0 = \log \gamma / \log(\theta^{-1} \gamma)$ .

## Result

1.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \infty\right) = 1$$

2.

$$\mathbb{P}\left(\sum_{k=1}^{\infty} U_k < \zeta\right) > 0 \iff \zeta > \frac{\theta}{1 - \theta}$$

## Tightness of analysis

---

## Almost zero gap

Let  $D_k = \{d_1, \dots, d_m\}$ , where  $d_i \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(\mathcal{S}^{n-1})$ .

Recall that PDS is convergent if

$$m > \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right).$$

# Almost zero gap

Let  $D_k = \{d_1, \dots, d_m\}$ , where  $d_i \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(\mathcal{S}^{n-1})$ .

Recall that PDS is convergent if

$$m > \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right).$$

With our non-convergence analysis, PDS is non-convergent if

$$m < \log_2 \left( 1 - \frac{\log \theta}{\log \gamma} \right).$$

## Tightness of assumption

Natural question:

$$\mathbb{P}(\text{cm}(D_k, -g_k) \leq 0 \mid D_0, \dots, D_{k-1}) \geq q \not\geq q_0,$$

## Tightness of assumption

Natural question:

$$\mathbb{P}(\text{cm}(D_k, -g_k) \leq 0 \mid D_0, \dots, D_{k-1}) \geq q \not\geq q_0,$$

Answer: NO

# Tightness of assumption

Natural question:

$$\mathbb{P}(\text{cm}(D_k, -g_k) \leq 0 \mid D_0, \dots, D_{k-1}) \geq q \not\geq q_0,$$

Answer: NO

## Example

We assume

- $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be gradient Lipschitz and strongly convex,
- $\theta = 1/2$  and  $\gamma = 2$ ,  $\Rightarrow q_0 = 1/2$
- $D_k = \{g_k/\|g_k\|\}$  or  $D_k = \{-g_k/\|g_k\|\}$  with probability  $1/2$ , respectively,

then we have

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} \|g_k\| = 0\right) = 1.$$

## Conclusions

---



In this talk

- Non-convergence analysis for probabilistic direct search
- Tightness of non-convergence analysis

In this talk

- Non-convergence analysis for probabilistic direct search
- Tightness of non-convergence analysis

Future work

- Estimate the non-convergence probability
- Conduct non-convergence analysis for other frameworks

Thank you!

# References I

- ▶ Biviano, A. et al. (2013). “CLASH-VLT: the mass, velocity-anisotropy, and pseudo-phase-space density profiles of the  $z = 0.44$  galaxy cluster MACS J1206.2-0847”. *A&A* 558, A1:1–A1:22.
- ▶ Conn, A. R., Scheinberg, K., and Vicente, L. N. (2009). *Introduction to Derivative-Free Optimization*. Vol. 8. MOS-SIAM Ser. Optim. Philadelphia: SIAM.
- ▶ Durrett, R. (2010). *Probability: Theory and Examples*. Fourth. Camb. Ser. Stat. Probab. Math. Cambridge: Cambridge University Press.
- ▶ Fermi, E. and Metropolis, N. (1952). *Numerical solution of a minimum problem*. Tech. rep. Alamos National Laboratory, Los Alamos, USA.

## References II

- ▶ Ghanbari, H. and Scheinberg, K. (2017). “Black-box optimization in machine learning with trust region based derivative free algorithm”. *arXiv:1703.06925*.
- ▶ Gratton, S. et al. (2015). “Direct search based on probabilistic descent”. *SIAM J. Optim.* 25, pp. 1515–1541.