Master thesis

Work Distribution of a Heterogeneous Library Staff - A Personnel Task Scheduling Problem

Claes Arvidson, Emelie Karlsson Lith - Mat - EX - - 04 / 04 - - SE

Work Distribution of a Heterogeneous Library Staff - A Personnel Task Scheduling Problem

Optimeringslära, Linköpings Universitet

Claes Arvidson, Emelie Karlsson

LiTH - MAT - EX - - 04 / 04 - - SE

Exam work: 30 hp

Level: ${\bf A}$

Supervisor: T. Larsson,

Optimeringslära, Linköpings Universitet

Examiner: E. Rönnberg,

Optimeringslära, Linköpings Universitet

Linköping: June 2016

Abstract

Here is where you can write your abstract. It may be very long, or it may be very short, the reason you have an abstract is for people not to be forced to read lots of crap.

But still, they will have to read your abstract. After all, the abstract is what everyone reads. . .

Keywords: Keyword One, Chemostat, Another Key-Word, Key, Clé, Mot de cle, Nyckelhål, XBOX, Dagens viktigaste nyckelord, and Keywords.

URL for electronic version:

http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-77777

Acknowledgements

I would like to thank my supervisor, I would like to thank my supervisor, I would like to thank my supervisor...

I also have to thank, I would like to thank my supervisor, I would like to thank my supervisor, I would like to thank my supervisor...

My opponent NN also deserves my thanks, I would like to thank my supervisor, I would like to thank my supervisor. I would like to thank my supervisor. . . .

Nomenclature

Most of the reoccurring definitions, symbols and abbreviations are described here.

Definitions

Plocklista Text Library on wheels Text Rostering Text

Symbols

- Y_0 The amount of the variable Y inserted into a system.
- \hat{Y} The unit-dimension of the variable Y, for example $\hat{t} = 1s$.
- \bar{Y}_i A steady state (number i) value of Y.
- K_i Constants used in kinetic expressions, for example K_I .
- **A** The system matrix.

Abbreviations

 $\begin{array}{ccc} \operatorname{Exp} & \operatorname{Text} \\ \operatorname{Info} & \operatorname{Text} \\ \operatorname{PL} & \operatorname{Text} \\ \operatorname{PTSP} & \operatorname{Text} \\ \operatorname{SMPTSP} & \operatorname{Text} \end{array}$

CPI Competitive Product Inhibition (or Inhibited)
CSI Competitive Substrate Inhibition (or Inhibited)

CSTR Continuous Stirred Tank (bio)Reactor MMI Michaelis-Menten Inhibition (or Inhibited)

Contents

1	Intr	oduction	1
•	1.1	Background	1
	1.2	Problem description	1
	1.2	1.2.1 Description of the daily tasks at the library	1
		1.2.2 Personnel attributes	2
		1.2.3 Main objective: increase number of stand in personnel	3
		1.2.4 Secondary objectives: repetitiveness of the schedule	3
	1.3	Method	3
	1.4	Topics Covered	3
2	Lite	rature review	5
	2.1	Personnel Task Scheduling Problem	5
		2.1.1 Applications	7
	2.2	Shift Minimisation Personnel Task Scheduling Problem	7
	2.3	Tour Scheduling Problem	8
	2.4	Other similar problems	9
		2.4.1 Fixed Job Schedule Problem $\dots \dots \dots \dots$	9
		2.4.2 Tactical Fixed Interval Scheduling Problem	10
		2.4.3 Operational Fixed Interval Scheduling Problem	10
		2.4.4 Stochastic job problems	10
	2.5	Problems with soft constraints	11
	2.6	Summary	12
	2.7	Relevance to our problem	12
3	The	mathematical model	13
	3.1	Set and variable definitions	13
	3.2	Objective function	13
	3.3	Constraints	14
4		ideal CSTR: the chemostat	15
	4.1	Some simple models of biological growth	15
		4.1.1 Exponential growth	15
		4.1.2 The logistic equation	15
	4.2	The chemostat	15
A	Set	and variable list	19
В	Para	ameter list	21

xii	Contents
•	

\mathbf{C}	The	Linearized stability	23
	C.1	The Linearization	. 23

List of Figures

List of Tables

1.1	Outer tasks can be performed exclusively by librarians or by as-
	sistants also
1.2	Demand of staff for the three daily tasks
1.3	Demand of staff at Hageby and Library on Wheels
2.1	PTSP variants
3.1	Significant sets and variables

xiv List of Tables

Chapter 1

Introduction

1.1 Background

At a library absence can cause problems, both due to lack of personnel as well as due to the qualifications required to perform a task varies. If a worker were to be unavailable a day because of a meeting or being ill it would require for a stand-in to fill the vacancy. Therefore, it is of great interest to have a schedule with as many skilled stand-ins as possible to overcome such disturbances. Furthermore, the library personnel have certain demands and preferences as to how a satisfactory working schedule should be. For instance, it is neither preferable to work more than one evening each week nor work more weekends than required.

The central library of Norrköping is more than 100 years old. The library has more than 40 employees. These consist of both librarians and assistants, who handle simpler tasks. The library is open both weekdays until 20:00 and during weekend days until 16, which is challenge for the schedulers of the library since this requires time compensation for the staff. The library also provides its services to more than five other smaller libraries.

1.2 Problem description

1.2.1 Description of the daily tasks at the library

The most important activity at a library is the activity directed towards the public. This includes lending books services as well as providing customers with helpful information about the resources at their disposal. These are referred to as "outer tasks". At the same time, the uppsättning of books must be maintained, the returned books must be sorted and put back, the web page must be up to date and so on. Such work is often referred to as "inner work" and is equally part of the everyday tasks of a librarian.

At the library of Norrköping, three main outer tasks can be identified as working in the service counter (sv. expiditionsdisken), working in the information counter (sv. informationsdiken) and assembling books according to the "fetch list" (sv. plocklista). These tasks can be performed by either librarians or assistants, as descirbed in able 1.1.

Table 1.1: Outer	tasks can	be performed	exclusively	y by	librarians or	by assis-
------------------	-----------	--------------	-------------	------	---------------	-----------

tants also.

Task	Description	Qualification
Service Counter	Administring loans, library cards	Assistants, Librar-
(Exp)	and the loaning machine	ians
Information	Handling questions	Librarians
Counter (Info)	about the library's resources.	
Fetch List (FL)	Fetching books that are to be	Assistants, Librar-
	sent to other libraries.	ians

As the number of visitors in the library differs at different times of the day and during different days so does the demand for people for the three tasks. The demand of people at for the different tasks is illustrated in table 1.2.

Table 1.2: Demand of staff for the three daily tasks.

Day	Time	Exp demand	Info demand	FL demand
Mon-Fri	08:00-10:00	2	2	1
Mon-Fri	10:00-13:00	3	3	1
Mon-Fri	13:00-16:00	3	3	1
Mon-Fri	16:00-20:00	3	3	-
Sat	11:00-16:00	3	3	-
Sun	11:00-16:00	3	3	-

As is the case with most libraries, the Central Library of Norrköping also has responsibilities that fall outside of it's normal daily activities. One such resposibility is the running of a smaller library filial in Hageby, situated in a suburbian area of Norrköping, during weekends. For this tasks only librarians are qualified as the placement implies all types of library tasks.

Similarly, only librarians are qualified for the task known as "Library on Wheels" (sv. Bokbussen), which is a type of library bus, providing citizens in remoter areas of the city with books and other library services. The Library on Wheels has only operates a few times a week and the schedule differs between even and odd weeks.

1.2.2 Personnel attributes

The Central Library of Norrköping currently has 39 workers, 23 of which are librarians and 16 of which are assistants. All staff have different availability for performing tasks, depending on their working hours and the amount of inner work they are in charge of.

Personnel works whole weekends!

1.3. Method 3

Table 1.3: Demand of staff at Hageby and Library on Wheels

Day	Time	Hageby	LoW - odd week	LoW - even week
Mon	08:00-10:00	-	1	1
Mon	16:00-20:00	-	1	-
Tue	08:00-10:00	-	-	-
Tue	16:00-20:00	-	-	-
Wed	08:00-10:00	-	1	1
Wed	16:00-20:00	-	1	1
Thu	08:00-10:00	-	1	1
Thu	16:00-20:00	-	1	1
Fri	08:00-10:00	-	1	-
Fri	16:00-20:00	-	-	-
Sat	11:00-16:00	1	-	-
Sun	11:00-16:00	1	-	-

- 1.2.3 Main objective: increase number of stand in personnel
- 1.2.4 Secondary objectives: repetitiveness of the schedule

1.3 Method

1.4 Topics Covered

[&]quot;The thesis is divided into..."

Chapter 2

Literature review

The scheduling problem is a mathematical optimization problem which has been studied since the 1950's and concerns creating a feasible and satisfactory schedule for workers or machines performing tasks. One of the most extensive overviews in the area is provided by Ernst et al. (2004). They state that, although the complexity of the scheduling problem has not increased in recent years, the mathematical models used to solve the scheduling problems have become more realistic and refined. Modern scheduling problems often concern the distribution of tasks and the creation of worker shifts, taking also into account softer values such as worker satisfaction and worker fatigue. Due to this modeling refinement as well as the development of more powerful computational methods, it is possible today to solve scheduling problems in a more satisfactory way than before.

In this section, the scheduling problem is classified into different subcategories which are areas related to the work of this paper. A few relevant areas for our work include Personnel Task Scheduling Problems (PTSP), Shift Minimization Task Scheduling Problems (SMTSP), Tour Scheduling Problems (TSP) and a few variations of these. Models and solutions methods of these problems will be discussed during each topic and a summary of the discussion will be provided.

2.1 Personnel Task Scheduling Problem

In many practical instances production managers will face the Personnel Task Scheduling Problem (PTSP) while scheduling plant operations. It occurs when the rosterer or shift supervisor need to allocate tasks with specified start and end times to available personnel who have the required qualifications. Furthermore, it also occurs in situations where tasks of fixed times shall be assigned to machines. Decisions will then have to be made regarding the amount of maintenance workers needed and which machine the workers are assigned to look after Krishnamoorthy and Ernst (2001).

There are numerous variants to the PTSP. Studies on these have been made in article Krishnamoorthy and Ernst (2001) by Krishnamoorthy et al. who gives a list of attributes that commonly appear in a PTSP and which are listed in Table 2.1 below. There are furthermore traits that always appear in a PTSP;

tasks with fixed start and end time are to be distributed to staff members that possess certain skills, allowing them to perform only a subset of the available tasks. Start and end time of their shifts are also predetermined for each day.

One variant, which also is the most simple, is mentioned in Krishnamoorthy and Ernst (2001) and is called the *Feasibility Problem* where the aim is to just find a feasible solution. This requires that each task is allocated to a qualified and available worker. It is also required that a worker cannot be assigned more than one task simultaneously as well as tasks cannot be pre-empted, meaning that each task has to be completed by one and the same worker.

In Table 2.1 one can see attributes of PTSP variants. The nomenclature of the attributes T, S, Q, O refer to the *Task type*, *Shift type*, *Qualifications* and *Objective function* respectively.

Table 2.1: PTSP variants

Attribute	Type	Explanation
Т	F	Fixed contiguous tasks
	V	Variable task durations
	\mathbf{S}	Split (non-contiguous) tasks
	\mathbf{C}	Changeover times between consecutive tasks
S	F	Fixed, given shift lengths
	I	Identical shifts which are effectively of infinite duration
	D	Maximum duration without given start or end times
	U	Unlimited number of shifts of each type available
Q	I	Identical qualification for all staff (homogeneous workforce)
	${ m H}$	Heterogeneous workforce
O	F	No objective, just find a feasible schedule
	A	Minimise assignment cost
	Τ	Worktime costs including overtime
	W	Minimise number of workers
	U	Minimise unallocated tasks

Many of the most basic problems and a few more complex ones can be described with this definition of PTSP attributes. It is, however, not possible to describe all of the numerous types of PTSP using these nomenclatures Krishnamoorthy and Ernst (2001).

By combining attributes it is possible to obtain more complex variants of the PTSP. An example would be the PTSP[F;F;H;A-T-W] mentioned in Krishnamoorthy and Ernst (2001) where multiple objectives are used. This problem has fixed contiguous tasks, fixed shift lengths, heterogeneous workforce and three objective functions; A-T-W, which represent assignment costs, work time with overtime included and requirements to minimize the number of workers respectively. For this problem the objective function is then a linear combination with different parameters used to prioritize (weigh) them against each other.

Given the nomenclature above, our problem would be most related to the PTSP[F;F;H;F]. The difference is that the objective function is not empty. We are looking to maximize the number of qualified stand-ins each day as well as maximize employee satisfaction by meeting their recommendations. Since we have a fix number of workers, no costs and no unallocated tasks when a feasible solution is found, this cannot be described with the type of objective attributes

given in Table 2.1. Therefore, none of the objective function types are relevant in our case.

Different variants of PTSP are given names in the literature. An example is when the shifts and qualifications are identical (S=I and Q=I) and the objective function is to minimize the number of workers that are used (O=W). This variant, PTSP[F;I;I;W], has been published as the "fixed job schedule problem" and is described in Section 2.4 Krishnamoorthy and Ernst (2001).

2.1.1 Applications

An example where PTSP can be found is when developing a rostering solution for ground personnel at an airport. Such a problem can be dealt with by first assigning the workers to days in order to satisfy all the labour constraints, followed by assigning the tasks to the scheduled workers Krishnamoorthy and Ernst (2001).

Three problems of type PTSP related to airplanes can be found when scheduling for either airport maintenance staff, planes to gates or staff that do not stay in one location, such as airline stewards. Scheduling for airport maintenance staff can lead to either PTSP[F;I;H;U-A] or PTSP[F;I-U;H;W], which are similar problems but are given two different names; Operational Fixed Interval Scheduling Problem and Tactical Fixed Interval Scheduling Problem respectively. These are described further in Section 2.4 Krishnamoorthy and Ernst (2001).

Another application, which has been frequently studied, is classroom assignments and is discussed in Krishnamoorthy and Ernst (2001). Based on specifications such as the amount of students in a class or the duration of a class, different classrooms have to be considered. Requirements of equipment, e.g. for a laboratory, may also greatly limit the available classrooms to choose from. A majority of the complications of this problem is due to the fact that lessons can span over multiple periods.

Worth noting for classroom assignment problems is that there are no start or end times for the shifts, as they represent the rooms. The aim in the present problem would be to simply find a feasible assignment of classrooms. Therefore the nomenclature of the problem would be PTSP[S;I;H;F], with the possibility of adding preferences to the objective function. An example of a preference would be to assign the lessons as close to each other as possible on a day, preventing traveling distances between classes for teachers and students Krishnamoorthy and Ernst (2001).

2.2 Shift Minimisation Personnel Task Scheduling Problem

A close relative to the PTSP is the Shift Minimisation Personnel Task Scheduling Problem (SMPTSP) and is a special case in which the aim is to minimize the cost occurring due to the number of personnel (shifts) that are used. The same common traits are valid in this problem as in the PTSP; workers with fixed work hours are to be assigned tasks, with specified start and end times, that they are qualified for Krishnamoorthy et al. (2012).

In article Krishnamoorthy et al. (2012) they "... concentrate mainly on a variant of the PTSP in which the number of personnel (shifts) required is to be

minimised.". In doing so, it is possible to determine the lowest number and mix of skilled staff a company should have to be able to complete the tasks and still be operational. They also presumed that the pool of workers are unlimited for either skill group, which is not the case in our problem due to the limitations on the amount of librarians and assistants available.

SMPTSP can be applied when there are a large number of workers available with different qualifications and it is needed to ensure that the tasks for that day are performed. The PTSP and SMPTSP are therefore useful day-to-day management tools that commonly occurs in many practical instances where tasks are allocated on a daily basis Krishnamoorthy et al. (2012).

It is shown in Kroon et al. (1995) that SMPTSP is a complex problem even if the preemption constraint were to be removed. However, if the qualifications of the workers were identical it would become an easily solvable problem Krishnamoorthy et al. (2012).

SMPTSP is almost identical to another problem introduced by Kroon et al. which is called the Tactical Fixed Interval Scheduling Problem and is described in Section 2.4 below Krishnamoorthy et al. (2012).

2.3 Tour Scheduling Problem

The Tour Scheduling Problem (TSP) is described by Loucks and Jacobs (1991) as a combination of shift scheduling and days-off scheduling. Using the notation used described in the section 2.1, this would be be classified as a variant of the PTSP[F;D;I;W] or PTSP[F;D;H;W]. Shift scheduling refers to creating sets of contiguous hours during which a worker is assigned for work. The need for days off scheduling typically occurs when the time horizon for scheduling is weekly or more and when weekend staff is needed.

According to Loucks and Jacobs (1991), the vast majority of all tour scheduling problems up to 1991 involved a homogeneous workforce, that is, any worker can perform any assigned task. One such early study of the our scheduling problem often mentioned in literature is provided by Roberts and Escudero (1983a). The problem studied in this PhD thesis concern only homogeneous work forces and the task assignment part is lacking.

In the article by Loucks and Jacobs, the authors study a tour scheduling problem with a heterogeneous work force. The problem both involves tour scheduling and task assignment, where the latter part is most interesting to us. The problem is studied in the context of fast food restaurants, where certain personnel is qualified only for certain stations in the restaurant. In such industries, the demand of staff differs between different weekdays and different times of the day. Two worker attributes are considered; their availability for work and their qualification for performing different tasks. The problem concerns finding shifts for all workers which are to have a length between a minimum and maximum number of hours per day.

The representative problem studied in the article involves creating a one-week schedule for 40 workers in a fast food restaurant, available for eight different tasks with a seven-day, 128-hour workweek. Several synthetic problems are studied in the article, all, however with minimum shift length three hours, maximum shift length eight hours and five maximum number of work days.

A similar problem to the one descibed by Loucks and Jacobs is studied by

Choi et al. (2009). They focus on a particular fast food restaurant in Seoul, which is made a representative of fast food chains in general. In this study, only two types of workers are available; fulltime and part time workers, with no other reference to difference in skill. The different shifts are already given by the reastaurant managers and the task is to combine them into a tour. The task assignment aspect is lacking in this article.

In both articles the main objective is to minimize both overstaffing and understaffing, which will both have economical consequences for the fast food chain. This is done by redusing or increasing the work force. For a problem with a fixed work force, such as ours, this objective is not relevant. In the example studied by Loucks and Jacobs there is also a goal to meet staff demand on total working hours. This is modeled as a secondary goal and is similar to our goal and somehow models a "soft" value, which is of interest to us.

A more recent tour scheduling problems concern monthly tour scheduling, as opposed to most literature which concerns only weekly scheduling. Such a study was done by Rong (2010). The main advantage of monthly scheduling over shorter time periods, as stated in the article, is the possibility to plan a schedule with respect to fairness and balance over a longer period of time. The problem concerns workers with different skills, where each worker also can possess multiple skills. This is referred to as a mixed skill problem. Thus the problem is similar to our problem, where mixed skill is also present. In the study, workers have individual weekend-off requirements. The problem does not involve task assignment, which makes it less relevant for us.

The solution methods used to solve the tour scheduling problem differs greatly between the articles studied. In the older articles, such as Roberts and Escudero (1983a) and Loucks and Jacobs (1991), custom made algorithms very similar to the methods used in manual scheduling are proposed to solve the problem. These solution methods involve classifying staff and distributing them according to some rule (for example, the staff with the most scarce skill is assigned first). General commercial solvers are not proposed, due to lack of efficiency during these times. In Hojati and Patil (2011), the same model and data is used as in Loucks and Jacobs (1991). Also this article states that commercial solvers are insufficient for solving the problem. Instead two methods are proposed, one which decomposes the problem into two problems solvable by commercial solvers and one customized heuristic method based on the Lagrangian relaxation method. in Choi et al. (2009), a pure integer programming method is used.

2.4 Other similar problems

In this section a couple of problems similar to our own will be described in order to give clarity as to how closely related many of these problem types are.

2.4.1 Fixed Job Schedule Problem

Variations of the task assignment problem relevant for our problem include for example the Fixed Job Schedule Problem (FJSP). The FJSP has been studied since the 1970s in the context of task assignment in processors. The problem concerns the distribution of tasks with fixed starting and ending times over a

workforce with identical skills, such as processing units Krishnamoorthy et al. (2012). Such problems have been solved by Gertsbakh and Stern (1978) and Fischetti et al. (1992).

In the article by Gertsbakh and Stern (1978) a situation where n jobs need to be scheduled over an unlimited number of procesors is studied. The objective function of such a problem becomes to minimize the number of machines needed to perform all tasks. Fischetti solves a similar problem, but adds time constraints, saying that no processor is allowed to work for more than a fixed time T during a day as well as a spread time constraint forcing tasks to spread out with time gap S over a processor.

2.4.2 Tactical Fixed Interval Scheduling Problem

Another type of problem is the Tactical Fixed Interval Scheduling Problem (TFISP). This is a problem very closely related to the SMPTSP problem with the only difference being that the TFISP concerns workers which always are available, such as industrial machines or processors. The problem is studied by for example Kroon et al. Kroon et al. (1995). A typical TFISP can be expressed using the nomenclature in Table 2.1 and written as PTSP[F;I-U;H;W] Krishnamoorthy and Ernst (2001).

As opposed to the FJSP, the TFISP deals with a heterogeneous workforce. Two different contexts are studied by Kroon et al. One of them concerns the handling of arriving aircraft passengers at an airport. Two modes of transport from the aeroplane to the airport are investigated; directly by gate or by bus. The two transportation modes thus correspond to two processing units which can only handle a number of jobs at the same time.

2.4.3 Operational Fixed Interval Scheduling Problem

The Operational Fixed Interval Scheduling Problem (OFISP) is a close relative to TFISP, where both types are restricted by the following; each machine (worker) cannot handle more than one job at a time, each machine can only handle a subset of the jobs and preemption is not allowed. The difference between them occurs in the objective function, as TFISP tries to minimize the number of workers while OFISP tries to minimize the operational costs and the number of unallocated tasks Kroon et al. (1995). In the present nomenclature this would give rise to the problem PTSP[F;I;H;U-A] Krishnamoorthy and Ernst (2001). Given the problem definition above, working shifts are to be created for the workers and tasks are to be allocated on a day-to-day basis. OFISP can therefore be seen both as a job scheduling problem and a task assignment problem Kroon et al. (1995).

2.4.4 Stochastic job problems

What differs mostly between the problem types described above and the problem studied in this thesis work, is the difference in objective. The main objective is often to minimize staff for a number of fixed jobs, not taking into consideration the need for stand ins to perform critical tasks. An area where the need for stand in personnel appears is in the maintenance industry, where some jobs can be forseen and other (emergency) jobs are of a stochastic nature, that is, there is a

probability that such jobs will occur a certain hour. The problem combing both unplanned and planned maintenance worker scheduling was studied by Duffuaa and Al-Sultan (1999), as a continuation of Roberts and Escudero (1983b).

In the article, a fixed, heterogeneous work force consisting of electricians, plumbers and mechanics is studied. The shifts of the personnel are predetermined by their given work times and thus the problem becomes a pure task assignment problem. An objective function is used where the goal is to maximize the number of planned and unplanned jobs performed by the workers, by taking into account the probability of unplanned work to occur. Thus, certain workers will be left at the station as stand-ins in the case an unplanned job arrives.

2.5 Problems with soft constraints

For most scheduling problems, the main objective is to minimize worker-related costs by reducing the number of workers needed to perform a task, or by reducing the working time for part-time employees. Equivalently, the goal in production industries is to reduce the number of machines needed. Recently, however, many studies have started to focus more or softer values such as worker satisfaction as an objective. Such values are usually considered when scheduling is done manually, but have been forgotten or set aside in mathematical modeling.

In an article by Akbari et al. (2013) a scheduling problem for part-time workers with different preferences, seniority level and productivity is investigated. In this article, these aspects are reflected in the objective function and weighted against each other. A similar problem was also studied by Mohan (2008), but for a work force of only part-time workers .

Other factors which may affect worker satisfaction, and in the long run efficiency and presence at work are fatigue, fairness and boredom. These are discussed by Eiselt and Marianov (2008). Repetitiveness of a job as well as the level of challenge can cause bordom in workers. Increasing variance is done by Eiselt and Marianov (2008) through providing an upper bound of how many tasks can be performed in a given time span. The article suggests some sort of measurement of the distance between the task requirements and the worker abilities is used. This will then be minimized in the objective function.

Another modeling method which is relevant specifically for scheduling problems featuring soft constraints is fuzzy goal programming. The method is discussed by Shahnazari-Shahrezaei et al. (2013), who model soft constraints as "fuzzy goals". These goal can become contradictory, for example could a preference of high seniority level workers come in conflict with a preference in working hours by an employee. The article uses fuzzy set theory, and uses a solution approach involving Li's two-phase method, Lee (1989). Soft constraints are modeled as trapezoid functions and an optimal solution which finds the best average of all of them is found.

The solution methods proposed by Akbari et al. (2013) to solve a scheduling problem with soft constraints are two matheuristics; Simulated Annealing (SA) and Variable Neighbourhood Search (VNS). According to Akbari et al. (2013), SA has been studied as a solution method for the scheduling problem since the 1990's and many studies have shown that it is capable of providing near-optimal solutions in a short time compared to optimal integer-programing models for a

variety of problems. The method is a random meta heuristic, which finds new random solutions and chooses it to be the current solution if sufficiently good. With time, the threshold for what is considered good is increasing.

VNS is the other proposed method by Akbari et al. (2013). A big difference between this and other methods is that VNS requires very little parameter tuning while often providing good solutions. The method uses involves a random and a systematic phase. In the random phase, new shifts are generated randomly and better solutions are saved. In the systematic phase, two shifts are swapped and again better solutions are saved.

In the article by Eiselt and Marianov (2008), a commercial solver was used. This was also the case in Mohan (2008), although the article compares these results with the results obtained from a branch-ans-cut algorithm. Also Shahnazari-Shahrezaei et al. (2013) uses a commercial solver.

2.6 Summary

Stycke ett: Modeller Stycke två: Metoder historiskt kronologiskt

2.7 Relevance to our problem

Kan börjas med syftesformulering. Referera tillbaka i texten, samt lägg till nya reflektioner.

Chapter 3

The mathematical model

In this chapter the mathematical model implemented to solve this problem will be presented. Variables and sets will be stated prior to the objective function and constraints. A list of all the parameters can be found in Appendix B.

3.1 Set and variable definitions

To solve the problem many sets and definitions had to be declared as there are many unique requirements that have to be met. An example is that some want a day free from outer tasks so that they can focus on other assignments or attend meetings. Another one is that some personnel have different schedules for odd and even weeks. These specific cases have to be modeled and result in a variety of set and variable definitions. Hence, a bigger part of the definitions are instead listed in Appendix A.

Table 3.1: Significant sets and variables

I	Set of workers
W	Set of weeks
D	Set of days in a week
S_d	Set of shifts day d
J_d	Set of task types day d
x_{iwdsj}	1 if worker i is assigned task j in week w , day d , shift s , 0 otherwise
$h1_{iw}$	1 if worker i works its first weekend week w , 0 otherwise
$h2_{iw}$	1 if worker i works its second weekend week w , 0 otherwise

3.2 Objective function

Due to multiple objective functions, it has been necessary to weigh them against each other to prioritize using parameters. As can be seen in equation ... below

$$maxN1l*lowest_stand_in_amount_lib + N1a*lowest_stand_in_amount_ass$$
 (3.1)

$$\begin{split} maxN1l*lowest_stand_in_amount_lib \\ + N1a*lowest_stand_in_amount_ass \\ - N2*\sum_{n=1}^{\infty} 2^{-n} = 1 \end{split}$$

3.3 Constraints

To model this problem it has been of relevance to divide many of the constraints into weekend- and weekday constraints.

Chapter 4

The ideal CSTR: the chemostat

In this chapter we study exponential growth, the logistic. . . .

4.1 Some simple models of biological growth

4.1.1 Exponential growth

If $\mu = \text{constant} > 0$, we get $X(t) = X_0 e^{\mu t}$.

4.1.2 The logistic equation

Let us assume that $\frac{dX}{dt} = \mu \cdot X$, with $\mu = \mu(S) = k \cdot S \dots$

$$\begin{cases} \frac{dX}{dt} = kSX & (a) \\ \frac{dS}{dt} = -\alpha kSX & (b) \end{cases}$$

$$\frac{dX}{dt} = r(1 - \frac{X}{B})X \tag{4.1}$$

An explicit solution to (4.1) is: $X(t) = \frac{X_0B}{X_0 + (B - X_0)e^{-rt}}$, if $0 < X_0 < B$. It can be found by separating variables in equation (4.1)

4.2 The chemostat

A chemostat is made of two main parts; a nutrient reservoir, and a growth-chamber, reactor, in which the bacteria reproduces.

$$\begin{cases}
\frac{dX}{dt} = \mu(S)X - X \frac{F}{V} \\
\frac{dS}{dt} = -\alpha\mu(S)X - S \frac{F}{V} + S_0 \frac{F}{V}
\end{cases}$$
(4.2)

$$\mathbf{A} = \left(\begin{array}{cc} 0 & \sigma \alpha_1 \\ -\frac{1}{\alpha_1} & -\sigma - 1 \end{array} \right)$$

The invariant line: conclusions

Model	Monods Chemostat	CSI-CSTR
μ	$\frac{S}{1+S}$	$\frac{S}{1+S+\frac{S^2}{K_I}}$
$\frac{dX}{dt}$	$\alpha_1 \frac{S}{1+S} X - X$	$\alpha_1 \frac{S}{1 + S + \frac{S^2}{K_I}} X - X$
$\frac{dS}{dt}$	$-\frac{S}{1+S}X - S + \alpha_2$	$-\frac{S}{1+S+\frac{S^2}{K_I}}X - S + \alpha_2$
XNC	$S = \frac{1}{\alpha_1 - 1}$	$S = \frac{K_I(\alpha_1 - 1)}{2} \pm \sqrt{\left(\frac{K_I(\alpha_1 - 1)}{2}\right)^2 - K_I}$
SNC	$X = \frac{(\alpha_2 - S)(1+S)}{S}$	$X = \frac{(\alpha_2 - S)(1 + S + \frac{S^2}{K_I})}{S}$
limit	_	$K_I o \infty$

The other three models, the chemostat, the MMI-CSTR and the CPI-CSTR are quite similar in comparison to the CSI-CSTR.

Monods chemostat does not "feel" this inhibition and does not care...

This document is an example of BibTeX using in bibliography management. Three items are cited: *The LaTeX Companion* book?, the Einstein journal paper?, and the Donald Knuth's website?. The LaTeX related items are??.

Bibliography

- Andreas T Ernst, Houyuan Jiang, Mohan Krishnamoorthy, Bowie Owens, and David Sier. An annotated bibliography of personnel scheduling and rostering. *Annals of Operations Research*, 127(1-4):21–144, 2004.
- Mohan Krishnamoorthy and Andreas T Ernst. The personnel task scheduling problem. In *Optimization methods and applications*, pages 343–368. Springer, 2001.
- Mohan Krishnamoorthy, Andreas T Ernst, and Davaa Baatar. Algorithms for large scale shift minimisation personnel task scheduling problems. *European Journal of Operational Research*, 219(1):34–48, 2012.
- Leo G Kroon, Marc Salomon, and Luk N Van Wassenhove. Exact and approximation algorithms for the operational fixed interval scheduling problem. European Journal of Operational Research, 82(1):190–205, 1995.
- John S Loucks and F Robert Jacobs. Tour scheduling and task assignment of a heterogeneous work force: A heuristic approach. *Decision Sciences*, 22(4): 719–738, 1991.
- SM Roberts and LF Escudero. Scheduling of plant maintenance personnel. Journal of Optimization theory and Applications, 39(3):323–343, 1983a.
- Kyuwan Choi, JOHyE HWaNG, and Myoungju Park. Scheduling restaurant workers to minimize labor cost and meet service standards. *Cornell Hospitality Quarterly*, 50(2):155–167, 2009.
- Aiying Rong. Monthly tour scheduling models with mixed skills considering weekend off requirements. Computers & Industrial Engineering, 59(2):334–343, 2010.
- Mehran Hojati and Ashok S Patil. An integer linear programming-based heuristic for scheduling heterogeneous, part-time service employees. *European Journal of Operational Research*, 209(1):37–50, 2011.
- Ilya Gertsbakh and Helman I Stern. Minimal resources for fixed and variable job schedules. *Operations Research*, 26(1):68–85, 1978.
- Matteo Fischetti, Silvano Martello, and Paolo Toth. Approximation algorithms for fixed job schedule problems. *Operations Research*, 40(1-supplement-1): S96–S108, 1992.

18 Bibliography

Salih O Duffuaa and KS Al-Sultan. A stochastic programming model for scheduling maintenance personnel. *Applied Mathematical Modelling*, 23(5): 385–397, 1999.

- SM Roberts and LF Escudero. Scheduling of plant maintenance personnel. Journal of Optimization theory and Applications, 39(3):323–343, 1983b.
- Mohammad Akbari, M Zandieh, and Behrouz Dorri. Scheduling part-time and mixed-skilled workers to maximize employee satisfaction. *The International Journal of Advanced Manufacturing Technology*, 64(5-8):1017–1027, 2013.
- Srimathy Mohan. Scheduling part-time personnel with availability restrictions and preferences to maximize employee satisfaction. *Mathematical and Computer Modelling*, 48(11):1806–1813, 2008.
- Horst A Eiselt and Vladimir Marianov. Employee positioning and workload allocation. *Computers & operations research*, 35(2):513–524, 2008.
- Parisa Shahnazari-Shahrezaei, Reza Tavakkoli-Moghaddam, and Hamed Kazemipoor. Solving a multi-objective multi-skilled manpower scheduling model by a fuzzy goal programming approach. *Applied Mathematical Modelling*, 37(7):5424–5443, 2013.
- Bum-Il Lee. Multiple objective decision making in a fuzzy environment. 1989.

Appendix A

Set and variable list

 $D=1\ 2\ 3\ 4\ 5\ 6\ 7$

• • •

Appendix B

Parameter list

 $\begin{array}{c} A=1 \\ B=2 \end{array}$

Appendix C

The Linearized stability

C.1 The Linearization

F(x), a one-variable function of x can be Taylor-expanded around a fix X. We get $F(X+x)=F(X)+F'(X)x+O(x^2)$. For small perturbations of x around X we get the linearization: $F(X+x)\approx F(X)+F'(X)x$, containing only the constant and the linear terms.

For functions of two variables F(X + x, S + s) and G(X + x, S + s):

```
 \left\{ \begin{array}{l} F(X+x,S+s) = F(X,S) + F_X'(X,S)x + F_S'(X,S)s + O((x+s)^2) \\ G(X+x,S+s) = G(X,S) + G_X'(X,S)x + G_S'(X,S)s + O((x+s)^2) \end{array} \right.
```

```
function chemostat_inhibited(alpha1, alpha2, xp0, sp0, xc)
%chemostat_inhibited Displays a phaseportrait, nullclines
    and an Euler-path of an inhibited Chemostat.
    chemostat_inhibited(alfa1, alfa2, np0, cp0, nc) will run if
    alpha1 > 1/xc, thus there is a reproduction.
    alpha2 > 1/(xc*alpha1-1), thus there is sufficient stock-nutrition. xp0 > 0 , you can not have a nonpositive population.
    sp0 > 0 , you can not have a nonpositive concentration.
    The blue arrows represent the vectorfield.
    The black lines are two of the three nullclines.
    The black dotted line is the invariant line (no solution crosses it). The red line is an Eulerpath, starting in + and ending in \ast.
    chemostat_inhibited(5, 3, 0.2, 0.3, 6)
    by Per Erik Strandberg, 2003-2004.
% Start-condition:
if ((alpha1>1) & (alpha2>0) & (sp0>0) & (xp0>0) & xc>0),
    if (alpha2<1/(alpha1-1)),
         disp(' ')
disp(' (HINT: Only the trivial steady state, alpha2 is too small...)')
         disp (' (HINT: Two steady states, alpha2 is quite large...)')
```

```
% The illegal indata case:
%-----
else
    disp(' chemostat_inhibited.m by Per Erik Strandberg, 2003-2004.')
    disp(' Did not Finish OK. (You used illegal indata.)')
    disp(' ')
    disp(' For syntax help type: help chemostat_inhibited .')
    disp(' ')
end
```



Copyright

The publishers will keep this document online on the Internet - or its possible replacement - for a period of 25 years from the date of publication barring exceptional circumstances. The online availability of the document implies a permanent permission for anyone to read, to download, to print out single copies for your own use and to use it unchanged for any non-commercial research and educational purpose. Subsequent transfers of copyright cannot revoke this permission. All other uses of the document are conditional on the consent of the copyright owner. The publisher has taken technical and administrative measures to assure authenticity, security and accessibility. According to intellectual property law the author has the right to be mentioned when his/her work is accessed as described above and to be protected against infringement. For additional information about the Linköping University Electronic Press and its procedures for publication and for assurance of document integrity, please refer to its WWW home page: http://www.ep.liu.se/

Upphovsrätt

Detta dokument hålls tillgängligt på Internet - eller dess framtida ersättare - under 25 år från publiceringsdatum under förutsättning att inga extraordinära omständigheter uppstår. Tillgång till dokumentet innebär tillstånd för var och en att läsa, ladda ner, skriva ut enstaka kopior för enskilt bruk och att använda det oförändrat för ickekommersiell forskning och för undervisning. Överföring av upphovsrätten vid en senare tidpunkt kan inte upphäva detta tillstånd. All annan användning av dokumentet kräver upphovsmannens medgivande. För att garantera äktheten, säkerheten och tillgängligheten finns det lösningar av teknisk och administrativ art. Upphovsmannens ideella rätt innefattar rätt att bli nämnd som upphovsman i den omfattning som god sed kräver vid användning av dokumentet på ovan beskrivna sätt samt skydd mot att dokumentet ändras eller presenteras i sådan form eller i sådant sammanhang som är kränkande för upphovsmannens litterära eller konstnärliga anseende eller egenart. För ytterligare information om Linköping University Electronic Press se förlagets hemsida http://www.ep.liu.se/

© 2016, Claes Arvidson, Emelie Karlsson