

# Trace Equivalence in Abstract GSOS

Oberseminar des Lehrstuhls für Theoretische Informatik

---

**Robin Jourde**<sup>1</sup>, Stelios Tsampas, Sergey Goncharov, Henning Urbat, Pouya Partow, Jonas Forster

14th January 2025

<sup>1</sup>ENS de Lyon – robin.jourde@ens-lyon.fr

# Outline

## 1. Preliminaries

### 1.1 GSOS

### 1.2 **Abstract** GSOS

### 1.3 Trace & Kleisli categories

### 1.4 Trace-GSOS

### 1.5 Trace equivalence & congruence

### 1.6 Strong and affine monads

## 2. Result

### 2.1 The theorem

### 2.2 Sketch of the proof

### 2.3 Focus on hypothesis : Smoothness

### 2.4 Focus on hypothesis : Affineness

### 2.5 And for non affine monads ?

## 3. Conclusion

# 1. Preliminaries

---

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$



## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t \rightsquigarrow \Sigma X = 1 + A \times X + X^2 + X$

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t \rightsquigarrow \Sigma X = 1 + A \times X + X^2 + X$

Eg. for  $a, b, c, \tau \in A : a.(b.a.0 + ?\tau.a.c.0)$

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t \rightsquigarrow \Sigma X = 1 + A \times X + X^2 + X$

Eg. for  $a, b, c, \tau \in A : a.(b.a.0 + ?\tau.a.c.0) = a(ba + ?\tau ac)$

## 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t \rightsquigarrow \Sigma X = 1 + A \times X + X^2 + X$

Eg. for  $a, b, c, \tau \in A$  :  $a.(b.a.0 + ?\tau.a.c.0) = a(ba + ?\tau ac)$

- **behaviour** (with endofunctor  $H$ )

# 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \mid \forall a \in A \mid t + t \mid ?t \rightsquigarrow \Sigma X = 1 + A \times X + X^2 + X$

Eg. for  $a, b, c, \tau \in A$  :  $a.(b.a.0 + ?\tau.a.c.0) = a(ba + ?\tau ac)$

- **behaviour** (with endofunctor  $H$ )

*Example:*  $x$  **terminates** ( $x \downarrow$ ) or **progresses** to  $x'$  with label  $a \in A$  ( $x \xrightarrow{a} x'$ )

# 1.1 GSOS

- a **framework** for specifying reduction rules and semantics  
→ rule format
- given a **syntax** (with endofunctor  $\Sigma$ )

*Example:* Set of operations  $\mathcal{O}$  with arity map  $\text{ar} : \mathcal{O} \rightarrow \mathbb{N}$ ,  $\Sigma X = \sum_{\sigma \in \mathcal{O}} X^{\text{ar } \sigma}$

*Example:*  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t \rightsquigarrow \Sigma X = 1 + A \times X + X^2 + X$

Eg. for  $a, b, c, \tau \in A$  :  $a.(b.a.0 + ?\tau.a.c.0) = a(ba + ?\tau ac)$

- **behaviour** (with endofunctor  $H$ )

*Example:*  $x$  **terminates** ( $x \downarrow$ ) or **progresses** to  $x'$  with label  $a \in A$  ( $x \xrightarrow{a} x'$ )  $\rightsquigarrow$   
 $HX = \mathcal{P}(1 + A \times X)$

## 1.1 GSOS

- $k : X \rightarrow HX$  a ***H-coalgebra***  $\longrightarrow$  set equipped with semantics

## 1.1 GSOS

- $k : X \rightarrow HX$  a  **$H$ -coalgebra**  $\longrightarrow$  set equipped with semantics

*Example:* Let  $k : X \rightarrow HX$ , for  $x \in X$ ,  $x \downarrow \Leftrightarrow * \in k(x)$  and  $x \xrightarrow{a} x' \Leftrightarrow (a, x') \in k(x)$



# 1.1 GSOS

- $k : X \rightarrow HX$  a **H-coalgebra**  $\longrightarrow$  set equipped with semantics

*Example:* Let  $k : X \rightarrow HX$ , for  $x \in X$ ,  $x \downarrow \Leftrightarrow * \in k(x)$  and  $x \xrightarrow{a} x' \Leftrightarrow (a, x') \in k(x)$

- **GSOS rules**

$$\frac{\left\{ x_i \xrightarrow{a_{i,k}} y_{i,k} \right\}_{i \in I, k \in K_i} \quad \left\{ x_j \downarrow \right\}_{j \in J}}{\sigma(x_1 \dots x_n) \xrightarrow{b} u[x_1 \dots x_n, y_{i,k} \dots]} \quad \text{or} \quad \frac{\left\{ x_i \xrightarrow{a_{i,k}} y_{i,k} \right\}_{i \in I, k \in K_i} \quad \left\{ x_j \downarrow \right\}_{j \in J}}{\sigma(x_1 \dots x_n) \downarrow}$$

with  $\sigma \in \mathcal{O}$ ,  $n = \text{ar } \sigma$ ,  $u \in \Sigma^*$ ,  $a_{i,k}, b \in A$ ,  $I, J, K_i \subset \llbracket 1, n \rrbracket$

# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$

# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules

# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules

$$\frac{}{0 \downarrow}$$

# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules

$$\frac{}{0 \downarrow} \quad \frac{}{a.t \xrightarrow{a} t} \forall a$$

# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules

$$\begin{array}{c} \frac{}{0 \downarrow} \quad \frac{}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a \quad \frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a \quad \frac{t \downarrow \quad u \downarrow}{t + u \downarrow} \end{array}$$

# A full example


- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules

$$\begin{array}{c} \frac{}{0 \downarrow} \quad \frac{}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a \quad \frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a \quad \frac{t \downarrow \quad u \downarrow}{t + u \downarrow} \quad \frac{t \xrightarrow{\tau} t' \quad t \xrightarrow{\tau} t''}{?t \xrightarrow{\tau} t' + t''} \end{array}$$

# A full example

- syntax:  $t ::= 0 \mid a.t \mid \forall a \in A \mid t + t \mid ?t$
- rules

$$\begin{array}{c}
 \frac{}{0 \downarrow} \quad \frac{}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a \quad \frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a \quad \frac{t \downarrow \quad u \downarrow}{t + u \downarrow} \quad \frac{t \xrightarrow{\tau} t' \quad t \xrightarrow{\tau} t''}{?t \xrightarrow{\tau} t' + t''}
 \end{array}$$







# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules




$$\begin{array}{c}
 \frac{}{0 \downarrow} \quad \frac{}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a \quad \frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a \quad \frac{t \downarrow \quad u \downarrow}{t + u \downarrow} \quad \frac{t \xrightarrow{\tau} t' \quad t \xrightarrow{\tau} t''}{?t \xrightarrow{\tau} t' + t''}
 \end{array}$$



# A full example

- syntax:  $t ::= 0 \mid a.t \ \forall a \in A \mid t + t \mid ?t$
- rules

$$\begin{array}{c}
 \frac{}{0 \downarrow} \quad \frac{}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a \quad \frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a \quad \frac{t \downarrow \quad u \downarrow}{t + u \downarrow} \quad \frac{t \xrightarrow{\tau} t' \quad t \xrightarrow{\tau} t''}{?t \xrightarrow{\tau} t' + t''}
 \end{array}$$



## 1.2 **Abstract** GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$

## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^* X$

## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^*X$

*Example:* For the previous example without  $?$  :  $\Sigma X = 1 + A \times X + X^2$ ,

$$\rho : 1 + A \times (X \times \mathcal{P}(1 + A \times X)) + (X \times \mathcal{P}(1 + A \times X))^2 \rightarrow \mathcal{P}(1 + A \times \Sigma^*X)$$

## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^*X$

*Example:* For the previous example without  $?$  :  $\Sigma X = 1 + A \times X + X^2$ ,

$$\rho : 1 + A \times (X \times \mathcal{P}(1 + A \times X)) + (X \times \mathcal{P}(1 + A \times X))^2 \rightarrow \mathcal{P}(1 + A \times \Sigma^*X)$$

- $\rho(*) = \{*\}$

$\overline{\phantom{x}}$   
 $0 \downarrow$

## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^*X$

*Example:* For the previous example without  $?$  :  $\Sigma X = 1 + A \times X + X^2$ ,

$$\rho : 1 + A \times (X \times \mathcal{P}(1 + A \times X)) + (X \times \mathcal{P}(1 + A \times X))^2 \rightarrow \mathcal{P}(1 + A \times \Sigma^*X)$$

- $\rho(*) = \{*\}$
- $\rho((a, t, T)) = \{(a, t)\}$

$$\frac{}{a.t \xrightarrow{a} t} \forall a$$

## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^*X$

*Example:* For the previous example without  $?$  :  $\Sigma X = 1 + A \times X + X^2$ ,

$$\rho : 1 + A \times (X \times \mathcal{P}(1 + A \times X)) + (X \times \mathcal{P}(1 + A \times X))^2 \rightarrow \mathcal{P}(1 + A \times \Sigma^*X)$$

- $\rho(*) = \{*\}$
- $\rho((a, t, T)) = \{(a, t)\}$
- $\rho((t, T), (u, U)) = \{(a, t') \mid \forall (a, t') \in T\} \cup$

$$\frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a$$



## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^* X$

*Example:* For the previous example without  $?$  :  $\Sigma X = 1 + A \times X + X^2$ ,

$$\rho : 1 + A \times (X \times \mathcal{P}(1 + A \times X)) + (X \times \mathcal{P}(1 + A \times X))^2 \rightarrow \mathcal{P}(1 + A \times \Sigma^* X)$$

- $\rho(*) = \{*\}$
- $\rho((a, t, T)) = \{(a, t)\}$
- $\rho((t, T), (u, U)) = \{(a, t') \mid \forall (a, t') \in T\} \cup \{(a, u') \mid \forall (a, u') \in U\} \cup$

$$\frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a$$

$$\frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a$$

## 1.2 Abstract GSOS

- any syntax functor  $\Sigma$  and behaviour functor  $H$
- rules  $\rightsquigarrow$  a **natural transformation**  $\rho_X : \Sigma(X \times HX) \rightarrow H\Sigma^* X$

*Example:* For the previous example without  $?$  :  $\Sigma X = 1 + A \times X + X^2$ ,

$$\rho : 1 + A \times (X \times \mathcal{P}(1 + A \times X)) + (X \times \mathcal{P}(1 + A \times X))^2 \rightarrow \mathcal{P}(1 + A \times \Sigma^* X)$$

- $\rho(*) = \{*\}$
- $\rho((a, t, T)) = \{(a, t)\}$
- $\rho((t, T), (u, U)) = \{(a, t') \mid \forall (a, t') \in T\} \cup \{(a, u') \mid \forall (a, u') \in U\} \cup \{* \mid * \in T \wedge * \in U\}$

$$\frac{t \xrightarrow{a} t'}{t + u \xrightarrow{a} t'} \forall a$$

$$\frac{u \xrightarrow{a} u'}{t + u \xrightarrow{a} u'} \forall a$$

$$\frac{t \downarrow \quad u \downarrow}{t + u \downarrow}$$

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow$$

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$ , defined by coinduction


$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \qquad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) =$




## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t = \text{set of words of } A^* \text{ that can be produced by } t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \qquad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,




## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t = \text{set of words of } A^* \text{ that can be produced by } t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) =$






## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t = \text{set of words of } A^* \text{ that can be produced by } t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,




## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t = \text{set of words of } A^* \text{ that can be produced by } t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,  $\text{tr } (a + b? c) =$




## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t = \text{set of words of } A^* \text{ that can be produced by } t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,  $\text{tr } (a + b?c) = \{a\}$



## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,  $\text{tr } (a + b? c) = \{a\}$

- recall  $HX = \mathcal{P}(1 + A \times X)$

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,  $\text{tr } (a + b? c) = \{a\}$

- recall  $HX = \mathcal{P}(1 + A \times X) = TBX$ 
  - $T = \mathcal{P}$  **effectful** behaviour
  - $B = 1 + A \times X$  **pure** behaviour

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t = \text{set of words of } A^* \text{ that can be produced by } t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,  $\text{tr } (a + b? c) = \{a\}$

- recall  $HX = \mathcal{P}(1 + A \times X) = TBX$ 
  - ▶  $T = \mathcal{P}$  **effectful** behaviour  $\leadsto$  powerset : non-determinism
  - ▶  $B = 1 + A \times X$  **pure** behaviour  $\leadsto$  words :  $A^*$  (initial  $B$ -algebra)

## 1.3 Trace & Kleisli categories

- **trace** of a term  $t$ :  $\text{tr } t =$  set of words of  $A^*$  that can be produced by  $t$ , defined by coinduction

$$\varepsilon \in \text{tr } t \Leftrightarrow t \downarrow \quad a.w \in \text{tr } t \Leftrightarrow t \xrightarrow{a} u \wedge w \in \text{tr } u$$

*Example:*  $\text{tr } a(b + c) = \{ab, ac\}$ ,  $\text{tr } (ab + ac) = \{ab, ac\}$ ,  $\text{tr } (a + b? c) = \{a\}$

- recall  $HX = \mathcal{P}(1 + A \times X) = TBX$ 
  - $T = \mathcal{P}$  **effectful** behaviour  $\leadsto$  powerset : non-determinism
  - $B = 1 + A \times X$  **pure** behaviour  $\leadsto$  words :  $A^*$  (initial  $B$ -algebra)
- $\text{tr } t \in \mathcal{P}(A^*)$

## 1.3 Trace & Kleisli categories

Trace **abstractly**



## 1.3 Trace & Kleisli categories

Trace **abstractly**

- in the **Kleisli category** of  $T$

## 1.3 Trace & Kleisli categories

Trace **abstractly**

- in the **Kleisli category** of  $T$

$$A \in \mathbf{Kl}(T) \Leftrightarrow A \in \mathbb{C}$$

$$A \multimap B \in \mathbf{Kl}(T) \Leftrightarrow A \rightarrow TB \in \mathbb{C}$$

## 1.3 Trace & Kleisli categories

Trace **abstractly**

- in the **Kleisli category** of  $T$

$$A \in \mathbf{Kl}(T) \Leftrightarrow A \in \mathbb{C}$$

$$A \multimap B \in \mathbf{Kl}(T) \Leftrightarrow A \rightarrow TB \in \mathbb{C}$$

- $A^*$  is the final  $B$ -coalgebra in  $\mathbf{Kl}(T)$

## 1.3 Trace & Kleisli categories

Trace **abstractly**

- in the **Kleisli category** of  $T$

$$A \in \mathbf{Kl}(T) \Leftrightarrow A \in \mathbb{C}$$

$$A \multimap B \in \mathbf{Kl}(T) \Leftrightarrow A \rightarrow TB \in \mathbb{C}$$

- $A^*$  is the final  $B$ -coalgebra in  $\mathbf{Kl}(T)$

$$\zeta : A^* \multimap BA^* \text{ or } A^* \rightarrow TBA^*$$

$$\zeta(\varepsilon) = \{*\}, \quad \zeta(a.w) = \{(a, w)\}$$

## 1.3 Trace & Kleisli categories

Trace **abstractly**

- in the **Kleisli category** of  $T$

$$A \in \mathbf{Kl}(T) \Leftrightarrow A \in \mathbb{C}$$

$$A \multimap B \in \mathbf{Kl}(T) \Leftrightarrow A \rightarrow TB \in \mathbb{C}$$

- $A^*$  is the final  $B$ -coalgebra in  $\mathbf{Kl}(T)$

$$\zeta : A^* \multimap BA^* \text{ or } A^* \rightarrow TBA^*$$

$$\zeta(\varepsilon) = \{*\}, \quad \zeta(a.w) = \{(a, w)\}$$

- for any  $k : X \multimap BX$ ,

$$\begin{array}{ccc} X & \xrightarrow{k} & BX \\ \text{tr}_k \downarrow & & \downarrow B\text{tr}_k \\ A^* & \xrightarrow{\zeta} & BA^* \end{array}$$

## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times HX) \rightarrow H\Sigma^* X$$

## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times TBX) \rightarrow TB\Sigma^*X$$

## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times TBX) \rightarrow TB\Sigma^*X$$

- **Trace**-GSOS rule

$$\rho : \Sigma(X \times BX) \rightarrow TB\Sigma^*X$$



## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times TBX) \rightarrow TB\Sigma^* X$$

- **Trace**-GSOS rule

$$\rho : \Sigma(X \times BX) \rightarrow TB\Sigma^* X$$

$\leadsto$  only pure observations

## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times TBX) \rightarrow TB\Sigma^*X$$

- **Trace**-GSOS rule

$$\rho : \Sigma(X \times BX) \dashrightarrow B\Sigma^*X$$

$\leadsto$  only pure observations

## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times TBX) \rightarrow TB\Sigma^*X$$

- **Trace**-GSOS rule

$$\rho : \Sigma(X \times BX) \multimap B\Sigma^*X$$

$\leadsto$  only pure observations

- Rules observe each variable **once and only once**

## 1.4 Trace-GSOS

- GSOS rule

$$\rho : \Sigma(X \times TBX) \rightarrow TB\Sigma^*X$$

- Trace**-GSOS rule

$$\rho : \Sigma(X \times BX) \rightarrow B\Sigma^*X$$

$\leadsto$  only pure observations

- Rules observe each variable **once and only once**

*Example:*

$$\frac{t \xrightarrow{\tau} t' \quad t \xrightarrow{\tau} t''}{?t \xrightarrow{\tau} t' + t''}$$



$$\frac{}{a.t \xrightarrow{a} t} \forall a$$



$$\frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b$$



$$\frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a$$



## 1.5 Trace equivalence & congruence

- **trace equivalence:**


## 1.5 Trace equivalence & congruence

- **trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

## 1.5 Trace equivalence & congruence

- trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$


*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .



## 1.5 Trace equivalence & congruence

- trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .




$$a(b + c) \equiv_{\text{tr}} ab + ac$$



## 1.5 Trace equivalence & congruence

- trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .



$a(b + c) \equiv_{\text{tr}} ab + ac$  but not bisimilar  $\leadsto \equiv_{\text{tr}}$  coarsest

## 1.5 Trace equivalence & congruence

- **trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .  $a(b + c) \equiv_{\text{tr}} ab + ac$  but not bisimilar  $\sim_{\text{tr}} \equiv$  coarsest

- **congruence:**

## 1.5 Trace equivalence & congruence

- **trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .  $a(b + c) \equiv_{\text{tr}} ab + ac$  but not bisimilar  $\sim_{\text{tr}} \equiv$  coarsest

- **congruence:**  $\forall \sigma, (\forall i, t_i \equiv u_i) \Rightarrow \sigma(t_1 \dots t_n) \equiv \sigma(u_1 \dots u_n)$

## 1.5 Trace equivalence & congruence

- **trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .  $a(b + c) \equiv_{\text{tr}} ab + ac$  but not bisimilar  $\sim_{\text{tr}} \equiv$  coarsest


- **congruence:**  $\forall \sigma, (\forall i, t_i \equiv u_i) \Rightarrow \sigma(t_1 \dots t_n) \equiv \sigma(u_1 \dots u_n)$
- prove  $\text{tr}(\sigma(t_1 \dots t_n)) = \llbracket \sigma \rrbracket(\text{tr } t_1 \dots \text{tr } t_n)$

## 1.5 Trace equivalence & congruence

- trace equivalence:**  $t \equiv_{\text{tr}} u \Leftrightarrow \text{tr } t = \text{tr } u$

*Example:* Recall  $\text{tr } a(b + c) = \{ab, ac\} = \text{tr } (ab + ac)$ .  $a(b + c) \equiv_{\text{tr}} ab + ac$  but not bisimilar  $\sim_{\text{tr}} \equiv$  coarsest

- congruence:**  $\forall \sigma, (\forall i, t_i \equiv u_i) \Rightarrow \sigma(t_1 \dots t_n) \equiv \sigma(u_1 \dots u_n)$
- prove  $\text{tr}(\sigma(t_1 \dots t_n)) = \llbracket \sigma \rrbracket(\text{tr } t_1 \dots \text{tr } t_n)$



## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y)$

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$



## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )
- **affine monad**:  $TX \times TY \xrightarrow{\text{dst}} T(X \times Y) \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} TX \times TY = \text{id}$  or  $\eta_1 : 1 \xrightarrow{\simeq} T1$

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )
- **affine monad**:  $TX \times TY \xrightarrow{\text{dst}} T(X \times Y) \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} TX \times TY = \text{id}$  or  $\eta_1 : 1 \xrightarrow{\simeq} T1$
- **affine part**: greatest affine submonad

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )
- **affine monad**:  $TX \times TY \xrightarrow{\text{dst}} T(X \times Y) \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} TX \times TY = \text{id}$  or  $\eta_1 : 1 \xrightarrow{\simeq} T1$
- **affine part**: greatest affine submonad

*Example:*

- Powerset  $\mathcal{P} \rightsquigarrow \mathcal{P}_{\text{ne}}$

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )
- **affine monad**:  $TX \times TY \xrightarrow{\text{dst}} T(X \times Y) \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} TX \times TY = \text{id}$  or  $\eta_1 : 1 \xrightarrow{\simeq} T1$
- **affine part**: greatest affine submonad

*Example:*

- Powerset  $\mathcal{P} \rightsquigarrow \mathcal{P}_{\text{ne}}$
- (Sub)distribution  $\mathcal{S} \rightsquigarrow \mathcal{D}$

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )
- **affine monad**:  $TX \times TY \xrightarrow{\text{dst}} T(X \times Y) \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} TX \times TY = \text{id}$  or  $\eta_1 : 1 \xrightarrow{\simeq} T1$
- **affine part**: greatest affine submonad

*Example:*

- Powerset  $\mathcal{P} \rightsquigarrow \mathcal{P}_{\text{ne}}$
- (Sub)distribution  $\mathcal{S} \rightsquigarrow \mathcal{D}$  with  $\mathcal{D}X = \left\{ \sum_{i \in I} p_i x_i \mid \sum p_i = 1, x_i \in X, I \text{ finite} \right\}$   
and  $\mathcal{S}X = \left\{ \sum_{i \in I} p_i x_i \mid \sum p_i \leq 1, x_i \in X, I \text{ finite} \right\}$

## 1.6 Strong and affine monads

- **strong monad**:  $\text{st}_{X,Y} : X \times TY \rightarrow T(X \times Y) \rightsquigarrow \text{st}' : TX \times Y \rightarrow T(X \times Y)$
- double strength  $\text{dst} : TX \times TY \xrightarrow{\text{st}} T(TX \times Y) \xrightarrow{T\text{st}'} T^2(X \times Y) \xrightarrow{\mu} T(X \times Y)$  (and  $\text{dst}'$ )
- **affine monad**:  $TX \times TY \xrightarrow{\text{dst}} T(X \times Y) \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} TX \times TY = \text{id}$  or  $\eta_1 : 1 \xrightarrow{\simeq} T1$
- **affine part**: greatest affine submonad

*Example:*

- Powerset  $\mathcal{P} \rightsquigarrow \mathcal{P}_{\text{ne}}$
- (Sub)distribution  $\mathcal{S} \rightsquigarrow \mathcal{D}$  with  $\mathcal{D}X = \left\{ \sum_{i \in I} p_i x_i \mid \sum p_i = 1, x_i \in X, I \text{ finite} \right\}$   
and  $\mathcal{S}X = \left\{ \sum_{i \in I} p_i x_i \mid \sum p_i \leq 1, x_i \in X, I \text{ finite} \right\}$
- Maybe  $-+1 \rightsquigarrow \text{Id}$

## 2. Result

---



## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,  $B$  a *behaviour* endofunctor that extends to  $\mathbf{Kl}(T)$ ,

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,  $B$  a *behaviour* endofunctor that extends to  $\mathbf{Kl}(T)$ ,  $\Sigma$  a *syntax* endofunctor that extends to  $\mathbf{Kl}(T)$  with all free objects  $(\Sigma^* X)$ ,

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,  $B$  a *behaviour* endofunctor that extends to  $\text{Kl}(T)$ ,  $\Sigma$  a *syntax* endofunctor that extends to  $\text{Kl}(T)$  with all free objects  $(\Sigma^* X)$ , let  $\zeta : Z \rightarrowtail BZ$  be the final  $\overline{B}$ -coalgebra (with  $\exists z, \zeta = \eta \circ z$ ) and

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,  $B$  a *behaviour* endofunctor that extends to  $\mathbf{Kl}(T)$ ,  $\Sigma$  a *syntax* endofunctor that extends to  $\mathbf{Kl}(T)$  with all free objects  $(\Sigma^* X)$ , let  $\zeta : Z \rightarrowtail BZ$  be the final  $\overline{B}$ -coalgebra (with  $\exists z, \zeta = \eta \circ z$ ) and let  $\rho : \Sigma(X \times BX) \rightarrow TB\Sigma^* X$  be a natural transformation *representing Trace-GSOS rules*

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,  $B$  a *behaviour* endofunctor that extends to  $\mathbf{Kl}(T)$ ,  $\Sigma$  a *syntax* endofunctor that extends to  $\mathbf{Kl}(T)$  with all free objects  $(\Sigma^* X)$ , let  $\zeta : Z \rightarrowtail BZ$  be the final  $\overline{B}$ -coalgebra (with  $\exists z, \zeta = \eta \circ z$ ) and let  $\rho : \Sigma(X \times BX) \rightarrow TB\Sigma^* X$  be a natural transformation representing *Trace-GSOS rules* such that  $\rho$  is **smooth** and is a map of distributive laws,

## 2.1 The theorem

**Theorem 2.1.1:** Let  $\mathbb{C}$  be a cartesian category,  $T$  be a strong **affine** *effectful* monad,  $B$  a *behaviour* endofunctor that extends to  $\mathbf{Kl}(T)$ ,  $\Sigma$  a *syntax* endofunctor that extends to  $\mathbf{Kl}(T)$  with all free objects  $(\Sigma^* X)$ , let  $\zeta : Z \rightarrowtail BZ$  be the final  $\overline{B}$ -coalgebra (with  $\exists z, \zeta = \eta \circ z$ ) and let  $\rho : \Sigma(X \times BX) \rightarrow TB\Sigma^* X$  be a natural transformation *representing Trace-GSOS rules* such that  $\rho$  is **smooth** and is a map of distributive laws, then trace equivalence is a congruence.



## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{[-]} & Z \end{array}$$

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace
- $\Sigma^* X \rightarrow B\Sigma^* X$  (with  $\rho^*$ )

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace
- $\Sigma^* X \rightarrow B\Sigma^* X$  (with  $\rho^*$ ) and  $Z \rightarrow BZ$

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace
- $\Sigma^* X \rightarrow B\Sigma^* X$  (with  $\rho^*$ ) and  $Z \rightarrow BZ$
- show  $\overline{B}$ -coalgebra morphisms

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace
- $\Sigma^* X \rightarrow B\Sigma^* X$  (with  $\rho^*$ ) and  $Z \rightarrow BZ$
- show  $\overline{B}$ -coalgebra morphisms
- $\text{tr} \circ i \quad \checkmark$

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace
- $\Sigma^* X \rightarrow \bullet B\Sigma^* X$  (with  $\rho^*$ ) and  $Z \rightarrow \bullet BZ$
- show  $\overline{B}$ -coalgebra morphisms
- $\text{tr} \circ i \quad \checkmark$
- $\llbracket - \rrbracket \circ \Sigma^* \text{tr}$  more complicated : naturality + smoothness + map of distributive law of  $\rho^*$

## 2.2 Sketch of the proof

- Recall

$$\begin{array}{ccc} \Sigma^* X & \xrightarrow{i} & X \\ \Sigma^* \text{tr}_k \downarrow & & \downarrow \text{tr}_k \\ \Sigma^* Z & \xrightarrow{\llbracket - \rrbracket} & Z \end{array}$$

- define  $\llbracket - \rrbracket$  : semantics of  $Z$  + induction + trace
- $\Sigma^* X \rightarrow \bullet B\Sigma^* X$  (with  $\rho^*$ ) and  $Z \rightarrow \bullet BZ$
- show  $\overline{B}$ -coalgebra morphisms
- $\text{tr} \circ i \quad \checkmark$
- $\llbracket - \rrbracket \circ \Sigma^* \text{tr}$  more complicated : naturality + smoothness + map of distributive law of  $\rho^*$

*Remark:* need dst



## 2.3 Focus on hypothesis : Smoothness

*Example:*  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0$ ,  $a.$ ,  $+$  and

## 2.3 Focus on hypothesis : Smoothness


*Example:*  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0$ ,  $a.$ ,  $+$  and

$$\frac{t \xrightarrow{a} t'}{\quad} \forall a \qquad \frac{t \xrightarrow{a} t'}{\quad} \forall a$$
$$!t \xrightarrow{a} ?t' \qquad ?t \xrightarrow{a} t + t'$$

## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0$ ,  $a.$ ,  $+$  and


$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{a} t'}{?t \xrightarrow{a} t + t'} \forall a$$



## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0$ ,  $a.$ ,  $+$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{a} t'}{?t \xrightarrow{a} t + t'} \forall a$$




$$\text{tr } !a(b + c) = \{ab, ac, abb, \underline{abc}, \underline{acb}, acc\}$$


## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for 0,  $a.$ ,  $+$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{a} t'}{?t \xrightarrow{a} t + t'} \forall a$$




$$\text{tr } !a(b + c) = \{ab, ac, abb, \underline{abc}, \underline{acb}, acc\}$$



## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for 0,  $a.$ ,  $+$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{a} t'}{?t \xrightarrow{a} t + t'} \forall a$$



$$\text{tr } !a(b+c) = \{ab, ac, abb, \underline{abc}, \underline{acb}, acc\} \neq \{ab, ac, abb, acc\} = \text{tr } !(ab+ac)$$

## 2.3 Focus on hypothesis : Smoothness


*Example:*  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0, a., +$  and

$$\frac{t \xrightarrow{a} t'}{\quad} \forall a \qquad \frac{t \xrightarrow{\tau} t'}{\quad}$$
$$!t \xrightarrow{a} ?t' \qquad ?t \xrightarrow{\tau} t$$

## 2.3 Focus on hypothesis : Smoothness

*Example:*  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0$ ,  $a.$ ,  $+$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t}$$






## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0, a., +$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t}$$




$$\text{tr } !a(b + \tau) = \{\underline{a\tau b}, a\tau\tau\}$$

## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0$ ,  $a$ .,  $+$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t}$$




$$\text{tr } !a(b + \tau) = \{\underline{a\tau b}, a\tau\tau\}$$

## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0, a., +$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t}$$




$$\text{tr } !a(b + \tau) = \{\underline{a\tau b}, a\tau\tau\} \neq \{a\tau\tau\} = \text{tr } !(ab + a\tau)$$

## 2.3 Focus on hypothesis : Smoothness

Example:  $t ::= 0 \mid a.t \mid t + t \mid ?t \mid !t$  with the previous rules for  $0, a., +$  and

$$\frac{t \xrightarrow{a} t'}{!t \xrightarrow{a} ?t'} \forall a \qquad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t}$$



$$\text{tr } !a(b + \tau) = \{\underline{a\tau b}, a\tau\tau\} \neq \{a\tau\tau\} = \text{tr } !(ab + a\tau)$$

→ observations that are “not used” 😞

## 2.3 Focus on hypothesis : Smoothness

- **smoothness**

## 2.3 Focus on hypothesis : Smoothness

- **smoothness**
  - **linear**: if  $x_i \rightarrow x_{i'}$  then not  $x_i$  and  $x_{i'}$  in the target

## 2.3 Focus on hypothesis : Smoothness

- **smoothness**
  - ▶ **linear**: if  $x_i \rightarrow x_{i'}$  then not  $x_i$  and  $x_{i'}$  in the target
  - ▶ if  $x_i$  in the target, the observation on  $x_i$  is **irrelevant** ie. any other observation could have been done (the same rule for each other possible observation exists)

## 2.3 Focus on hypothesis : Smoothness

- **smoothness**
  - **linear**: if  $x_i \rightarrow x_{i'}$ , then not  $x_i$  and  $x_{i'}$  in the target
  - if  $x_i$  in the target, the observation on  $x_i$  is **irrelevant** ie. any other observation could have been done (the same rule for each other possible observation exists)
- **abstract smoothness**

$$\begin{array}{ccccc}
 & & \text{mix} & & \\
 & \swarrow & & \searrow & \\
 \Sigma T(X \times BX) & \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} & \Sigma(TX \times TBX) & \xrightarrow{\text{dst}} & \Sigma T(X \times BX) \\
 \lambda \downarrow & & & & \lambda \downarrow \\
 T\Sigma(X \times BX) & & & & T\Sigma(X \times BX) \\
 T\rho \downarrow & & & & T\rho \downarrow \\
 T^2 B\Sigma^* X & \xrightarrow{\mu} & TB\Sigma^* X & \xleftarrow{\mu} & T^2 B\Sigma^* X
 \end{array}$$



## 2.3 Focus on hypothesis : Smoothness

- **smoothness**

- ▶ **linear**: if  $x_i \rightarrow x_{i'}$ , then not  $x_i$  and  $x_{i'}$  in the target
- ▶ if  $x_i$  in the target, the observation on  $x_i$  is **irrelevant** ie. any other observation could have been done (the same rule for each other possible observation exists)

- **abstract smoothness**

$$\begin{array}{ccc}
 & \text{mix} & \\
 & \text{-----} & \\
 \Sigma T(X \times BX) & \xrightarrow{\langle T\pi_1, T\pi_2 \rangle} & \Sigma(TX \times TBX) \xrightarrow{\text{dst}} \Sigma T(X \times BX) \\
 \lambda \downarrow & & \lambda \downarrow \\
 T\Sigma(X \times BX) & & T\Sigma(X \times BX) \\
 T\rho \downarrow & & T\rho \downarrow \\
 T^2 B\Sigma^* X & \xrightarrow{\mu} & TB\Sigma^* X \xleftarrow{\mu} T^2 B\Sigma^* X
 \end{array}$$

$$\begin{aligned}
 \Phi(\rho)(\sigma)(\text{mix } X_1 \dots \text{mix } X_n) = \\
 \Phi(\rho)(\sigma)(X_1 \dots X_n) \\
 \text{where } X_i \subset X \times BX
 \end{aligned}$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

*Example:*  $t ::= 0 \mid a.t \mid ? t$  with the following smooth rules

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

*Example:*  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

*Example:*  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

$$\text{let } X_1 = \left\{ t \xrightarrow{\tau} t', u \xrightarrow{a} u' \right\}$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

*Example:*  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

$$\text{let } X_1 = \left\{ t \xrightarrow{\tau} t', u \xrightarrow{a} u' \right\} \text{ then mix } X_1 = \left\{ t \xrightarrow{\tau} t', t \xrightarrow{a} u', u \xrightarrow{\tau} t', u \xrightarrow{a} u' \right\}$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

Example:  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

let  $X_1 = \{t \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$  then  $\text{mix } X_1 = \{t \xrightarrow{\tau} t', t \xrightarrow{a} u', u \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$

$$\frac{t \xrightarrow{\tau} t' \in X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} ? \notin X_1}{?u \nrightarrow}$$

$$\frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \nrightarrow}{b.?u \nrightarrow}$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

Example:  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

let  $X_1 = \left\{ t \xrightarrow{\tau} t', u \xrightarrow{a} u' \right\}$  then  $\text{mix } X_1 = \left\{ t \xrightarrow{\tau} t', t \xrightarrow{a} u', u \xrightarrow{\tau} t', u \xrightarrow{a} u' \right\}$

$$\frac{t \xrightarrow{\tau} t' \in X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} ? \notin X_1}{?u \nrightarrow}$$

$$\frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \nrightarrow}{b.?u \nrightarrow}$$

$$\Phi(\rho^*)(b.?x_1)(X_1) = \left\{ \xrightarrow{b} ?t \right\}$$



## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

Example:  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

let  $X_1 = \{t \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$  then  $\text{mix } X_1 = \{t \xrightarrow{\tau} t', t \xrightarrow{a} u', u \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$

$$\frac{t \xrightarrow{\tau} t' \in X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} ? \notin X_1}{?u \nrightarrow} \quad \frac{t \xrightarrow{\tau} t' \in \text{mix } X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} t' \in \text{mix } X_1}{?u \xrightarrow{\tau} t'}$$

$$\frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \nrightarrow}{b.?u \nrightarrow} \quad \frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \xrightarrow{\tau} t'}{b.?u \xrightarrow{b} ?u}$$

$$\Phi(\rho^*)(b.?x_1)(X_1) = \left\{ \xrightarrow{b} ?t \right\}$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

Example:  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

let  $X_1 = \{t \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$  then  $\text{mix } X_1 = \{t \xrightarrow{\tau} t', t \xrightarrow{a} u', u \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$

$$\frac{t \xrightarrow{\tau} t' \in X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} ? \notin X_1}{?u \nrightarrow} \quad \frac{t \xrightarrow{\tau} t' \in \text{mix } X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} t' \in \text{mix } X_1}{?u \xrightarrow{\tau} t'}$$

$$\frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \nrightarrow}{b.?u \nrightarrow} \quad \frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \xrightarrow{\tau} t'}{b.?u \xrightarrow{b} ?u}$$

$$\Phi(\rho^*)(b.?x_1)(X_1) = \left\{ \xrightarrow{b} ?t \right\} \neq \left\{ \xrightarrow{b} ?t, \xrightarrow{b} ?u \right\} = \Phi(\rho^*)(b.?x_1)(\text{mix } X_1)$$

## 2.3 Focus on hypothesis : Smoothness

- need smoothness **for**  $\rho^*$  (terms with more than one layer)

Example:  $t ::= 0 \mid a.t \mid ?t$  with the following smooth rules

$$\frac{}{0 \downarrow} \quad \frac{t \xrightarrow{b} t'}{a.t \xrightarrow{a} t} \forall a, b \quad \frac{t \downarrow}{a.t \xrightarrow{a} t} \forall a \quad \frac{t \xrightarrow{\tau} t'}{?t \xrightarrow{\tau} t'}$$

let  $X_1 = \{t \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$  then  $\text{mix } X_1 = \{t \xrightarrow{\tau} t', t \xrightarrow{a} u', u \xrightarrow{\tau} t', u \xrightarrow{a} u'\}$

$$\frac{t \xrightarrow{\tau} t' \in X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} ? \notin X_1}{?u \nrightarrow} \quad \frac{t \xrightarrow{\tau} t' \in \text{mix } X_1}{?t \xrightarrow{\tau} t'} \quad \frac{u \xrightarrow{\tau} t' \in \text{mix } X_1}{?u \xrightarrow{\tau} t'}$$

$$\frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \nrightarrow}{b.?u \nrightarrow} \quad \frac{?t \xrightarrow{\tau} t'}{b.?t \xrightarrow{b} ?t} \quad \frac{?u \xrightarrow{\tau} t'}{b.?u \xrightarrow{b} ?u}$$

$\Phi(\rho^*)(b.?x_1)(X_1) = \left\{ \xrightarrow{b} ?t \right\} \neq \left\{ \xrightarrow{b} ?t, \xrightarrow{b} ?u \right\} = \Phi(\rho^*)(b.?x_1)(\text{mix } X_1) \rightarrow$  the

**stuck** computation is messing with smoothness 😞

## 2.4 Focus on hypothesis : Affineness

**Theorem 2.4.1:** If  $T$  is an **affine** monad then the smoothness of  $\rho$  entails the smoothness of  $\rho^*$ .

## 2.4 Focus on hypothesis : Affineness

**Theorem 2.4.1:** If  $T$  is an **affine** monad then the smoothness of  $\rho$  entails the smoothness of  $\rho^*$ .

- affine part of  $\mathcal{P}$  is  $\mathcal{P}_{\text{ne}} \longrightarrow$  no stuckness !

## 2.4 Focus on hypothesis : Affineness

**Theorem 2.4.1:** If  $T$  is an **affine** monad then the smoothness of  $\rho$  entails the smoothness of  $\rho^*$ .

- affine part of  $\mathcal{P}$  is  $\mathcal{P}_{\text{ne}} \longrightarrow$  no stuckness !
- at the level of rules: give a semantics to **every situation**, nothing unspecified

## 2.4 Focus on hypothesis : Affineness

**Theorem 2.4.1:** If  $T$  is an **affine** monad then the smoothness of  $\rho$  entails the smoothness of  $\rho^*$ .

- affine part of  $\mathcal{P}$  is  $\mathcal{P}_{\text{ne}} \longrightarrow$  no stuckness !
- at the level of rules: give a semantics to **every situation**, nothing unspecified

*Example:*

$$\frac{t \xrightarrow{\tau} t'}{? t \xrightarrow{\tau} t'}$$

## 2.4 Focus on hypothesis : Affineness

**Theorem 2.4.1:** If  $T$  is an **affine** monad then the smoothness of  $\rho$  entails the smoothness of  $\rho^*$ .

- affine part of  $\mathcal{P}$  is  $\mathcal{P}_{\text{ne}} \longrightarrow$  no stuckness !
- at the level of rules: give a semantics to **every situation**, nothing unspecified

*Example:*

$$\frac{t \xrightarrow{\tau} t'}{? t \xrightarrow{\tau} t'} \quad + \quad \frac{t \xrightarrow{a} t'}{? t \dots} \forall a \neq \tau \quad \frac{t \downarrow}{? t \dots}$$

need to have some semantics



## 2.4 Focus on hypothesis : Affineness

**Theorem 2.4.1:** If  $T$  is an **affine** monad then the smoothness of  $\rho$  entails the smoothness of  $\rho^*$ .

- affine part of  $\mathcal{P}$  is  $\mathcal{P}_{\text{ne}} \longrightarrow$  no stuckness !
- at the level of rules: give a semantics to **every situation**, nothing unspecified

*Example:*

$$\frac{t \xrightarrow{\tau} t'}{? t \xrightarrow{\tau} t'} \quad + \quad \frac{t \xrightarrow{a} t'}{? t \downarrow} \quad \forall a \neq \tau \quad \frac{t \downarrow}{? t \downarrow}$$

need to have some semantics eg. termination  $\downarrow$

## 2.5 And for non affine monads ?

- still under investigation 🚧

## 2.5 And for non affine monads ?

- still under investigation 🚧
- idea 1: add an **extra sink state**  $\perp$  for stuck computations

## 2.5 And for non affine monads ?

- still under investigation 🚧
- idea 1: add an **extra sink state**  $\perp$  for stuck computations
- idea 2: map stuckness to **explicit termination** (cf. previous example) 🚨 change of semantics

## 2.5 And for non affine monads ?

- still under investigation 🚧
  - idea 1: add an **extra sink state**  $\perp$  for stuck computations
  - idea 2: map stuckness to **explicit termination** (cf. previous example) 🚨 change of semantics
- Can we get back information on the original system ?

### 3. Conclusion

---

### 3. Conclusion

- For an affine monadic effect, under reasonable assumptions, trace equivalence is a congruence 🎉 !

### 3. Conclusion

- For an affine monadic effect, under reasonable assumptions, trace equivalence is a congruence 🎉 !
- Can we do better ?



### 3. Conclusion

- For an affine monadic effect, under reasonable assumptions, trace equivalence is a congruence 🎉 !
- Can we do better ? Can we find a good reduction to the affine case for non affine monads ?

### 3. Conclusion

- For an affine monadic effect, under reasonable assumptions, trace equivalence is a congruence 🎉 !
- Can we do better ? Can we find a good reduction to the affine case for non affine monads ?
- Thank you all for welcoming me in the chair ❤️

*~ The End ~*

Powered by Typst