



NORMALITY TEST

HYPOTHESIS TESTING

prepared by:

Gyro A. Madrona
Electronics Engineer

TOPIC OUTLINE

Shapiro-Wilk

Anderson-Darling



SHAPIRO-WILK



SHAPIRO-WILK

The Shapiro-Wilk normality test calculates a statistic based on the correlation between the data and the corresponding normal distribution; effective for sample size ($n \leq 50$).

Shapiro Function

`shapiro()` is a statistical tool used to perform the Shapiro-Wilk test for normality.

Syntax

```
w_stat, p_value = stats.shapiro(data)
```

Null Hypothesis

H_0 : Normal data

Alternative Hypothesis

H_a : Non-normal data ($p\text{-value} \leq \alpha$)



EXERCISE

Perform Shapiro-Wilk normality test for the given dataset.

dataset

“defects-data-30-samples.csv”

Solution

```
w_stat, p_value =  
stats.shapiro(df['Defects'])
```

Let $\alpha = 0.05$

Null Hypothesis

H_o : Normal data

Alternative Hypothesis

H_a : Non-normal data (p-value ≤ 0.05)



ANDERSON-DARLING



ANDERSON-DARLING

The Anderson-Darling normality test provides a more sensitive test by giving weight to the tails of the distribution; effective for larger sample size ($n > 50$).

Anderson Function

`anderson ()` is a statistical tool used to perform the Anderson-Darling test for normality.

Syntax

```
a2_stat, critical_values, alpha =  
stats.anderson(data)
```

Null Hypothesis

$H_o: A^2 \leq \text{critical value}$ (Normal data)

Alternative Hypothesis

$H_a: A^2 > \text{critical value}$ (Non-normal data)



EXERCISE

Perform Anderson-Darling normality test for the given dataset.

dataset

“defects-dataset.csv”

Solution

```
a2_stat, critical_values, alpha =  
stats.anderson(df['Defects'])
```

Let $\alpha = 0.05$

Null Hypothesis

$H_0: A^2 \leq \text{critical value}$ (Normal data)

Alternative Hypothesis

$H_a: A^2 > \text{critical value}$ (Non-normal data)



LABORATORY

