



HYPOTHESIS TESTING

INFERENTIAL STATISTICS

prepared by:

Gyro A. Madrona
Electronics Engineer

TOPIC OUTLINE

Hypothesis Test

Rejection Region

Critical Value and Z-score

p-Value



HYPOTHESIS TEST



HYPOTHESIS

A hypothesis is an initial assumption formed before collecting data, and it serves as a statement about a population parameter rather than about the sample data.



HYPOTHESIS TEST

A hypothesis test is simply comparing reality to an assumption and asking, “Did things change?”

Null Hypothesis (H_o)

Represents no change, no effect, or the status quo.

Alternative Hypothesis (H_a)

Represents the possibility that things did change or that there is a significant difference.



IS YOUR DATA GUILTY?

Hypothesis testing is like a legal system where the defendant is assumed innocent until proven guilty.

H_0 : Innocent

	The Truth	
	Innocent	Guilty
Accept	✓	✗
Reject	✗	✓




TYPES OF ERROR

1. Type I Error :

Rejecting the null hypothesis when it is actually true. The risk of making type I error is denoted by α (e.g., 0.05).

2. Type II Error :

Failing to reject the null hypothesis when it is actually false. The risk of making a type II error is denoted by β (e.g. 0.20)



		The Truth	
		Innocent	Guilty
H_0 : Innocent	Accept	✓	Type II Error (False negative)
	Reject	Type I Error (False positive)	✓

EXERCISE

A fitness tracker company claims their device measures heart rate with **95%** accuracy compared to medical-grade monitors. An independent lab wants to verify this claim.

Null Hypothesis

$$H_o: \mu_o = 95$$

The average accuracy is 95%.

Alternative Hypothesis

$$H_a: \mu_o \neq 95$$

The average accuracy differs from 95%.



EXERCISE

A manufacturer claims that their new energy-efficient LED bulbs have an average lifespan of **at least 25,000 hours**. A consumer group suspects that the actual lifespan is shorter and decides to test this claim.

Null Hypothesis

$$H_o: \mu_o = 25,000$$

The average lifespan of the LED is 25,000 hours.

Alternative Hypothesis

$$H_a: \mu_o < 25,000$$

The average lifespan of the LED is less than 25,000 hours.



EXERCISE

A study suggests that storing apples in a controlled atmosphere **extends** their shelf life beyond **30 days**. A food scientist wants to verify if this method truly increases shelf life compared to conventional storage.

Null Hypothesis

$$H_o: \mu_o = 30$$

Controlled-atmosphere storage shelf life is 30 days.

Alternative Hypothesis

$$H_a: \mu_o > 30$$

Controlled-atmosphere storage increases shelf life beyond 30 days.



REJECTION REGION



SIGNIFICANCE LEVEL

The significance level (α) determines the threshold for deciding whether to reject the null hypothesis (H_o).


Typical values for α :

0.01

0.05


0.1

Standard Normal Distribution



REJECTION REGION


Two-sided Test



$$H_o: \mu_1 = \mu_o$$

$$H_a: \mu_1 \neq \mu_o$$

One-sided Test




$$H_o: \mu_1 = \mu_o$$

$$H_a: \mu_1 < \mu_o$$



REJECTION REGION


Two-sided Test



$$H_o: \mu_1 = \mu_o$$

$$H_a: \mu_1 \neq \mu_o$$

One-sided Test



$$H_o: \mu_1 = \mu_o$$

$$H_a: \mu_1 > \mu_o$$



CRITICAL VALUE AND Z-SCORE



CRITICAL VALUE AND Z-SCORE

One-sided Test

lowercase z


z refers to the critical value obtained from the standard normal distribution table (z-table).

uppercase Z

Z is a standardized variable associated with the test called the Z-score.


Formula:

$$Z = \frac{\bar{x} - \mu_o}{\sigma/\sqrt{n}}$$



EXERCISE

Two-sided Test



$$\alpha = 0.05$$

$$z_{0.025} = 1.96$$

Null Hypothesis

$$H_o: \mu_o = 95$$

The average accuracy is 95%.

Alternative Hypothesis


$$H_a: \mu_o \neq 95$$

The average accuracy differs from 95%.



EXERCISE

One-sided Test



$$\alpha = 0.05$$

$$z_{0.05} = 1.645$$

Null Hypothesis

$$H_o: \mu_o = 25,000$$

The average lifespan of the LED is 25,000 hours.

Alternative Hypothesis


$$H_a: \mu_o < 25,000$$

The average lifespan of the LED is less than 25,000 hours.



EXERCISE

One-sided Test



$$\alpha = 0.05$$

$$z_{0.05} = 1.645$$

Null Hypothesis

$$H_o: \mu_o = 30$$

Controlled-atmosphere storage shelf life is 30 days.

Alternative Hypothesis

$$H_a: \mu_o > 30$$

Controlled-atmosphere storage increases shelf life beyond 30 days.



EXERCISE

A manufacturing process is claimed to have an average defect rate of **10.32** units, with a known standard deviation of **3.17** units. The Statistical Process Control (SPC) department suspects this claim may no longer be valid and collects a **random sample** of **30** production units to test whether the true average **defect rate differs** significantly from **10.32**.

Dataset

defects-data-30-samples.csv

Solution



P-VALUE



P-VALUE

The **p-value** (probability value) is the **smallest level of significance** at which we can still reject the null hypothesis, given the observed sample statistic.

One-sided Test

$$\mathbf{p\text{-}value = 1 - \text{value from the table}}$$


Two-sided Test

$$\mathbf{p\text{-}value = (1 - \text{value from the table}) \times 2}$$

Hypothesis Test

Reject H_o if **p-value $\leq \alpha$**

Fail to reject H_o if p-value $> \alpha$



P-VALUE

The p-value (probability value) is the smallest level of significance at which we can still reject the null hypothesis, given the observed sample statistic.

One-sided Test

$$\text{p-value} = 1 - \text{value from the table}$$

Two-sided Test

$$\text{p-value} = (1 - \text{value from the table}) \times 2$$

Cumulative Distribution Function (CDF)

`cdf()` returns the probability that a random variable **Z** (Z-score) from a standard normal distribution is less than or equal to a given critical value (**z**).

Syntax

One-sided Test

$$\text{p_value} = 1 - \text{stats.norm.cdf}(\text{Z_score})$$

Two-sided Test

$$\text{p_value} = 2 * (1 - \text{stats.norm.cdf}(\text{Z_score}))$$



EXERCISE

A manufacturing process is claimed to have an average defect rate of **10.32** units, with a known standard deviation of **3.17** units. The Statistical Process Control (SPC) department suspects this claim may no longer be valid and collects a **random sample** of **30** production units to test whether the true average **defect rate differs** significantly from **10.32**.

Dataset

defects-data-30-samples.csv

Solution



LABORATORY

