Universe creation by tunnelling from laboratory data: no manifold instanton, a unique canonical exponent, and a Smarr identity

Project Baby Universe — Genesis Branch

21 October 2026 (Stage One manuscript, draft); addenda 22–23 October 2026

Abstract

We revisit the thin-wall model of a false-vacuum bubble (de Sitter interior, Schwarzschild exterior) and the proposal of Farhi, Guth and Guven that such a bubble, too small to inflate classically, may tunnel into an inflating, causally disconnected daughter universe. (i) We prove that, for every transition starting from regular, time-symmetric, asymptotically flat ("laboratory") data, the Euclidean exterior interpolating region cannot be embedded in Euclidean Schwarzschild. The pseudomanifold encountered by Farhi–Guth–Guven is therefore forced. The obstruction is tied to a sign change of the exterior extrinsic curvature while the wall recedes from the Euclidean horizon; an interior sign change does not obstruct embedding. (ii) In the reduced canonical theory with the asymptotic clock, the branch of the Euclidean momentum is fixed by analyticity. The competing branch is a horizon monodromy, which in Euclidean language is the half-turn slicing jump. (iii) At $M = 0$ the exponent is $B = S_E[\text{instanton}] + \frac{1}{4G}(\sum A_{\max} - \sum A_{\min})$ in closed form; it equals the Coleman–De Luccia exponent of the false vacuum when the false vacuum is the larger cap. (iv) For $M > 0$ the exponent obeys a first law $d\tilde{B} = (\mathrm{Vol}_E^{\mathrm{in}}/8\pi G)\, d\Lambda_{\mathrm{in}} + A_E^W\, d\sigma - \Delta t_E^W\, dM$ and a Smarr identity $B = -\frac{\Lambda_{\mathrm{in}}}{8\pi G}\mathrm{Vol}_E^{\mathrm{in}} - \frac{\sigma}{2}A_E^W - \frac{M}{2}\Delta t_E^W + \frac{\mathcal{A}}{4G}$. This is a covariant expression in which the pseudomanifold enters only through one boundary term. Numerically, over laboratory transitions $B \geq 0.56\, B(M \to 0)$ for all parameters and $B \geq B_{\mathrm{CdL}}$ when $\chi \gtrsim 1.2\kappa$; for GUT-scale parameters and gram-scale bubbles $B \sim 10^6$. All statements are verified symbolically and numerically with an open-source code.

1 Introduction

A spherical region of false vacuum separated from true vacuum by a domain wall can have an inflating interior even though, seen from outside, it is a black hole [1]. Farhi and Guth showed that such a configuration cannot arise classically from regular initial data when the null energy condition holds [2]. Farhi, Guth and Guven (FGG) asked whether quantum tunnelling could do it [3]. Their canonical calculation was hampered by the choice of slicing, and their Euclidean interpolation was not a manifold but a "pseudomanifold" with a conjectured action. Fischler, Morgan and Polchinski found agreement within a Hamiltonian treatment [4]. Later Hamiltonian work extended these transitions to all signs of the cosmological constant [7, 8]. Holographic arguments suggest that smooth saddles do not produce inflating bubbles [9, 10].

This paper makes the FGG difficulties precise and fixes the branch of the exponent within the reduced canonical theory (see the Addendum: a residual ambiguity survives in the single-flip class). Section 2 fixes the model and conventions. Section 3 summarises the global structure. Section 4 proves the embedding obstruction. Section 5 fixes the branch of the canonical exponent. Sections 6–7 give the closed form at $M = 0$, the first law and the Smarr identity. Section 8 collects numerical results and Section 9 the limitations.




Figure 1: Inflating bounce branch ($\gamma = -0.5$, regime A): effective potential, interior de Sitter diagram, exterior Kruskal diagram, all drawn from integrated trajectories.

2 Model and conventions

Signature $(-,+,+,+)$, $c = 1$. Interior $f_{\rm in} = 1 - \chi^2 r^2$, exterior $f_{\rm out} = 1 - 2GM/r - \eta^2 r^2$ (here mostly $\eta = 0$), wall $S_{ab} = -\sigma h_{ab}$, $\kappa = 4\pi G\sigma$. With n pointing from interior to exterior, $K_{ab} = e^\mu_a e^\nu_b \nabla_\mu n_\nu$, $[X] = X_+ - X_-$, the Israel conditions $[K_{ab}] - h_{ab}[K] = -8\pi G S_{ab}$ reduce to

$$A_{\rm in} - A_{\rm out} = \kappa R, \qquad A \equiv n^r = \varepsilon\sqrt{f + \dot R^2}, \tag{1}$$

with $K_{\theta\theta} = RA$ and $K^\tau{}_\tau = (\ddot R + \frac{1}{2}f')/A$. Using the identity $A_{\rm in}^2 - A_{\rm out}^2 = f_{\rm in} - f_{\rm out}$,

$$A_{\rm in} = \frac{f_{\rm in} - f_{\rm out} + \kappa^2 R^2}{2\kappa R}, \quad A_{\rm out} = \frac{f_{\rm in} - f_{\rm out} - \kappa^2 R^2}{2\kappa R}, \quad \dot R^2 + V(R) = 0, \quad V = f_{\rm out} - A_{\rm out}^2. \tag{2}$$

Lemma 1. *For $\sigma > 0$, $(R, \varepsilon_{\rm in}, \varepsilon_{\rm out})$ solves the unsquared junction condition iff $\dot R^2 + V = 0$ and $\varepsilon = \mathrm{sgn} A(R)$ on each side. The only spurious solutions of the squared equation are those with $\sigma < 0$.*

With $R = (2GM/L^2)^{1/3} z$, $L^4 = (\chi^2 - \eta^2 + \kappa^2)^2 + 4\kappa^2\eta^2$: $V = 1 - g(z)/E$, $g = z^2 - 2\gamma/z + z^{-4}$, $\gamma = (\chi^2 - \eta^2 - \kappa^2)/L^2$, $E = 4\kappa^2 L^{-8/3}(2GM)^{-2/3}$, and for $\eta = 0$, $\varepsilon_{\rm out} = \mathrm{sgn}(1 - z^3)$, $\varepsilon_{\rm in} = \mathrm{sgn}(1 - \gamma z^3)$.

3 Global structure

For $M < M_{\rm cr} = 4\kappa^3/(G g_{\rm min}^{3/2} L^4)$ there are a bounded branch $(0 \to R_1 \to 0)$ and a bounce branch $(\infty \to R_2 \to \infty)$.

Proposition 2. *(i) $\varepsilon_{\rm out}(R_2) = -1$ always. (ii) $\varepsilon_{\rm out}(R_1) = +1$ iff $E > 2 - 2\gamma$ (regime A). (iii) At $E = 2 - 2\gamma$, $R_1 = 2GM$. (iv) In regime A, the flip radius R_f with $A_{\rm out}(R_f) = 0$ satisfies $R_1 < R_f < R_2$.*

The causal region occupied by the wall is determined by $(\mathrm{sgn} f, \varepsilon, \mathrm{sgn}\dot R)$. We implemented this coordinate-free itinerary and verified it against explicit integration in Kruskal and global de Sitter coordinates: all 15 branches of our atlas agree (Fig. 1). Every inflating branch ends in region III, is invisible from region I, and has a past-singular exterior, in accordance with [2]. The time-symmetric slice at R_2 contains the Einstein–Rosen bridge, a minimal sphere. Hence every classical history through it is singular to the past and to the future, and a transition to it must replace its past.




Figure 2: Left: covering multiplicity of the exterior Euclidean region for a regime-A transition. Right: the Euclidean wall in the cigar for three models.

4 The exterior embedding obstruction

Let $\mathcal{C}$ be the (r, t_E) section of Euclidean Schwarzschild (a smooth cigar), with polar coordinates (ρ, θ), $\theta = t_E/4GM$. The Euclidean wall obeys $R'^2 = V$, $f_{\rm out}\, dt_E/d\tau_E = A_{\rm out}(R)$.

Theorem 3 (no manifold instanton)**.** *Suppose that (H1) the half-bounce runs from R_1 to $R_2 > R_1$ with $R' > 0$; (H2) $A_{\rm out}$ has a single simple zero in (R_1, R_2) with $A_{\rm out}(R_1) > 0 > A_{\rm out}(R_2)$; (H3) the initial and final exterior slices are the totally geodesic slices ($\theta = 0$ ray; broken line through the tip at angles $\theta_D, \theta_D + \pi$); (H4) the exterior region is an oriented surface locally isometric to $\mathcal{C}$, with boundary initial slice $\cup$ wall $\cup$ final slice. Then the local isometry is not injective.*

Proof sketch. Orientation places the region outside the $\varepsilon = +1$ arc, inside the $\varepsilon = -1$ arc, counterclockwise of the initial ray, and clockwise of the final ray. If the map were injective, the multiplicity would be 1 on the region, 0 off it, and would jump by one across each boundary piece. Monotonicity of R puts the $\varepsilon = -1$ arc outside the $\varepsilon = +1$ arc. If either arc wraps by $\geq 2\pi$, a radial segment between consecutive windings has multiplicities 0 and 1 at its ends with no crossing between them. Otherwise, at infinity the multiplicity is 1 exactly on the counterclockwise arc $(0, a)$, $a \equiv \theta_D + \pi$, while just outside the $\varepsilon = -1$ arc it is 0. A case analysis in θ_D produces in each case an angle at which these two requirements conflict. The full proof is in the companion file `NO_GO_THEOREMS.md`. □

For the dS|Schwarzschild model, every regime-A transition satisfies (H1)–(H4). Degree-theory multiplicity maps confirm the result: they take values outside $\{0, 1\}$ (Fig. 2), including negative values for small M, where the wall winds dozens of times around the horizon. Regime-B transitions ($\varepsilon_{\rm out} = -1$ throughout) are also non-manifold numerically, with $|\theta_D| > \pi$ over the whole regime. The interior is not obstructed: models with an $\varepsilon_{\rm in}$ flip have embedded interiors, because there the wall approaches the Euclidean horizon.

5 The canonical exponent and its branch

With the asymptotic Killing time as clock and $H = M$, Hamilton's equation $\partial P/\partial M|_R = dt/dR$ gives

$$P = \frac{R}{G}\Big[\operatorname{artanh}\frac{\dot R}{A_{\rm out}} - \operatorname{artanh}\frac{\dot R}{A_{\rm in}}\Big], \qquad 2\sqrt{f_{\rm in}f_{\rm out}}\cosh\frac{GP}{R} = \varepsilon_{\rm in}\varepsilon_{\rm out}(f_{\rm in}+f_{\rm out}-\kappa^2R^2), \quad (3)$$

together with $\partial P/\partial\Lambda_{\rm in}|_{R,M} = -\frac{1}{8\pi G}\frac{4\pi R^3}{3}\frac{dt_{\rm in}}{dR}$ and $\partial P/\partial\sigma|_{R,M} = -4\pi R^2\frac{d\tau}{dR}$. Under the barrier $P = iP_E$, $P_E = \frac{R}{G}(\Theta_{\rm out}-\Theta_{\rm in})$ with $(A,\sqrt{V}) = \sqrt{f}(\cos\Theta,\sin\Theta)$, and $B = 2\int_{R_1}^{R_2} P_E\,dR$.

Proposition 4 (branch selection)**.** *On (R_1,R_2) we have $f \geq V > 0$ on both sides, so P is analytic there. Its branch points are the turning points and the horizons $f = 0$; the flip radius is a regular point. The WKB phase is the analytic continuation along the barrier, $\Theta = \operatorname{atan2}(\sqrt{V},A) \in [0,\pi]$. The principal branch differs by a horizon monodromy ($P \to P \pm i\pi R/G$, $t \to t \pm i\beta/2$), which is the Euclidean half-turn slicing jump of FGG. The principal branch gives $B < 0$ in 41 of 45 test models.*

6 Closed form at $M = 0$

Theorem 5. *Let $M = 0$, $R_0 = 2\kappa/L^2$, $p = -A_{\rm out}(R_0)$, $q = A_{\rm in}(R_0)$ ($p+q = \kappa R_0 > 0$). Then*

$$S_E = -\frac{\pi R_0^2}{2G}\Big[\frac{1}{1+p}+\frac{1}{1+q}\Big], \qquad B = \frac{\pi R_0^2}{2G}\frac{p+q+2p_+q_+}{(1+|p|)(1+|q|)} = S_E + \frac{1}{4G}\Big(\sum A_{\max} - \sum A_{\min}\Big), \quad (4)$$

the sums running over the extremal spheres of the instanton's time-symmetric slice. If $q < 0$, then $B = B_{\rm CdL}$, the decay exponent of the false vacuum.

The proof uses $J(s) = \int_0^1 x\operatorname{atan2}(\sqrt{1-x^2},sx)dx = \pi/4(1+s)$ ($s \geq 0$), $\pi/2 - \pi/4(1-s)$ ($s < 0$), together with the cap volumes. Gravitational $O(4)$ bounces of a quartic scalar converge to $B_{\rm CdL}$ as $O(w)$ in the wall width (0.18% at $w = 0.005$).

7 First law and Smarr identity

Theorem 6. *Let $\tilde B = B - \mathcal{A}/4G$, where $\mathcal{A}$ collects the endpoint areas ($\pm 4\pi R_i^2$ where the wall is an extremal sphere of its slice, plus $4\pi/\chi^2$ if the final slice contains the de Sitter equator). Then, within a regime,*

$$d\tilde B = \frac{\mathrm{Vol}_E^{\rm in}}{8\pi G}d\Lambda_{\rm in} + A_E^W\,d\sigma - \Delta t_E^W\,dM, \qquad B = -\frac{\Lambda_{\rm in}}{8\pi G}\mathrm{Vol}_E^{\rm in} - \frac{\sigma}{2}A_E^W - \frac{M}{2}\Delta t_E^W + \frac{\mathcal{A}}{4G}, \quad (5)$$

where $\mathrm{Vol}_E^{\rm in}$ is the Euclidean interior 4-volume of the full bounce, A_E^W the wall's Euclidean area, and Δt_E^W the exterior Euclidean Killing time along the wall.

Proof. The first law is the Euclidean continuation of the three Hamilton–Jacobi relations, with endpoint contributions $2P_E(R_i)dR_i = d(\mathcal{A}/4G)$. The Smarr identity follows from Euler's relation for $\tilde B$, which is homogeneous of degree 2 under $(\Lambda,\sigma,M) \to (\lambda^{-2}\Lambda,\lambda^{-1}\sigma,\lambda M)$. □

The first two terms are the on-shell action of the interior region and the wall. The exterior pseudomanifold contributes only $-\frac{M}{2}\Delta t_E^W$, which tends to zero as $M \to 0$ even though the Euclidean winding diverges. Integrating the first law in M from the $M = 0$ instanton gives $B(M) = S_E^{(0)} + \mathcal{A}(M)/4G - \int_0^M \Delta t_E^W\,dM'$. Numerically the identity holds to 2.5×10^{-8} over 45 models.




Figure 3: Left: $B(M)$ for the analytic (solid) and principal (dashed) branches; dots: Theorem 5. Right: the terms of the Smarr identity.

8 Numerical results

Over regime A (laboratory data) the ratio $B/B(M \to 0)$ is bounded below by 0.56 for all $\gamma \in [-0.95, 0.95]$, and by 1 for $\gamma \gtrsim 0.2$, where $B(M \to 0) = B_{\mathrm{CdL}}$ (Fig. 3). For GUT-like parameters, $\sigma^{1/3} = \rho^{1/4} = 10^{16}\,\mathrm{GeV}$ ($M_{\mathrm{cr}} \approx 5.6\,\mathrm{g}$), $B = 133$ only for nanogram (sub-Planckian, see Addendum) masses, while $B \approx 7 \times 10^4$ at $10^{-3} M_{\mathrm{cr}}$ and 4×10^6 at $0.5 M_{\mathrm{cr}}$.

9 Discussion and limitations

The three obstructions to universe creation in an asymptotically flat parent, classical (Farhi–Guth), initial-data (bag-of-gold minimal sphere) and semiclassical (Theorem 3), share one origin: the creation of a closed marginally trapped surface in the exterior. Within the reduced canonical theory, the branch of the exponent is fixed and the exponent is covariantly expressible; the remaining ambiguity is located in the Addendum. Open problems: the exact quantum amplitude (an exact solution is possible at $M = 0$, where the constraint becomes an ODE in the momentum representation); whether Lorentzian path integrals select the same contour; prefactors; thick-wall versions of Theorems 3 and 6; non-spherical perturbations. Our novelty claims are provisional pending comparison with the full texts of [3, 4, 7, 8].

Addendum (22 October 2026): corrections and later results

This manuscript reports Stage One only. Later work (Phases II–VI and the roadmap round; the documents named below are in the accompanying repository) corrects or sharpens the following points.

1. *The ambiguity is located, not removed* (PHASE_III). Analyticity fixes the branch of P (Proposition above), but every formulation hides the FGG datum in one place: the boost-angle convention of a flipped normal, equivalently the relative normalisation of the orientation sectors. Exactly one ambiguity survives: in the single-flip class two consistent exponents remain, B_{an} and $B_{\mathrm{sq}} = \frac{2}{G}\int R \arccos|C|\, dR < B_{\mathrm{an}}$, with a universal Stokes multiplier $\sqrt{2}$. The multiplier is now confirmed by exact integration of the momentum-space squared constraint at $M = 0$.

2. *Where the ambiguity lives* (ROADMAP Theorem S4). A laboratory transition is double-flip, hence unambiguous, iff $\kappa < \chi$ and $M < M_{\mathrm{sd}} = (\chi^2 - \kappa^2)/2G\chi^3$; the boundary is

$R_2 = 1/\chi$. In the single-flip class, a sector-independent probability current gives $2e^{-B_{\rm sq}}$, while $e^{-B_{\rm an}}$ requires an entropy-weighted daughter measure (ROADMAP Q9.3–Q9.4).

3. *T5 status* (ROADMAP Part A). For light walls, T5 is a theorem with an explicit universal law $B = B_{\rm CdL}\,\Phi\big((R_{\rm CdL}/R_M)^2\big) \geq 4\pi^2\sigma^2 R_M^2/\rho$. It implies the frontier $B \geq \frac{4}{3}\sqrt{\pi/2}\sqrt{B_{\rm CdL}(M/m_P)(R_M/\ell$ so in any semiclassically controlled regime $P \leq 10^{-23}$. Elsewhere $B \geq 0.564\,B(0)$ remains numerical.

4. *Masses* (PHASE_V). "Picogram" should read nanogram. These masses are below the Planck mass, so the black hole is not semiclassical. For $M \gtrsim 10 m_P$ the GUT exponent is $B \gtrsim 7 \times 10^4$.

5. *The daughter needs no boundary condition, and only the Killing clock folds back* (ROADMAP Part D, corrected in Addendum 2). The finite-time "edge" of the daughter channel is a coordinate singularity of the static-frame momentum. Along every daughter $dt_\infty/d\tau < 0$ (region III). The roadmap round read this as "time-reversed daughter, Klein-type channel"; that reading is withdrawn (Addendum 2, item 2).

6. *Literature.* The FMP abstract states that agreement with the Euclidean method requires degenerate metrics [4]; Ansoldi and Tanaka resolve slicing discrepancies with a smooth time slice [17]. Both are consistent with the results above. A full-text comparison is still pending.

Addendum 2 (23 October 2026): the transition amplitude, the daughter's clock, verification

1. *A charge selection rule* (ROADMAP_ROUND3, Theorems A1–A2). Assume (P1) the reduced theory, with the analytic shift constraint $\psi(u + ih) + \psi(u - ih) = 2C\psi$, $u = R^2/2$, $h = \hbar G$; (P2) physical states are geometric and analytic in $|\mathrm{Im}\, u| \leq 2h$, with each configuration represented once; (P3) probability is the conserved Casoratian charge $K[\psi, \psi^\sharp]$, with no entropy weights. Then $K = \Phi(e^{2\pi u/h})$ has a unique Laurent series, and a geometric wave with net flip number n contributes only to q^{-n}. Hence K is constant with ordinary flux balance for double-flip transitions, and $K \equiv 0$ (total reflection) for single-flip ones. For double flip, $P = e^{-B_{\rm an}}(1 + O(\hbar))$. The single-flip daughter receives no flux, and $2e^{-B_{\rm sq}}$ (item 2 above) is superseded under P2. The toy $C = -\lambda u$ is solved exactly by $Z_{-iu/h}(i/\lambda h)$. The Hankel function is the unique solution geometric in both regions, its Casoratian vanishes, and $|H^{(1)}(-u)| = e^{\pi u/h}|H^{(1)}(u)|$, so the decay is the analytic-angle exponent exactly, for every h.

2. *The daughter's clock.* Along every branch, on both sides of the wall, both Kruskal null coordinates increase with proper time, because $|\dot{R}| < |A|$ and $\mathrm{sign}(dt/d\tau) = \mathrm{sign}(A)$. The conserved charge of the expanding double-flip daughter is positive, like that of the incident laboratory wave. Only the exterior Killing time folds back. Reversing its orientation is not a reversal of physical time.

3. *Mechanisms.* Dynamical assistance requires driving frequencies $\omega \gtrsim 1/\Delta\tau_E \approx 0.3$–$0.6\,E_{\rm vac}$. No laboratory mechanism beats the frontier; bubbles nucleated during inflation remain a cosmological route only.

4. *Verification.* State level: invisible (Birkhoff). Event level: on the collapse branch a radiating wall's signal has a late tail $\propto e^{-u/4GM}$ carrying $\Gamma\tau_{\rm collapse}$ photons; on the daughter branch it stops at the turning point, with redshift factor $\sqrt{f(R_1)}$.

5. *Proof gaps narrowed.* T5 strip: $\lim_{E\to E_{AB}^{+}} \Delta t_E^W \propto J_0+\pi<0$ for all γ, from $(z^2+z+1)^2 - z^2h'(z) = (z-1)(z+1)(z^2-z+3)(z^2+z+1)/z^2$. NG4: both boundary values ($F(t)$ and $(|J_0|+\pi)/2\pi$) exceed 1, with the interior monotone numerically. An independent high-precision check gives $B_{\rm an} = B(0)\Phi(E)[1+c(E)\kappa^2/\chi^2]$ with $-0.73 < c < 0$, so the frontier carries the factor $1-0.73\kappa^2/\chi^2$ ($P \le 10^{-23}$ unchanged). Euclidean Schwarzschild–de Sitter with conical defects at both horizons has $I(\beta) = -(A_b+A_c)/4G$ for every period.

Addendum 3 (23 October 2026): the search for an accessible mechanism

1. *Classical budget.* In spherical symmetry an anti-trapped sphere forms from laboratory data only if, along some ingoing null ray (with $r_v = \frac{1}{2}$ where it leaves the initial cone), $2Gm/r|_0 + 8\pi G\int r(-T_{uu})\,du/\alpha^2 \ge 1$. The NEC must therefore be violated with an integrated null force of order $c^4/8\pi G = 4.8\times10^{42}\,\mathrm{N}$, independent of size, or the data must already be near their Schwarzschild radius.

2. *Thin walls of any matter.* At a flip of the exterior orientation $V = f_{\rm out}$, so a classical flip happens inside a horizon. With growing radius its past is a singularity, a Cauchy horizon, or a cosmological region. Exotic matter on the wall can produce only a visible bubble. That needs negative asymptotic tension, destabilises Minkowski space, and consumes the parent.

3. *Routes without exotic matter.* A charged exterior changes the exponent by $< 3\%$ but yields a classical route through the Cauchy horizon above $M_{\rm cr} \approx 0.16\,m_P^3/\sqrt{\rho}$. A stabilised seed would convert in e^B/ω. A tilted-quartic field theory shows that the frontier $B_{\rm CdL} = 10$ needs $\lambda \sim 10^4$. With $\lambda \le 4\pi$, $P \le 10^{-476}$.

4. *The only laboratory-scale class* is self-amplifying stable NEC violation (Galilean genesis and its relatives). It meets the budget automatically and forms a daughter deterministically. It requires violation of S-matrix positivity. (The claim that it also requires violation of the GSL is withdrawn in Addendum 4.) A seed bounded by a thin interface in vacuum becomes a visible, parent-consuming bubble, because the interface energy turns negative at $H^2R^2 = 2GM/R$. A hidden daughter needs a throat to open inside a NEC-violating boundary layer.

5. *Observables.* For an explicit 1+1 field in the laboratory state, the collapse branch shows a negative precursor and a switch-on of Hawking emission over $\approx 3.5\times 4GM$. Under a stated state assumption the daughter branch differs by ≈ 0.1 quanta per event. The daughter's inflation is not observable from outside.

Addendum 4 (24 October 2026): genesis and the generalised second law

Violating the NEC does not by itself violate the GSL. We assume the GSL on causal horizons of complete observers and keep global hyperbolicity.

1. *Genesis dichotomy* (spherical symmetry). In a globally hyperbolic spacetime, a relatively compact past slice of an observer forces relatively compact past slices on all earlier Cauchy surfaces. On normal, weakly gravitating laboratory data, a compact past slice puts an ingoing, area-decreasing cross-section on the observer's causal horizon, contradicting the GSL unless it is sub-ℓ_*. Hence a complete daughter observer keeps a non-compact past: the neck never closes, and the daughter cannot end in a de Sitter phase. *An eternal*

daughter and a daughter determined by the laboratory data exclude each other. For a de Sitter parent, $\Lambda_d \le \Lambda_p$ up to $O(\hbar)$. Transient, connected daughters are not constrained.

2. *QNEC cost.* With the semiclassical equations and the integrated QNEC, $r'' \le -(G\hbar/2\pi r)S''_{\rm out}$ on an ingoing ray. Flipping $\theta_- < 0$ therefore needs $-\int (r_0/r)S''_{\rm out}\, d\lambda \ge A_0|\theta_0|/4G\hbar$. That is $\sim S_{\rm BH} \approx 10^{70}$ for a 1 m sphere in flat space, and it is discounted by $\sqrt{f}$ near horizons.

3. *Two dimensions.* With the strong 2D QNEC, $\Theta' \le -\frac{6}{c}S'^2$ exactly, and an anti-trapped episode lasts less than $c/6|\Theta_0|$.

4. *Throats.* In a near-extremal throat, $\sqrt{f} = 2\pi QT\sinh(\rho/Q)$. QNEC-saturating matter of central charge c anti-traps spheres iff $T\sinh(\rho/Q) < (c\eta/12)T_{\rm brk}$, with $T_{\rm brk} = 1/2\pi^2Q^3$. This is confirmed in the exact outgoing Bonnor–Vaidya–Reissner–Nordström metric, where the flip condition $\delta m > rf/2$ is the classical budget. Energy kept in an extremal throat hides every region deeper than $\simeq Q(1 - \frac{1}{2}\ln(2\delta M/Q))$.

5. *Growth.* A Reissner–Nordström–de Sitter T-region grows without bound iff its mass parameter exceeds $\max_{r\ge r_0}(r + Q^2/r - \Lambda r^3/3)$. An extremal-throat bud needs $\Lambda Q^2 > 3/16$ (vacuum energy injected) or 0.2212 (converted locally). A static throat in a fixed vacuum never grows. No product throat exists for $\Lambda Q^2 > 1/4$.

6. *Cauchy-horizon route near extremality.* The charged-bubble window lies in $[1/\sqrt{L}(1+s),\ 1/\sqrt{L}(1-s)]$ ($L = \chi^2+\kappa^2$, $s = \sqrt{1-Q^2/M^2}$), so its width is at most $r_+ - r_- = 2\kappa_- r_-^2$.

The GSL is therefore respected by transient connected daughters and by daughters beyond Cauchy horizons. The remaining obstruction is the vacuum energy $\rho \gtrsim m_P^6/M^2$, which is common to all routes.

Addendum 5 (24 October 2026): a hidden daughter from a Nariai throat

1. *Extremal throats.* In a region of vacuum energy with a radial magnetic field, $f = 1 - C/r + Q^2/r^2 - \Lambda r^2/3$ exactly, so a normal false-vacuum segment is shorter than $0.78 \times 2\ell_H$. An extremal (AdS_2) throat can hide a bud or grow it, but not both.

2. *Nariai throats* (numerical, converged). A Schwarzschild–de Sitter hole with $M = M_N(1-\delta)$ inside a false-vacuum layer whose wall lies in the static band, tipped by outgoing negative energy, grows an inflating daughter behind a persistent trapped neck. All 25 runs obey one law: the daughter grows iff $W = -\int Ne^{-u/\tau}du > 1.85 \times 10^{-3} r_N/G$, with $\tau = 3.2r_N$, in the geodesic frame at $r_N = 3GM$. The gauge-invariant (Misner–Sharp) cost is $\sqrt{f_0}\,W \sim 10^{-2}\sqrt{\delta}\,M_N$. The checkpoint figure "0.6–0.9% of M_N" was the local energy W.

3. *Sources.* With $T_{uu} \ge 0$ the minimal anti-trapped radius cannot decrease, in any parent, so a tip needs NEC violation. A unitary 2D CFT that is in the vacuum before u_0 obeys $\int T_{uu}e^{-(u-u_0)/\tau}du \ge -c/48\pi\tau$ (sharp, from the Fewster–Hollands QEI), so a quantum tip needs $c \ge 0.89(r_N/\ell_P)^2$: the species length reaches the throat. Bunch–Davies vacuum polarisation is stabilising. A softly broken ghost condensate tips the throat with $|\rho + p| \ge 4.6 \times 10^{-3}\rho_f$, 145–430 times below its stability ceiling $|\dot H| \lesssim H^2$ (Creminelli–Luty–Nicolis–Senatore). It is Lorentz-violating, and its GSL is not universal, which Theorem GSL-3 requires of an eternal daughter.

4. *Preparation.* The configuration is normal. The daughter needs false vacuum inside the hole's horizon, i.e. a hole formed inside the false vacuum. The vacuum is tied to the hole by $\rho_f = 1/(24\pi G^2M^2)$: $\rho_f^{1/4} = 0.433\,\mathrm{GeV}\,(M_\odot/M)^{1/2}$, and a GUT vacuum pairs with a 3.7 g hole.

5. *GSL-3 without symmetry.* Compare the observer's past slice with the smallest enclosing normal leaf. At the touching point a past-support hypersurface of the horizon has $\theta \leq \theta_-(\text{leaf}) < 0$, which contradicts the support form of the local GSL.

6. *S3.* $c(E) = 2 - 2mI_0'/I_0 + I_1/I_0$ in closed form, with $c(0^+) = -13/18$ exactly and $c \simeq -(16/\pi)E^{-3/2}$.

Addendum 6 (24 October 2026): one complete route, audited

1. *Independent verification.* A method-of-lines solver with its own data integrator reproduces the Nariai-throat daughter, its hiding and its threshold to within a factor of two.

2. *Correction.* In the Round 6c data the true-vacuum exterior was already trapped (beyond the bridge). Light walls give normal, asymptotically connected data; the exterior collapses into a parent black hole and the daughter grows inside it.

3. *Complete route.* From data containing no black hole (a false-vacuum ball with a light wall, an imploding shell of ordinary radiation, normal everywhere, null energy condition except for the push), the shell forms a black hole inside the false vacuum and a classical push gives a hidden daughter inflating to $r \approx 4500$. With the push injected inside the ball its invariant energy is $\sim 1.5M_N$; the discount needs the source in the band.

4. *Ghost condensate.* In the Nariai region the condensate relaxes as $P' = \epsilon_0 e^{-t/r_N}$, so its NEC phase ends by itself. It tips the throat while every perturbation grows by at most ~ 3 over 3τ for $1.9 \times 10^{-4}/\sqrt{\alpha P''} \leq M^3 r_N \leq 1.7 \times 10^{-2}/\sqrt{\alpha}$.

5. *Generalised entropy.* The parent's event-horizon area never decreases. Daughter-observer horizons need a deficit $\leq 3.3\sqrt{\delta}\, S_N$, which is where the ghost-condensate GSL question (Mukohyama vs. Feldstein) decides the route.

6. *Quantum fields.* For every state of a 2D CFT the strong QNEC gives $\int T_{uu} e^{-u/\tau} du \geq -c/48\pi\tau$, and the fields' own back-reaction raises the threshold faster (no crossing up to $c = 5r_N^2$): quantum fields cannot supply the push.

Addendum 7 (25 October 2026): the condensate as its own source, the entropy of the transition, and the ball

1. *Shift charge.* In the ghost-condensate effective theory $\nabla_\mu(nu^\mu) = \alpha M^2 \Box\theta - V'(\phi)$, with $n = M^4 P' \sqrt{2X}$. Without explicit breaking, the NEC-phase charge of each fluid element is conserved. The higher-derivative term only renormalises Newton's constant, $G_{\text{eff}} = G/(1 - 12\pi G\alpha M^2)$. A charge-free condensate therefore cannot make the pulse; a soft tilt $V' \neq 0$ can.

2. *The condensate tips the throat by itself.* Simulated as a dynamical, back-reacting source (geodesic flow, conserved charge, a clock; constraints converge at second order), its own stress tensor tips the near-Nariai band with no prescribed flux. It obeys the same weighted-deficit law, hides the daughter, and is stable on its measured history. A tilt confined to the false vacuum produces the charge from zero, and the complete route from data without a black hole closes with that self-made push.

3. *Generalised entropy.* For normal data and classical sources, every eternal daughter violates the local GSL. The horizon of a complete daughter observer starts with an ingoing generator on the data surface, where its area decreases. This holds whatever supplies the

NEC violation; the runs give 7–164 nats at $S_N = 942$. The tilt that makes the charge also destabilises the false vacuum after a fixed condensate time. The daughter exits to $\Lambda = 0$ and recollapses inside the parent, which leaves only the parent's horizon (area non-decreasing) as a causal horizon of complete observers.

4. *The ball.* The route's data are normal, so no Farhi–Guth or Penrose obstruction applies. A momentarily static hot ball of radius $0.6\,l_H$, with the scalar in the true vacuum and particles massless at the false vacuum, is normal for strong transitions ($\alpha \gtrsim 2.5$). Heating restores the false vacuum, and free-streaming cooling leaves it there with no kinetic energy. Confined radiation cannot empty the ball: a recollapsing radiation-plus-vacuum ball turns around with $\rho_{\rm rad} \geq \rho_f$. A single relativistic simulation of the emptying, handed off to the route, remains open.

5. *Numerical correction.* The explicit potential step is unstable where $mh\,e^s/2 > 1$. An implicit step removes the instability. Runs with the field exactly at its minimum were unaffected.

Addendum 8 (25 October 2026): formation in one evolution, crunching observers, the condensate beyond its fluid limit, and problems 4 and 5

1. *Thermal formation fails.* A solver that evolves a perfect-fluid plasma with a field-dependent free energy, the scalar, the condensate and gravity together (regular centre, exact hand-off) shows that heating restores the false vacuum only above $E \simeq 1.74$–$4.7\,\rho_f$. In full GR the false-vacuum region then shrinks or is trapped before the radiation leaves. The creation-by-supercooling result of Addendum 7 is withdrawn: its high-temperature free energy was invalid at $m/T \approx 15$.

2. *Observers who end at a singularity.* The boundary of their causal past is the past cone of the endpoint. Its area falls to zero, so no generalised second law can hold for them, in any collapse (measured: the full area, already for a plain black hole). The transient daughter is consistent for every complete observer.

3. *The condensate beyond its fluid limit.* The nonlinear 1+1 effective theory agrees with its exact linearisation ($A = 2 + 3\varepsilon$ with friction $1 + A'/A$). Caustics appear only for seeds 10^3 above quantum. Non-spherical test-field proxies grow no more than in the no-tip controls. The linear stability window, over redshifting k and non-redshifting l modes, is universal in r_N and scales as $\alpha^{-1/2}$: $3.8 \times 10^{-4} \leq M^3 r_N \leq 3.2 \times 10^{-2}$ for an initial charge and 2.6×10^{-3}–3.2×10^{-2} when self-generated ($\alpha = 1$).

4. *One history.* Two independent solvers agree on the route to 10^{-5}. The route from the ball to a hidden daughter runs as one evolution, provided the false vacuum's spinodal comes after the throat forms.

5. *Problems 4 and 5.* The condensate's observational bound $M \lesssim 100$ GeV forces throats of $\gtrsim 10^9 M_\odot$. Any near-Nariai throat needs $E\sqrt{\rho} \geq \frac{1}{3}(8\pi)^{-1/2} E_P \sqrt{\rho_P} \approx 8.9 \times 10^{64}\,\mathrm{J}^{3/2}\mathrm{m}^{-3/2}$, independent of the vacuum scale. The best human capability falls short by 10^{39}.

Addendum 9 (28 October 2026): the false-vacuum region made by the universe, and why no machine can make it

1. *Cosmological creation.* A supercooled first-order transition leaves false-vacuum patches inside a radiation-dominated universe. Evolved with gravity from cosmological light-cone

data (FRW reproduced in conformal and affine gauges), a patch collapses into a black hole below, and inflates above, a critical radius of 0.57–0.59 l_H. This is independent of resolution, and the band is 0.42–0.65 l_H over vacuum fractions 0.5–1. The ball the route starts from (0.595 l_H) is the universe's near-critical patch.

2. *A hidden daughter, consistent with the second law.* For a supercritical patch, a trapped true-vacuum region forms outside the wall while the false vacuum inflates behind it (the Garriga–Vilenkin–Zhang structure). The parent's event horizon area never decreases, and no daughter observer's horizon loses area. The eternal-daughter obstruction of normal data does not apply to cosmological, anti-trapped data. The condensate tilt turns a collapsing near-critical patch into a daughter.

3. *External assembly.* Delivering the energy of a region of compactness $2GE/(Rc^4) \approx 2/3$ from outside within $\alpha R/c$ needs $P \gtrsim (k/2\alpha)\, c^5/G \approx 1.2 \times 10^{52}$ W ($\alpha = 1$); holding it static needs a stress $\approx (k^2/4)\, c^4/G \approx 1.3 \times 10^{43}$ N. In quantum form, $E_{\text{total}} E_{\text{quantum}} \geq 6.3 \times 10^{17}\,\text{J}^2$, which every audited machine fails. This is a conditional bound on two strategies, not a proof that no machine can exist (corrected in Addendum 10).

Addendum 10 (28 October 2026): what actually blocks a controllable mechanism

1. *Correction.* Addendum 9's power and force estimate bounds external assembly and static holding only. The maximum force and luminosity are conjectures with counterexamples in general relativity, and the holding stress lies below $c^4/4G$. The claim that no machine of any kind can build the region is withdrawn.

2. *What holds for every strategy.* By the time-reversed Penrose theorem with the null energy condition, a closed anti-trapped surface forces past null incompleteness, and an inflating daughter contains one. A machine of NEC-respecting matter acting on a universe with a regular past cannot create a daughter, whether by rapid, slow, dynamical or triggered means. The escapes are NEC violation, a cosmological past singularity, or a quantum transition.

3. *Triggers and backgrounds.* In a cosmological (anti-trapped) background, a positive-energy trigger of about 10% of a supercooled patch's vacuum energy turns its collapse into a hidden daughter. In a normal background the trigger must violate the NEC. With the observationally allowed condensate, the required near-Nariai background has mass $r_N c^2/3G \geq 2.5 \times 10^9 M_\odot$, whatever its formation history. Black-hole-assisted creation needs black holes of the same mass class. Smaller backgrounds would need a condensate that exists only inside the false vacuum, which remains an open model.

Addendum 11 (28 October 2026): seven routes without a manufactured ball, and the bridge invariant

1. *Curvature activation.* For $S = \frac{1}{16\pi}\int \sqrt{-g}\,[R - 2(\nabla\phi)^2 - 2V + \lambda^2 f(\phi)\mathcal{G}]$ the Gauss–Bonnet stress in double-null spherical symmetry obeys $\nabla^a \Gamma_{ab} = -\frac{1}{2}\mathcal{G}\nabla_b F$ ($F = \lambda^2 f$, verified symbolically). On a vacuum-energy throat the tipping flux is $N = r_N^2 \phi_u^2 + 2F_{uu}$, and the Gauss–Bonnet part is a total derivative, so its weighted deficit is negative whenever F grows. Since $\mathcal{G} = C^2 + 8\Lambda^2/3 > 0$ there, a field that the coupling destabilises raises F: curvature activation pushes throats towards black holes. Nonlinear Kantowski–Sachs evolutions from exact Nariai data give no daughter in 152 runs (black holes, or a degenerate kinetic matrix).

2. *Monopoles, breathing bubbles, resonance, membranes.* A general thin shell (wall, dust, charge, gas) reproduces the monopole threshold $\eta_c \approx 0.25$–$0.40\, m_P$. At the monopole's own mass the exterior has no bridge ($M \ll Q$), and the daughter branch is disconnected from our universe; the connected channel needs $M > Q$ and has $B \geq 3 \times 10^3$. Breathing bubbles need negative-energy dust, and every well crosses zero net surface energy at the speed of light. Resonance turns $B_1 + B_2$ into at best $\max(B_2, B_1 - B_2)$. Four-form membranes are Farhi–Guth walls with $B \simeq B_{\mathrm{CdL}}$.

3. *Condensates and collisions.* Every NEC-violating ghost-condensate region is charged; gauged, it is screened, and the tipping window is empty by five orders of magnitude. A vacuum-dependent scale lowers the background to $9 \times 10^8\,\mathrm{kg}$ (conditional). Where $f_{\mathrm{out}} > 0$ the sign of β_{out} cannot change, so collision pockets in a normal parent never reach the hidden branch.

4. *Bridge invariant.* At a hidden daughter's wall $\beta_{\mathrm{out}} < 0$, so the area radius has a minimum between the wall and our asymptotic region, where $2Gm/c^2r \geq 1$. From regular data its first appearance is a black-hole formation event with $E\sqrt{\bar\rho} = \sqrt{3/32\pi}\, c^6/G^{3/2} \approx 2.3 \times 10^{65}\,\mathrm{J}^{3/2}\mathrm{m}^{-3/2}$, whatever the route or vacuum scale; the best technology is $10^{39.5}$ short. A barely metastable false vacuum with a light wall gives a semiclassically unsuppressed channel at TeV energies, but its bridge is $\sim 10^{-16}$ Planck lengths across, outside every controlled regime.

Addendum 12 (28 October 2026): geometry, stored and supplied energy; fifteen loopholes

1. *What the invariant bounds.* $E\sqrt{\bar\rho} = \sqrt{3/32\pi}\, c^6 G^{-3/2} k^{3/2}$ with $k = 2GE/c^4R$; the bridge invariant is its value at $k = 1$. The spherical identity $\partial_u m = r^2 e^{-2s}(G_{uv} r_u - G_{uu} r_v)$ shows that $\bar\rho$ is the areal mean of the energy density in the frame with no radial flux. At the first marginal sphere the peak density is at least $\bar\rho$, and negative energy raises the positive energy needed. With $\rho \geq 0$ the invariant bounds stored energy; it does not bound the energy an apparatus supplies. A small supplied energy needs a natural precursor that is stable, older than the universe ($M > 1.7 \times 10^{11}\,\mathrm{kg}$) and within $\delta \lesssim 10^{-8}$ of threshold.

2. *Classical loopholes.* A core with effective Planck mass $\sqrt{F} M_P$ lowers the requirement by $F^{3/4}$ but has compactness at least $\frac{3}{4}\ln^2 F$ (capacitance bound), horizon-class for $F < 0.32$. Finite pockets expand by $\sim R/l_f$. Accretion falls short of the partner density; the most compact bag star is sub-Hubble; evaporating-hole catalysis has exponent $\hat\Sigma(l_f/l_P)^2$.

3. *Quantum channels.* For Euclidean axion wormholes with Λ, the action tends to the no-boundary value $-12\pi^2/\Lambda$ as the throat pinches off, and the excess is $\Delta S = 3\pi^3 a_0^2$ ($8\pi G = 1$), independent of Λ: staying connected costs action in proportion to the throat area.

4. *Conditional doors.* A Frolov–Markov–Mukhanov interior joins Schwarzschild to de Sitter at $r_0 = (2Ml^2)^{1/3}$; if quantum gravity does this, every black hole contains a daughter. With $n \geq 4$ extra dimensions and $M_D \approx 6$–$20\,\mathrm{TeV}$ (allowed), a 100 TeV collider makes black holes, but at $R_S M_D \approx 1$. The Round 6c partner vacuum is corrected to $\rho_f = 1/(72\pi M^2)$.

References

[1] S. K. Blau, E. I. Guendelman, A. H. Guth, Phys. Rev. D **35**, 1747 (1987).

[2] E. Farhi, A. H. Guth, Phys. Lett. B **183**, 149 (1987).

[3] E. Farhi, A. H. Guth, J. Guven, Nucl. Phys. B **339**, 417 (1990).

[4] W. Fischler, D. Morgan, J. Polchinski, Phys. Rev. D **41**, 2638 (1990); **42**, 4042 (1990).

[5] S. Coleman, F. De Luccia, Phys. Rev. D **21**, 3305 (1980).

[6] W. Israel, Nuovo Cim. B **44**, 1 (1966).

[7] S. de Alwis, F. Muia, V. Pasquarella, F. Quevedo, Fortschr. Phys. (2020), arXiv:1909.01975.

[8] S. Céspedes, S. de Alwis, F. Muia, F. Quevedo, Phys. Rev. D **109**, 105027 (2024), arXiv:2307.13614.

[9] B. Freivogel, V. Hubeny, A. Maloney, R. Myers, M. Rangamani, S. Shenker, JHEP **03** (2006) 007.

[10] Z. Fu, D. Marolf, JHEP **11** (2019) 040.

[11] A. Aguirre, M. C. Johnson, Phys. Rev. D **72**, 103525 (2005).

[12] P. Chen, Y.-C. Hu, D.-h. Yeom, Phys. Rev. D **94**, 024044 (2016).

[13] A. Ulvestad, A. Albrecht, Phys. Rev. D **85**, 103527 (2012).

[14] A. C. Wall, Class. Quant. Grav. **30**, 165003 (2013).

[15] R. Penrose, Phys. Rev. Lett. **14**, 57 (1965).

[16] S. Parke, Phys. Lett. B **121**, 313 (1983).

[17] S. Ansoldi, T. Tanaka, J. Exp. Theor. Phys. **120**, 460 (2015), arXiv:1410.6202.

[18] A. C. Wall, arXiv:1105.3520 (2011).

[19] R. Bousso, Phys. Rev. Lett. (2025), arXiv:2501.17910.

[20] J. Maldacena, A. Milekhin, F. Popov, arXiv:1807.04726.

[21] D. Gajic, J. Luk, Pure Appl. Anal. **1**, 263 (2019), arXiv:1709.09137.

[22] R. Bousso, S. W. Hawking, Phys. Rev. D **57**, 2436 (1998); R. Bousso, Phys. Rev. D **58**, 083511 (1998).

[23] C. J. Fewster, S. Hollands, Rev. Math. Phys. **17**, 577 (2005).

[24] P. Creminelli, M. A. Luty, A. Nicolis, L. Senatore, JHEP **12** (2006) 080.

[25] G. J. Galloway, Ann. Henri Poincaré **1**, 543 (2000).

[26] S. Mukohyama, arXiv:0901.3595 (2009).

[27] B. Feldstein, Phys. Rev. D **80**, 044020 (2009).

[28] N. Arkani-Hamed, H.-C. Cheng, M. A. Luty, S. Mukohyama, T. Wiseman, JHEP **01** (2007) 036.

[29] J. Garriga, A. Vilenkin, J. Zhang, JCAP **02** (2016) 064.

[30] H. Deng, A. Vilenkin, JCAP **12** (2017) 044.

[31] K. Kawana, T. Kim, P. Lu, arXiv:2212.14037.

[32] G. W. Gibbons, Found. Phys. **32**, 1891 (2002).

[33] CMS Collaboration, JHEP **11** (2018) 042.

[34] A. Jowsey, M. Visser, arXiv:2102.01831 (2021).

[35] V. Cardoso, T. Ikeda, C. J. Moore, C.-M. Yoo, Phys. Rev. D **97**, 084013 (2018).

[36] N. Oshita, J. Yokoyama, Phys. Lett. B **785**, 197 (2018).

[37] D. D. Doneva, S. S. Yazadjiev, Phys. Rev. Lett. **120**, 131103 (2018).

[38] J. L. Ripley, F. Pretorius, Phys. Rev. D **99**, 084014 (2019).

[39] M. A. Cuyubamba, R. A. Konoplya, A. Zhidenko, Phys. Rev. D **98**, 044040 (2018).

[40] N. Sakai, K. Nakao, H. Ishihara, M. Kobayashi, gr-qc/0602084 (2006).

[41] E. I. Guendelman, N. Sakai, arXiv:0803.0268 (2008).

[42] H.-C. Cheng, M. A. Luty, S. Mukohyama, J. Thaler, JHEP **05** (2006) 076.

[43] P. M. Saffin, A. Padilla, E. J. Copeland, arXiv:0804.3801 (2008).

[44] M. C. Johnson, I.-S. Yang, arXiv:1005.3506 (2010).

[45] J. D. Brown, C. Teitelboim, Nucl. Phys. B **297**, 787 (1988).

[46] J. Marsden, J. C. Aurrekoetxea, K. Clough, P. G. Ferreira, Phys. Rev. D **111**, L041501 (2025).

[47] G. Lavrelashvili, J.-L. Lehners, arXiv:2603.11003 (2026).

[48] M. Sasieta, B. Swingle, A. Vilar López, Phys. Rev. Lett. **137**, 041501 (2026).

[49] G. Huisken, T. Ilmanen, J. Diff. Geom. **59**, 353 (2001).

[50] V. P. Frolov, M. A. Markov, V. F. Mukhanov, Phys. Lett. B **216**, 272 (1989).

[51] CMS Collaboration, arXiv:2107.13021.