

# Numerical Analysis

## Assignment 9

Chuan Lu

October 20, 2017

**Problem 1.** Problem 4.1, Page 239.

*Solution.* When  $n = 4$ ,

$$\begin{aligned} p_4(x) &= \sum_{k=0}^4 C_4^k f\left(\frac{k}{4}\right) x^k (1-x)^{4-k} = f(0)(1-x)^4 + 4f\left(\frac{1}{4}\right)x(1-x)^3 + 6f\left(\frac{1}{2}\right)x^2(1-x)^2 + 4f\left(\frac{3}{4}\right)x^3(1-x) + f(1)x^4 \\ &= (6 - 4\sqrt{2})x^4 + 8\sqrt{2}x^3 - (12 + 6\sqrt{2})x^2 + 2\sqrt{2}x + 6 \\ &= (6 - 4\sqrt{2})\left(x - \frac{1}{2}\right)^4 - 3\left(x - \frac{1}{2}\right)^2 + 3 + \frac{\sqrt{2}}{4} \end{aligned}$$

And the fourth degree Taylor polynomial expanded about  $\frac{1}{2}$  is

$$\begin{aligned} q_4(x) &= f\left(\frac{1}{2}\right) + f'\left(\frac{1}{2}\right)\left(x - \frac{1}{2}\right) + \frac{1}{2}f''\left(\frac{1}{2}\right)\left(x - \frac{1}{2}\right)^2 + \frac{1}{6}f^{(3)}\left(\frac{1}{2}\right)\left(x - \frac{1}{2}\right)^3 + \frac{1}{24}f^{(4)}\left(\frac{1}{2}\right)\left(x - \frac{1}{2}\right)^4 \\ &= 1 - \frac{1}{2}\pi^2\left(x - \frac{1}{2}\right)^2 + \frac{1}{24}\pi^4\left(x - \frac{1}{2}\right)^4 \end{aligned}$$

When  $x \rightarrow \frac{1}{2}$ ,  $q_4(x) \rightarrow 1 = f\left(\frac{1}{2}\right)$ , while  $p_4(x) \rightarrow 3 + \frac{\sqrt{2}}{4}$ . Thus Bernstein polynomials are poor approximations.

**Problem 2.** Problem 4.5, Page 240

**Problem 3.** Problem 4.6, Page 240

**Problem 4.** Problem 4.10, Page 241

(a). The linear Taylor polynomial to  $f = \ln(x)$  expanding about  $\frac{3}{2}$  is

$$p_1(x) = \frac{3}{2} + \frac{2}{3}\left(x - \frac{3}{2}\right) = \frac{2}{3}x + \frac{1}{2}.$$

The error of is as follows, the left is error graph for (a) and the right is for (b).

(b). The linear minimax approximation to  $f$  is

$$p_2(x) =$$

**Problem 5.** Problem 4.12, Page 241

*Solution.* The linear least square approximation to  $f$  is

