

Introduction to Analysis I
Homework 5
Sunday, October 8, 2017

Instructions: This and all subsequent homeworks must be submitted written in L^AT_EX.
If you use results from books, Royden or others, please be explicit about what results you are using.

Homework 5 is due by midnight, Saturday, October 21.

1. (Problem 24, Page 64) Let I be an interval in \mathbb{R} and let $f : I \rightarrow \mathbb{R}$ be increasing. Show that f is measurable by first showing that, for each natural number n , the strictly increasing function $x \rightarrow f(x) + x/n$ is measurable, and then taking pointwise limits.

Collaborators: None

Solution: Denote $f_n(x) = f(x) + \frac{x}{n}$. Then since $f : I \rightarrow \mathbb{R}$ is increasing, $f_n(x)$ is strictly increasing on $I \in \{[a, b], [a, b), (a, b], (a, b)\}$.

For each fixed number c , if $\exists x_0 \in I$, s.t. $f(x_0) = c$, then the set

$$\{x \mid f(x) < c\} = (a, x_0),$$

and the left side is the same as the set I , and it is an interval in \mathbb{R} , hence is measurable. Thus f_n is measurable for every n .

For each $x \in I$,

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) + \lim_{n \rightarrow \infty} \frac{x}{n} = f(x),$$

hence f_n converges to f pointwise. Using Proposition 9, we know f is measurable.

2. (Problem 8, Page 343) Let (X, \mathcal{M}, μ) be a measure space. The measure μ is said to be **semifinite** provided each measurable set of infinite measure contains measurable sets of arbitrarily large finite measure.
- (a) Show that each σ -finite measure is semifinite.
 - (b) For $E \in \mathcal{M}$, define $\mu_1(E) = \mu(E)$, if $\mu(E) < \infty$, and if $\mu(E) = \infty$, define $\mu_1(E) = \infty$ if E contains measurable sets of arbitrarily large finite measure and $\mu_1(E) = 0$ otherwise. Show μ_1 is a semifinite measure: it is called the semifinite part of μ .
 - (c) Find a measure μ_2 on \mathcal{M} that only takes the values 0 and ∞ and $\mu = \mu_1 + \mu_2$.

Collaborators: None

Solution: (a) If μ is a σ -finite measure, then $X = \bigcup_{n=1}^{\infty} X_n$, where X_n has finite measure. Assume E is a measurable set with infinite measure, then since $E \subset X$, $E = E \cap X = \bigcup_{n=1}^{\infty} E \cap X_n$. For $\forall M > 0$, there must exist $N > 0$, s.t. $\mu\left(\bigcup_{n=1}^N E \cap X_n\right) > M$, otherwise $\mu(E) \leq \sum_{n=1}^{\infty} \mu(E \cap X_n) < M_0$ for some $M_0 > 0$, which makes a contradiction with $\mu(E) = \infty$. Besides, we know from $\mu(X_n) < \infty$, that $\mu\left(\sum_{n=1}^N (E \cap X_n)\right) < \infty$. Thus μ is semifinite.

(b) Suppose E is a measurable set of infinite measure in the measure space (X, \mathcal{M}, μ_1) . Then according to the definite of μ_1 , E contains measurable sets of arbitrarily large finite measure. Thus μ_1 is semifinite.

(c) Define μ_2 like this: If E can be represented as countable union of measurable sets with finite measure, then $\mu_2(E) = 0$. Otherwise $\mu_2(E) = \infty$.

First, we show that $\mu = \mu_1 + \mu_2$. If $\mu(E) < \infty$, then $E = \bigcup_{n=1}^{\infty} E_n$, in which $E_n = \emptyset$ for $n \geq 2$. Thus $\mu_2(E) = 0 = \mu(E) - \mu_1(E)$. Now we assume $\mu(E) = \infty$. If E can be represented as countable union of measurable sets with finite measure, that is, $E = \bigcup_{n=1}^{\infty} E_n$ with $\mu(E_n) < \infty$, then with the continuity of measure, $\forall M > 0, \exists N > 0$, s.t. $\mu\left(\bigcup_{n=1}^N E_n\right) > M$. Since \mathcal{M} is a σ -algebra, it shows that E contains measurable sets of arbitrarily large finite measure. Thus $\mu(E) = \infty = \mu_1(E) + 0 = \mu_1(E) + \mu_2(E)$. If E can't be represented as countable union of measurable sets with finite measure, then $\mu(E) = \infty = \mu_1(E) + \mu_2(E)$.

Now we show that μ_2 is a measure. In fact, we only need to show that $\mu_2(\emptyset) = 0$, and μ_2 is countably additive. First, from the definition we know $\mu(\emptyset) = 0 < \infty$, so $\mu_2(\emptyset) = 0$. Suppose $\{E_n\}_{n=1}^{\infty}$ is a sequence of disjoint measurable sets. If $\exists i$, s.t. $\mu_2(E_i) = \infty$. Then E_i can't be represented as countable union of measurable sets with finite measure, thus $\bigcup_{n=1}^{\infty} E_n$ can't be represented as countable union of measurable sets with finite measure. Then

$$\mu_2\left(\bigcup_{n=1}^{\infty} E_n\right) = \infty = \sum_{n=1}^{\infty} \mu_2(E_n).$$

Now we suppose all $\mu_2(E_n) < \infty$. In this case, since μ_2 takes value only in $\{0, \infty\}$, we have $\mu_2(E_n) = 0$. In this case, all E_n is a countable union of measurable sets with finite measure, then $\bigcup_{n=1}^{\infty} E_n$ is a countable union of measurable sets with finite measure. Thus

$$\mu_2\left(\bigcup_{n=1}^{\infty} E_n\right) = 0 = \sum_{n=1}^{\infty} \mu_2(E_n).$$

3. (Problem 9, Page 343) Prove Proposition 3, that is, show that \mathcal{M}_0 is a σ -algebra, μ_0 is properly defined and $(X, \mathcal{M}_0, \mu_0)$ is complete. In what sense is \mathcal{M}_0 minimal?

Collaborators:

Solution:

4. (Problem 10, Page 343) If (X, \mathcal{M}, μ) is a measure space, we say that a subset E of X is **locally measurable** provided for each $B \in \mathcal{M}$ with $\mu(B) < \infty$, the intersection $E \cap B$ belongs to \mathcal{M} . The measure μ is called **saturated** provided every locally measurable set is measurable.
- Show that each σ -finite measure is saturated.
 - Show that the collection \mathcal{C} of locally measurable sets is a σ -algebra.
 - Let (X, \mathcal{M}, μ) be a measure space and \mathcal{C} the σ of locally measurable sets. For $E \in \mathcal{C}$, define $\bar{\mu}(E) = \mu(E)$ if $E \in \mathcal{M}$ and $\bar{\mu}(E) = \infty$ if $E \notin \mathcal{M}$. Show that $(X, \mathcal{C}, \bar{\mu})$ is a saturated measure space.
 - If μ is semifinite and $E \in \mathcal{C}$, set $\underline{\mu}(E) = \sup\{\mu(B) \mid B \in \mathcal{M}, B \subseteq E\}$. Show that $(X, \mathcal{C}, \underline{\mu})$ is a saturated measure space and that $\underline{\mu}$ is an extension of μ . Give an example to show that $\bar{\mu}$ and $\underline{\mu}$ may be different.

Collaborators:

Solution:

5. (Problem 18, Page 373) Let $\{u_n\}$ be a sequence of nonnegative measurable functions on X . For $x \in X$, define $f(x) = \sum_{n=1}^{\infty} u_n(x)$. Show that

$$\int_X f \, d\mu = \sum_{n=1}^{\infty} \left[\int_X u_n \, d\mu \right].$$