

**CLASS: A**

**PROTOCOL(S) USED IN:**

Seizure

**PHARMACOLOGY AND ACTIONS:**

Non-narcotic analgesic and antipyretic

**INDICATIONS:**

Reduction of fever associated with febrile seizures in the pediatric patient.

**CONTRAINDICATIONS:**

**A. Hypersensitivity**

**B. DO NOT use with any other products that contain acetaminophen**

**SIDE EFFECTS AND NOTES:**

May be administered via rectal suppository (same dose) if patient is vomiting, the patient's gag reflex is absent or in question or the patient is not alert. Can be given up to every 4 hours. Do not exceed five doses a day.

**DOSING/DOSING CHART:**

**A. 15 mg/kg PO/PR suppository or**

**B. See chart below for other concentrations/forms**

| Weight   | Age      | 80mg Tablets | 160mg/tsp Elixir | 80mg/0.8ml Drops | Mg    |
|----------|----------|--------------|------------------|------------------|-------|
| 6-11 lbs | 0-3 mos  | ---          | ¼ tsp            | .4ml             | 40mg  |
| 12-17lbs | 4-11mos  | ---          | ½ tsp            | .8ml             | 80mg  |
| 18-23lbs | 11-23mos | 1 ½ tab      | ¾ tsp            | 1.2ml            | 120mg |
| 24-35lbs | 2-3yrs   | 2 tabs       | 1 tsp            | 1.6ml            | 160mg |
| 36-47lbs | 4-5yrs   | 3 tabs       | 1 ½ tsp          | 2.4ml            | 240mg |
| 48-59lbs | 6-8yrs   | 4 tabs       | 2 tsp            | 3.2ml            | 320mg |
| 60-71lbs | 9-10yrs  | 5 tabs       | 2 ½ tsp          | 4.0ml            | 400mg |
| 72-95lbs | 11-12yrs | 6 tabs       | 3 tsp            | 4.8ml            | 480mg |