


Neonatal Resuscitation – 10.160



NOTES & PRECAUTIONS:

- A. Neonatal age is birth to 28 days.
- B. Do not use atropine in neonatal resuscitation.
- C. If meconium is lightly stained and infant is vigorous (strong respiratory effort, good muscle tone, heart rate > 100 bpm) endotracheal suctioning should not be performed.
- D. An infant may need resuscitation if intrapartum risk factors for asphyxia are present (prolapsed cord, painful bleeding, prolonged rupture of membranes, maternal fever, multiple births, abnormal presentation, maternal hypotension or seizure)

KEY CONSIDERATIONS:

Fetal presentation, recent trauma, maternal health/risk factors, maternal medications, previous birth difficulties, APGAR

APGAR SCORE:	0	1	2
Appearance	Blue/Pale	Body pink, blue extremities	Completely pink
Pulse	Absent	Slow (< 100 bpm)	≥ 100 bpm
Grimace	No response	Grimace	Cough or sneeze
Activity	Limp	Some flexion	Active motion
Respirations	Absent	Slow, irregular	Good, crying

Newborn Target Spo2 after birth:

1 min	60-65%
2 min	65-70%
3 min	70-75%
4 min	75-80%
5 min	80-85%
10 min	85-95%