

# Lower bounds on the sizes of integer programs without additional variables

Volker Kaibel · Stefan Weltge

Springer-Verlag Berlin Heidelberg and Mathematical Optimization Society 2014

# Presentation structure

- ① Simple facts on integer linear programming and combinatorial optimization
- ② Initial question and definitions
  - ① How many inequalities needed to formulate a specific combinatorial problem as an ILP?
  - ② No "extra" variables allowed.
- ③ Main technique and results
  - ① Hiding sets
  - ② Exponential lower bounds for many problems.
- ④ Later results on the topic

# Solving combinatorial optimization problems by linear programming

## Typical combinatorial optimization problem

- Finite ground set  $E$ .
- Feasible solution: Any set  $F \subseteq E$  with some property.
- Vector  $c \in \mathbb{R}^E$ .
- Objective value of feasible solution  $F$  is  $\sum_{e \in F} c_e$ .

# Solving combinatorial optimization problems by linear programming

## Typical combinatorial optimization problem

- Finite ground set  $E$ .
- Feasible solution: Any set  $F \subseteq E$  with some property.
- Vector  $c \in \mathbb{R}^E$ .
- Objective value of feasible solution  $F$  is  $\sum_{e \in F} c_e$ .

## Typical solution approach

- Make ordering for  $e_1, e_2, \dots$  of  $E$
- Identify each  $F$  with its characteristic vector  $\chi(F) \in \{0, 1\}^E$
- $(\chi(F))_i = 1 \iff e_i \in F$
- Vector  $c \in \mathbb{R}^E$ .
- Objective value of feasible solution  $F$  is  $c^T \chi(F)$ .

# Solving combinatorial optimization problems

Need algebraic description of set  $X := \{\chi(F) : F \subseteq E \text{ feasible}\}$ .

## Standard approach

$X = \{x \in \mathbb{Z}^E : Ax \leq b\}$ , for well chosen  $A, b$

# Solving combinatorial optimization problems

Need algebraic description of set  $X := \{\chi(F) : F \subseteq E \text{ feasible}\}$ .

## Standard approach

$X = \{x \in \mathbb{Z}^E : Ax \leq b\}$ , for well chosen  $A, b$

## Example

Traveling salesman problem:

- Let  $K_n = (V_n, E_n)$  be the undirected complete graph on  $n$  nodes.
- Let  $X = \text{STSP}_n$  be the set of characteristic vectors of Hamiltonian cycles in  $K_n$ .
- Formulate  $\text{STSP}_n$  as

# Solving combinatorial optimization problems

Need algebraic description of set  $X := \{\chi(F) : F \subseteq E \text{ feasible}\}$ .

## Standard approach

$X = \{x \in \mathbb{Z}^E : Ax \leq b\}$ , for well chosen  $A, b$

## Example

Traveling salesman problem:

- Let  $K_n = (V_n, E_n)$  be the undirected complete graph on  $n$  nodes.
- Let  $X = \text{STSP}_n$  be the set of characteristic vectors of Hamiltonian cycles in  $K_n$ .
- Formulate  $\text{STSP}_n$  as

$$\text{STSP}_n = \{x \in \{0, 1\}^{E_n} : x(\delta(S)) \geq 2, \forall S \neq \emptyset, V_n \quad (1)$$

$$x(\delta(v)) = 2 \forall v \in V_n \} \quad (2)$$

## Standard approach

$X = \{x \in \mathbb{Z}^E : Ax \leq b\}$ , for well chosen  $A, b$

- Feasible solution: Any set  $F \subseteq E$  with some property.
- Vector  $c \in \mathbb{R}^E$ .
- Objective value of feasible solution  $F$  is  $\sum_{e \in F} c_e$ .



## Standard approach

$X = \{x \in \mathbb{Z}^E : Ax \leq b\}$ , for well chosen  $A, b$

- Feasible solution: Any set  $F \subseteq E$  with some property.
- Vector  $c \in \mathbb{R}^E$ .
- Objective value of feasible solution  $F$  is  $\sum_{e \in F} c_e$ .

## Example

Traveling salesman problem:

- Let  $K_n = (V_n, E_n)$  be the undirected complete graph on  $n$  nodes.
- Let  $X = \text{STSP}_n$  be the set of characteristic vectors of Hamiltonian cycles in  $K_n$ .
- Formulate  $\text{STSP}_n$  as

$$\text{STSP}_n = \{x \in \{0, 1\}^{E_n} : x(\delta(S)) \geq 2 \forall S : \emptyset \neq S \subset V_n \quad (3)$$

$$x(\delta(v)) = 2 \forall v \in V_n \} \quad (4)$$

## Standard approach

$X = \{x \in \mathbb{Z}^E : Ax \leq b\}$ , for well chosen  $A, b$

- Feasible solution: Any set  $F \subseteq E$  with some property.
- Vector  $c \in \mathbb{R}^E$ .
- Objective value of feasible solution  $F$  is  $\sum_{e \in F} c_e$ .

## Example

Traveling salesman problem:

- Let  $K_n = (V_n, E_n)$  be the undirected complete graph on  $n$  nodes.
- Let  $X = \text{STSP}_n$  be the set of characteristic vectors of Hamiltonian cycles in  $K_n$ .
- Formulate  $\text{STSP}_n$  as

$$\text{STSP}_n = \{x \in \{0, 1\}^{E_n} : x(\delta(S)) \geq 2 \forall S : \emptyset \neq S \subset V_n \quad (3)$$

$$x(\delta(v)) = 2 \forall v \in V_n \} \quad (4)$$

## Example

Traveling salesman problem:

- Let  $K_n = (V_n, E_n)$  be the undirected complete graph on  $n$  nodes.
- Let  $X = \text{STSP}_n$  be the set of characteristic vectors of Hamiltonian cycles in  $K_n$ .
- Formulate  $\text{STSP}_n$  as

$$\text{STSP}_n = \{x \in \{0, 1\}^{E_n} : x(\delta(S)) \geq 2, \forall S \neq \emptyset, V_n \quad (5)$$

$$x(\delta(v)) = 2 \forall v \in V_n \} \quad (6)$$

## Example

Traveling salesman problem:

- Let  $K_n = (V_n, E_n)$  be the undirected complete graph on  $n$  nodes.
- Let  $X = \text{STSP}_n$  be the set of characteristic vectors of Hamiltonian cycles in  $K_n$ .
- Formulate  $\text{STSP}_n$  as

$$\text{STSP}_n = \{x \in \{0, 1\}^{E_n} : x(\delta(S)) \geq 2, \forall S \neq \emptyset, V_n \quad (5)$$

$$x(\delta(v)) = 2 \forall v \in V_n \} \quad (6)$$

- Uses exponentially many (in  $n$ ) linear inequalities.
- Nevertheless, computationally efficient (both in theory and practise).  
The separation problem associated with these inequalities can be solved efficiently.

## Initial question

Do all formulations of the form  $STSP_n = \{x \in \mathbb{Z}^E : Ax \leq b\}$  need exponential size?

Question answered for number of linear inequalities.

## Motivators

- Pure mathematical curiosity
- Simplicity of implementation may be more important issue than efficiency.
  - A small ILP-formulation that can be fed immediately into a black box ILP-solver is likely to be preferred over one for which a separation procedure has to be implemented and linked to the solver.

## Initial question

Do all formulations of the form  $STSP_n = \{x \in \mathbb{Z}^E : Ax \leq b\}$  need exponential size?

Question answered for number of linear inequalities.

## Motivators

- Pure mathematical curiosity
- Simplicity of implementation may be more important issue than efficiency.
  - A small ILP-formulation that can be fed immediately into a black box ILP-solver is likely to be preferred over one for which a separation procedure has to be implemented and linked to the solver.

We extend this question to many combinatorial problems.

# Extra inequalities and variables

- As for extra inequalities and variables?

# Extra inequalities and variables

- As for extra inequalities and variables?
- Formulations for  $X=\text{STSP}_n = \{x \in \mathcal{Z}^E : \exists y \in \mathcal{Z}^m : Ax + By \leq b\}$  of polynomial size known.
- Actually, using the reduction proving integer programming is NP-hard, this holds for any  $L \in \text{NP}$ .



# Extra inequalities and variables

- As for extra inequalities and variables?
- Formulations for  $X=\text{STSP}_n = \{x \in \mathbb{Z}^E : \exists y \in \mathbb{Z}^m : Ax + By \leq b\}$  of polynomial size known.
- Actually, using the reduction proving integer programming is NP-hard, this holds for any  $L \in \text{NP}$ .

## Integer programming

Given a language  $L \subseteq \{0,1\}^* \in \text{NP}$ , there is a system  $Ax + By \leq b$  such that

$$\{x \in \{0,1\}^k : x \in L\} = \{x \in \{0,1\}^k : \exists y \in \{0,1\}^m Ax + By \leq b\}$$


where the number of extra variables  $m$  and the number of inequalities is polynomially bounded by the size of the input  $x$ .

## Definition

Given a set  $X \subseteq \mathbb{Z}^d$ , let us call a polyhedron  $R \subseteq \mathbb{R}^d$  a relaxation for  $X$  if  $R \cap \mathbb{Z}^d = \text{conv}(X) \cap \mathbb{Z}^d$  holds.

## Definition


Given a set  $X \subseteq \mathbb{Z}^d$ , let us call a polyhedron  $R \subseteq \mathbb{R}^d$  a relaxation for  $X$  if  $R \cap \mathbb{Z}^d = \text{conv}(X) \cap \mathbb{Z}^d$  holds.



So they have the same set of integer solutions.

## Definition

Given a set  $X \subseteq \mathbb{Z}^d$ , let us call a polyhedron  $R \subseteq \mathbb{R}^d$  a relaxation for  $X$  if  $R \cap \mathbb{Z}^d = \text{conv}(X) \cap \mathbb{Z}^d$  holds.



So they have the same set of integer solutions.

If  $R = \{x \in \mathbb{R}^d \mid Ax \leq b\}$ , then  $\text{conv}(X) = \{x \in \mathbb{Z}^d \mid Ax \leq b\}$ , an ILP problem.

## Definition

The smallest number  $rc(X)$  of facets among any relaxation for  $X$  will be called the relaxation complexity of  $X$ .

## Definition

The smallest number  $rc(X)$  of facets among any relaxation for  $X$  will be called the relaxation complexity of  $X$ .

As we know, we need as many inequalities as facets.

## Definition

The smallest number  $rc(X)$  of facets among any relaxation for  $X$  will be called the relaxation complexity of  $X$ .

As we know, we need as many inequalities as facets.

With this notation, the initial question asks for the asymptotic behavior of  $rc(STSP_n)$ , the traveling salesman problem.

## Previous work

- No reference that deals with a similar quantity except for a paper by Jeroslow.
- For a set  $X \subseteq \{0, 1\}^d$  of binary vectors, Jeroslow introduces the term index of  $X$  (short:  $\text{ind}(X)$ ), defined as the smallest number of inequalities needed to separate  $X$  from the remaining points in  $\{0, 1\}^d$ .
- Thus, relaxation complexity can be seen as a natural extension of the index with respect to general subsets of  $\mathbb{Z}^d$ .



## Previous work

- No reference that deals with a similar quantity except for a paper by Jeroslow.
- For a set  $X \subseteq \{0, 1\}^d$  of binary vectors, Jeroslow introduces the term index of  $X$  (short:  $\text{ind}(X)$ ), defined as the smallest number of inequalities needed to separate  $X$  from the remaining points in  $\{0, 1\}^d$ .
- Thus, relaxation complexity can be seen as a natural extension of the index with respect to general subsets of  $\mathbb{Z}^d$ .

How to find  $\text{RC}(X)$ ?

# Lower bound framework

We introduce a simple framework to achieve that. Assume from now on  $X$  is polyhedral, that is,  $\text{conv}(X)$  is a polyhedron.

## Definition


Let  $X \subseteq \mathbb{Z}^d$ . A set  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus \text{conv}(X)$  is called a hiding set for  $X$  if for any two distinct points  $a, b \in H$  we have that  $\text{conv}\{a, b\} \cap \text{conv}(X) \neq \emptyset$ .

# Lower bound framework


We introduce a simple framework to achieve that. Assume from now on  $X$  is polyhedral, that is,  $\text{conv}(X)$  is a polyhedron.

## Definition

Let  $X \subseteq \mathbb{Z}^d$ . A set  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus \text{conv}(X)$  is called a hiding set for  $X$  if for any two distinct points  $a, b \in H$  we have that  $\text{conv}\{a, b\} \cap \text{conv}(X) \neq \emptyset$ .




# Lower bound framework



Thinking of  $\text{conv}(X)$  as an obstacle, every pair  $a, b \in H$  is "hidden" from each other.

# Lower bound framework




Thinking of  $\text{conv}(X)$  as an obstacle, every pair  $a, b \in H$  is "hidden" from each other.

## Proposition

Let  $X \subseteq \mathbb{Z}^d$  be polyhedral and  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus X$  a hiding set for  $X$ . Then,  $rc(X) \geq |H|$ .

# Lower bound framework



Thinking of  $\text{conv}(X)$  as an obstacle, every pair  $a, b \in H$  is "hidden" from each other.

## Proposition

Let  $X \subseteq \mathbb{Z}^d$  be polyhedral and  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus X$  a hiding set for  $X$ . Then,  $rc(X) \geq |H|$ .

## Proposition

Let  $X \subseteq \mathbb{Z}^d$  be polyhedral and  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus X$  a hiding set for  $X$ . Then,  $rc(X) \geq |H|$ .

## Proposition

Let  $X \subseteq \mathbb{Z}^d$  be polyhedral and  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus X$  a hiding set for  $X$ . Then,  $rc(X) \geq |H|$ .

## Proof.

Let  $R$  be a relaxation of  $X$ . We will prove that if an inequality  $\langle a, x \rangle \leq \beta$  of  $R$  removes more than 1 points of  $H$  from the set of feasible solutions, it must also remove a point of  $R$ , a contradiction.



## Proposition

Let  $X \subseteq \mathbb{Z}^d$  be polyhedral and  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus X$  a hiding set for  $X$ . Then,  $rc(X) \geq |H|$ .

## Proof.

Let  $R$  be a relaxation of  $X$ . We will prove that if an inequality  $\langle a, x \rangle \leq \beta$  of  $R$  removes more than 1 points of  $H$  from the set of feasible solutions, it must also remove a point of  $R$ , a contradiction. As such, to remove all points of  $H$ , we need  $|H|$  inequalities at least.

## Proposition

Let  $X \subseteq \mathbb{Z}^d$  be polyhedral and  $H \subseteq \text{aff}(X) \cap \mathbb{Z}^d \setminus X$  a hiding set for  $X$ . Then,  $rc(X) \geq |H|$ .

## Proof.

Let  $R$  be a relaxation of  $X$ . We will prove that if an inequality  $\langle a, x \rangle \leq \beta$  of  $R$  removes more than 1 points of  $H$  from the set of feasible solutions, it must also remove a point of  $R$ , a contradiction. As such, to remove all points of  $H$ , we need  $|H|$  inequalities at least.

Suppose  $h_1, h_2 \in H$  violate  $\langle a, x \rangle \leq \beta$ . Then, all of  $\text{conv}(h_1, h_2)$  does as well. But  $\text{conv}\{a, b\} \cap \text{conv}(X) \neq \emptyset$ , and  $\text{conv}(X) \subseteq R$ . □


- We are ready to prove that the subtour relaxation for  $STSP_n$  has asymptotically smallest size, i.e.,  $rc(STSP_n) = 2^{\Theta(n)}$ .

- We are ready to prove that the subtour relaxation for  $STSP_n$  has asymptotically smallest size, i.e.,  $rc(STSP_n) = 2^{\Theta(n)}$ .
- Assume number of nodes is even,  $n = 2(N + 1)$ ,  $N \geq 0$ .


- We are ready to prove that the subtour relaxation for  $STSP_n$  has asymptotically smallest size, i.e.,  $rc(STSP_n) = 2^{\Theta(n)}$ .
- Assume number of nodes is even,  $n = 2(N + 1)$ ,  $N \geq 0$ .
- Let node set be  $V := \{v_1, \dots, v_{N+1}, w_1, \dots, w_{N+1}\}$

- We are ready to prove that the subtour relaxation for  $STSP_n$  has asymptotically smallest size, i.e.,  $rc(STSP_n) = 2^{\Theta(n)}$ .
- Assume number of nodes is even,  $n = 2(N + 1)$ ,  $N \geq 0$ .
- Let node set be  $V := \{v_1, \dots, v_{N+1}, w_1, \dots, w_{N+1}\}$
- In fact, we first prove this for the directed version, ATSP. We have an arc set  $A$  instead of edge set  $E$ .

Let us keep only some edges to define the following subgraph (example for  $N=6$ ).




Let us keep only some edges to define the following subgraph (example for  $N=6$ ).




Not the only graph we care about, so to make notation easier, rather than  $\{(v_6, v_1), (w_6, w_1)\} \cup \{(v_1, v_2), (w_1, w_2)\} \cup \{(v_2, v_3), (w_2, w_3)\} \cup \{(v_3, v_4), (w_3, w_4)\} \cup \{(v_4, v_5), (w_4, w_5)\} \cup \{(v_5, w_6), (w_5, v_6)\}$  represent this graph with "01001".






Other variations possible.



Other variations possible.



## Definition

Let there be a binary vector  $b \in \{0, 1\}^N$ . Let us define an arc set as a function of  $b$ :  $E_b := \{(v_{N+1}, v_1), (w_{N+1}, w_1)\} \cup \bigcup_{i:b_i=0} \{(v_i, v_{i+1}), (w_i, w_{i+1})\} \cup \bigcup_{i:b_i=1} \{(v_i, w_{i+1}), (w_i, v_{i+1})\}$

## Definition

Let there be a binary vector  $b \in \{0, 1\}^N$ . Let us define an arc set as a function of  $b$ :  $E_b := \{(v_{N+1}, v_1), (w_{N+1}, w_1)\} \cup \bigcup_{i:b_i=0} \{(v_i, v_{i+1}), (w_i, w_{i+1})\} \cup \bigcup_{i:b_i=1} \{(v_i, w_{i+1}), (w_i, v_{i+1})\}$

## Hiding set

Note  $E_b$  is a directed Hamiltonian cycle on node set  $V$  if and only if  $\sum_{i=1}^N b_i$  is odd.

## Definition

Let there be a binary vector  $b \in \{0, 1\}^N$ . Let us define an arc set as a function of  $b$ :  $E_b := \{(v_{N+1}, v_1), (w_{N+1}, w_1)\} \cup \bigcup_{i:b_i=0} \{(v_i, v_{i+1}), (w_i, w_{i+1})\} \cup \bigcup_{i:b_i=1} \{(v_i, w_{i+1}), (w_i, v_{i+1})\}$

## Hiding set

Note  $E_b$  is a directed Hamiltonian cycle on node set  $V$  if and only if  $\sum_{i=1}^N b_i$  is odd. Define our hiding set as the other  $E_b$ s.  
 $H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$

# STSP and ASTP

$H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$  is a hiding set.

# STSP and ASTP

$H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$  is a hiding set.

$H_N \subseteq \text{aff}(ATSP_{2(N+1)}) \cap \mathbb{Z}^A \setminus \text{conv}(ATSP_{2(N+1)})$ :

# STSP and ASTP

$H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$  is a hiding set.

$H_N \subseteq \text{aff}(ATSP_{2(N+1)}) \cap \mathbb{Z}^A \setminus \text{conv}(ATSP_{2(N+1)})$ :

- $H_N$  only has values in  $\mathbb{Z}^A$



# STSP and ASTP

$H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$  is a hiding set.

$H_N \subseteq \text{aff}(ATSP_{2(N+1)}) \cap \mathbb{Z}^A \setminus \text{conv}(ATSP_{2(N+1)})$ :

- $H_N$  only has values in  $\mathbb{Z}^A$
- $\chi(\delta^{\text{out}}(S)) > 1$  for all  $S$  for any element of  $\text{conv}(X)$ , but not for all  $S$  when it comes to elements of  $H_N$ .

# STSP and ASTP

$H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$  is a hiding set.

$H_N \subseteq \text{aff}(ATSP_{2(N+1)}) \cap \mathbb{Z}^A \setminus \text{conv}(ATSP_{2(N+1)})$ :

- $H_N$  only has values in  $\mathbb{Z}^A$
- $\chi(\delta^{out}(S)) > 1$  for all  $S$  for any element of  $\text{conv}(X)$ , but not for all  $S$  when it comes to elements of  $H_N$ .
- $H_N \subseteq \text{aff}(ATSP_{2(N+1)}) = \{x \in \mathbb{R}^A : x(\delta^{in}(v)) = x(\delta^{out}(v)) = 1, \forall v \in V\}$



# STSP and ASTP

$H_N := \{\chi(E_b) : b \in \{0, 1\}^N, \sum_{i=1}^N b_i \text{ is even}\}$  is a hiding set.

$H_N \subseteq \text{aff}(\text{ATSP}_{2(N+1)}) \cap \mathbb{Z}^A \setminus \text{conv}(\text{ATSP}_{2(N+1)})$ :

- $H_N$  only has values in  $\mathbb{Z}^A$
- $\chi(\delta^{\text{out}}(S)) > 1$  for all  $S$  for any element of  $\text{conv}(X)$ , but not for all  $S$  when it comes to elements of  $H_N$ .
- $H_N \subseteq \text{aff}(\text{ATSP}_{2(N+1)}) = \{x \in \mathbb{R}^A : x(\delta^{\text{in}}(v)) = x(\delta^{\text{out}}(v)) = 1, \forall v \in V\}$



$\text{conv}(\chi(E_b), \chi(E_{b'})) \cap \text{conv}(\text{ATSP}_{2(N+1)}) \neq \emptyset$  :

Let  $b, b' \in \{0, 1\}^N$  be distinct with  $b$  and  $b'$  having an even number of ones. Let  $j$  be an index with  $b_j \neq b'_j$ . Flip  $b_j, b'_j$  to get  $c, c'$ .  $c, c'$  have an odd number of ones, hence  $\chi(E_c)$  and  $\chi(E_{c'})$  are both contained in  $\text{ATSP}_{2(N+1)}$ . Clearly  $\chi(E_b) + \chi(E_{b'}) = \chi(E_c) + \chi(E_{c'})$ .



The asymptotic growth of  $\text{rc}(\text{ATSP}_n)$  and  $\text{rc}(\text{STSP}_n)$  is  $2^{\Theta(n)}$ .

## Proof.

Clearly,  $|H_N| = 2^{\Theta(n)}$ . Note both ATSP and ASTP have formulations of size  $2^{O(n)}$ .  $\text{rc}(\text{ATSP}_n) = 2^{\Theta(n)}$  follows instantly.

For  $\text{STSP}_n$ , replace all directed arcs with undirected edges,  $H_N$  is still a hiding set. □

- Let  $CONN_n$  be the set of all characteristic vectors of edge sets that form a connected spanning subgraph in the complete graph on  $n$  nodes.

- Let  $CONN_n$  be the set of all characteristic vectors of edge sets that form a connected spanning subgraph in the complete graph on  $n$  nodes.
- The polytope  $\{x \in [0, 1]^{E_n} : x(\delta(S)) \geq 1, \forall \emptyset \neq S \subset V_n\}$  is a relaxation for  $CONN_n$ .

- Let  $CONN_n$  be the set of all characteristic vectors of edge sets that form a connected spanning subgraph in the complete graph on  $n$  nodes.
- The polytope  $\{x \in [0, 1]^{E_n} : x(\delta(S)) \geq 1, \forall \emptyset \neq S \subset V_n\}$  is a relaxation for  $CONN_n$ .
- Thus, we have that  $rc(CONN_n) \leq O(2^n)$ .

- Let  $CONN_n$  be the set of all characteristic vectors of edge sets that form a connected spanning subgraph in the complete graph on  $n$  nodes.
- The polytope  $\{x \in [0, 1]^{E_n} : x(\delta(S)) \geq 1, \forall \emptyset \neq S \subset V_n\}$  is a relaxation for  $CONN_n$ .
- Thus, we have that  $rc(CONN_n) \leq O(2^n)$ .
- We prove the asymptotic growth of  $rc(CONN_n)$  is  $2^{\Theta(n)}$ .



- Let  $CONN_n$  be the set of all characteristic vectors of edge sets that form a connected spanning subgraph in the complete graph on  $n$  nodes.
- The polytope  $\{x \in [0, 1]^{E_n} : x(\delta(S)) \geq 1, \forall \emptyset \neq S \subset V_n\}$  is a relaxation for  $CONN_n$ .
- Thus, we have that  $rc(CONN_n) \leq O(2^n)$ .
- We prove the asymptotic growth of  $rc(CONN_n)$  is  $2^{\Theta(n)}$ .
- So, something as simple as encoding connectivity on a graph requires an exponential in  $n$  number of inequalities.

- Let  $CONN_n$  be the set of all characteristic vectors of edge sets that form a connected spanning subgraph in the complete graph on  $n$  nodes.
  - The polytope  $\{x \in [0, 1]^{E_n} : x(\delta(S)) \geq 1, \forall \emptyset \neq S \subset V_n\}$  is a relaxation for  $CONN_n$ .
  - Thus, we have that  $rc(CONN_n) \leq O(2^n)$ .
  - We prove the asymptotic growth of  $rc(CONN_n)$  is  $2^{\Theta(n)}$ .
  - So, something as simple as encoding connectivity on a graph requires an exponential in  $n$  number of inequalities.
- $H_N$  from before for undirected graphs is still a hiding set:

Remember, the polytope  $\{x \in [0, 1]^{E_n} : x(\delta(S)) \geq 1, \forall \emptyset \neq S \subset V_n\}$  is a relaxation for  $\text{CONN}_n$ .

Proof.

$H_N \subseteq \text{aff}(\text{CONN}_n) \cap \mathbb{Z}^d \setminus \text{conv}(\text{CONN}_n)$ :

- $H_N$  only has values in  $\mathbb{Z}^{E_n}$
- $\chi(\delta(S)) \geq 1$  for all  $S$  for any element of  $\text{conv}(X)$ , and not for all  $S$  for elements of  $H_N$ .
- It holds that  $\text{aff}(\text{CONN}_n) = \mathbb{R}^{E_n}$ .

Let  $a, b \in H_N$ .

$\emptyset \neq \text{conv}\{a, b\} \cap \text{conv}(\text{STSP}_n) \subseteq \text{conv}\{a, b\} \cap \text{conv}(\text{CONN}_n)$ . □


# Spanning trees and arborescences

- Connectivity on a graph requires an exponential in  $n$  number of inequalities.
- Perhaps something like, such as spanning trees or an arborescence?
- Formulations of  $2^{\mathcal{O}(n)}$  known for both.


# Spanning trees and arborescences

- Connectivity on a graph requires an exponential in  $n$  number of inequalities.
- Perhaps something like, such as spanning trees or an arborescence?
- Formulations of  $2^{\mathcal{O}(n)}$  known for both.

Spanning tree



Arborescence



# Spanning trees and arborescences


- Connectivity on a graph requires an exponential in  $n$  number of inequalities.
- Perhaps something like, such as spanning trees or an arborescence?
- Formulations of  $2^{\mathcal{O}(n)}$  known for both.

## Towards a lower bound

Let us modify the definition of  $E_b$  by removing arc  $(w_{N+1}, w_1)$ . Then, if  $b \in \{0, 1\}^N$  with  $b$  having an even number of ones, we have that  $E_b$  is a node-disjoint union of a cycle and a path and hence not an arborescence.

We will obtain that the modified set  $H_N$  is a hiding set for  $\text{ARB}_n$ .

# Spanning trees and arborescences



## Proof.

$$H_N \subseteq \text{aff}(\text{ARB}_N) \cap \mathbb{Z}^d \setminus \text{conv}(\text{ARB}_N):$$

- $H_N$  only has values in  $\mathbb{Z}^{E_n}$
- $\chi(\delta^{\text{out}}(S)) \geq 1$  for all  $S$  for any element of  $\text{conv}(X)$ , and not for all  $S$  of  $H_N$ .
- It holds that  $\text{aff}(\text{ARB}_n) = \mathbb{R}^{A_n}$ .



# Spanning trees and arborescences

## Proof.

$H_N \subseteq \text{aff}(\text{ARB}_N) \cap \mathbb{Z}^d \setminus \text{conv}(\text{ARB}_N)$ :

- $H_N$  only has values in  $\mathbb{Z}^{E_n}$
- $\chi(\delta^{\text{out}}(S)) \geq 1$  for all  $S$  for any element of  $\text{conv}(X)$ , and not for all  $S$  of  $H_N$ .
- It holds that  $\text{aff}(\text{ARB}_n) = \mathbb{R}^{A_n}$ .

$\text{conv}(\text{ARB}_N) \cap \text{conv}(\chi(E_b), \chi(E_{b'})) \neq \emptyset$ :

- Choosing  $b, b'$  with even number of ones, and flipping a bit they differ on, we still have  $\chi(E_b) + \chi(E_{b'}) = \chi(E_c) + \chi(E_{c'})$ , where  $E_c$  and  $E_{c'}$  are hamilton paths, and thus arborescences, therefore  $\text{conv}(\text{ARB}_N) \cap \text{conv}(\chi(E_b), \chi(E_{b'})) \neq \emptyset$ .



# Spanning trees and arborescences

## Proof.

$H_N \subseteq \text{aff}(\text{ARB}_N) \cap \mathbb{Z}^d \setminus \text{conv}(\text{ARB}_N)$ :

- $H_N$  only has values in  $\mathbb{Z}^{E_n}$
- $\chi(\delta^{\text{out}}(S)) \geq 1$  for all  $S$  for any element of  $\text{conv}(X)$ , and not for all  $S$  of  $H_N$ .
- It holds that  $\text{aff}(\text{ARB}_n) = \mathbb{R}^{A_n}$ .

$\text{conv}(\text{ARB}_N) \cap \text{conv}(\chi(E_b), \chi(E_{b'})) \neq \emptyset$ :

- Choosing  $b, b'$  with even number of ones, and flipping a bit they differ on, we still have  $\chi(E_b) + \chi(E_{b'}) = \chi(E_c) + \chi(E_{c'})$ , where  $E_c$  and  $E_{c'}$  are hamilton paths, and thus arborescences, therefore  $\text{conv}(\text{ARB}_N) \cap \text{conv}(\chi(E_b), \chi(E_{b'})) \neq \emptyset$ .



By undirecting all arcs,  $H_N$  also yields a hiding set for  $\text{SPT}_n$ . We deduce a lower bound of  $|H_N| = 2^{\Omega(n)}$  for both  $\text{rc}(\text{ARB}_n)$  and  $\text{rc}(\text{SPT}_n)$ .

# Branchings and forests

Let  $\text{BRANCH}_n$  ( $\text{FORESTS}_n$ ) be the set of characteristic vectors of branchings (forests) in the complete directed (undirected) graph.

# Branchings and forests

Let  $\text{BRANCH}_n$  ( $\text{FORESTS}_n$ ) be the set of characteristic vectors of branchings (forests) in the complete directed (undirected) graph.

## Notice

$$\text{SPT}_n = \text{FORESTS}_n \cap \{x \in \mathbb{R}^{E_n} : \sum_{e \in \mathbb{R}^{E_n}} x_e = n - 1\}.$$

# Branchings and forests

Let  $\text{BRANCH}_n$  ( $\text{FORESTS}_n$ ) be the set of characteristic vectors of branchings (forests) in the complete directed (undirected) graph.

## Notice

$\text{SPT}_n = \text{FORESTS}_n \cap \{x \in \mathbb{R}^{E_n} : \sum_{e \in \mathbb{R}^{E_n}} x_e = n - 1\}$ .

So, if we could encode  $\text{FORESTS}_n$  in  $2^{\mathcal{O}(n)}$  inequalities, we could do the same with  $\text{SPT}_n$ .

# Branchings and forests

Let  $\text{BRANCH}_n$  ( $\text{FORESTS}_n$ ) be the set of characteristic vectors of branchings (forests) in the complete directed (undirected) graph.

## Notice

$$\text{SPT}_n = \text{FORESTS}_n \cap \{x \in \mathbb{R}^{E_n} : \sum_{e \in \mathbb{R}^{E_n}} x_e = n - 1\}.$$

So, if we could encode  $\text{FORESTS}_n$  in  $2^{O(n)}$  inequalities, we could do the same with  $\text{SPT}_n$ . Similarly,

$$\text{ARB}_n = \text{BRANCH}_n \cap \{x \in \mathbb{R}^{A_n} : \sum_{e \in \mathbb{R}^{E_n}} x_e = n - 1\}.$$

# Binary-all-different

How about encoding that an  $m \times n$  0-1 matrix has no double row?

# Binary-all-different

How about encoding that an  $m \times n$  0-1 matrix has no double row?

$$\text{DIFF}_{m,n} := \{x \in \{0,1\}^{m \times n} : x \text{ has pairwise distinct rows}\}$$



# Binary-all-different

How about encoding that an  $m \times n$  0-1 matrix has no double row?

$\text{DIFF}_{m,n} := \{x \in \{0,1\}^{m \times n} : x \text{ has pairwise distinct rows}\}$

Can be done in  $\binom{m}{2}2^n + 2mn$  inequalities.

# Binary-all-different

How about encoding that an  $m \times n$  0-1 matrix has no double row?

$\text{DIFF}_{m,n} := \{x \in \{0,1\}^{m \times n} : x \text{ has pairwise distinct rows}\}$

Can be done in  $\binom{m}{2}2^n + 2mn$  inequalities. Almost tight, even for 2 rows.

## Hiding set

$$H_{2,n} := \{(x, x)^T \in \{0,1\}^{2 \times n} : x \in \{0,1\}^n\}$$

# Binary-all-different

How about encoding that an  $m \times n$  0-1 matrix has no double row?

$\text{DIFF}_{m,n} := \{x \in \{0,1\}^{m \times n} : x \text{ has pairwise distinct rows}\}$

Can be done in  $\binom{m}{2}2^n + 2mn$  inequalities. Almost tight, even for 2 rows.

## Hiding set

$$H_{2,n} := \{(x, x)^T \in \{0,1\}^{2 \times n} : x \in \{0,1\}^n\}$$

## Proof.

$$H_{2,n} \subseteq \text{aff}(\text{DIFF}_{2 \times n}) \cap \mathbb{Z}^d \setminus \text{conv}(\text{DIFF}_{m,n}):$$

- $H_{2,n}$  only has values in  $\mathbb{Z}^{2 \times n}$
- $\text{conv}(\text{DIFF}_{m,n})$  has no common element with  $H_{2,n}$ , clearly.
- It holds that  $\text{aff}(\text{DIFF}_{2 \times n}) = \mathbb{R}^{2 \times n}$ .

Let  $(x, x)^T, (y, y)^T \in H_{2,n}$ .

$$\frac{(x, x)^T + (y, y)^T}{2} = \frac{(x, y)^T + (y, x)^T}{2} \in \text{CONV}(\text{DIFF}_{2,n}).$$



# Permutahedron

$PERM_n = (\pi(1), \dots, \pi(n)) \in \mathbb{Z}^n : \pi \text{ is a permutation of } [n].$

# Permutahedron

$PERM_n = (\pi(1), \dots, \pi(n)) \in \mathbb{Z}^n : \pi \text{ is a permutation of } [n]$ .

It's proven that

$$\begin{aligned} \text{conv}(Perm_n) = \{x \in \mathbb{R}^n : & \sum_{i=1}^n x_i = n(n+1)/2 \\ & \sum_{i \in S} x_i \geq |S|(|S|+1)/2 \text{ for all } \emptyset \neq S \subset [n] \\ & x \geq 0\} \end{aligned}$$

# Permutahedron

$PERM_n = (\pi(1), \dots, \pi(n)) \in \mathbb{Z}^n : \pi \text{ is a permutation of } [n]$ .

It's proven that

$$\begin{aligned} \text{conv}(Perm_n) = \{x \in \mathbb{R}^n : & \sum_{i=1}^n x_i = n(n+1)/2 \\ & \sum_{i \in S} x_i \geq |S|(|S|+1)/2 \text{ for all } \emptyset \neq S \subset [n] \\ & x \geq 0\} \end{aligned}$$

The asymptotic growth of  $\text{rc}(PERM_n)$  is  $2^{\Theta(n)}$

## Preliminary

Set  $S := \{i \in [n] : i \text{ is odd}\}$ .  $|S| = \lfloor \frac{n}{2} \rfloor$ . Select an integer vector  $x \in \mathbb{Z}^n$  with  $\{x_i : i \text{ is odd} (= S)\} = \{1, \dots, \lfloor \frac{n}{2} \rfloor - 1\}$  and  $\lfloor \frac{n}{2} \rfloor - 1$  occurring twice, and  $\{x_i : i \notin S\} = \{\lfloor \frac{n}{2} \rfloor + 2, \dots, n\}$  and  $\lfloor \frac{n}{2} \rfloor + 2$  occurring twice. Such a vector is not contained in  $\text{conv}(PERM_n)$ .

$$\begin{aligned} \text{conv}(\text{Perm}_n) = \{x \in \mathbb{R}^n : & \sum_{i=1}^n x_i = n(n+1)/2 \\ & \sum_{i \in S} x_i \geq |S|(|S|+1)/2 \text{ for all } \emptyset \neq S \subset [n] \\ & x \geq 0\} \end{aligned}$$

## Preliminary

$$\sum_{i \in S} x_i = 1 + 2 + \dots + (|S| - 1) + (|S| - 1) < |S|(|S| + 1)/2$$

### Note

- The only violated inequality is this one.
- $x \in \text{aff}(\text{Perm}_n)$  as it satisfies all implied equalities.
- We used only the cardinality of  $S$  so this holds for any  $S$  so that  $|S| = \lfloor \frac{n}{2} \rfloor$

$H := \{x^S : S \subseteq [n], |S| = \lfloor \frac{n}{2} \rfloor - 1\}$  is a hiding set for  $\text{PERM}_n$ , where  $x^S$  is defined as follows: Select an integer vector  $x \in \mathbb{Z}^n$  with
   
 $\{x_i : i \in S\} = \{1, \dots, \lfloor \frac{n}{2} \rfloor - 1\}$  and  $\lfloor \frac{n}{2} \rfloor - 1$  occurring twice, and
   
 $\{x_i : i \notin S\} = \{\lfloor \frac{n}{2} \rfloor + 2, \dots, n\}$  and  $\lfloor \frac{n}{2} \rfloor + 2$  occurring twice.



$H := \{x^S : S \subseteq [n], |S| = \lfloor \frac{n}{2} \rfloor - 1\}$  is a hiding set for  $\text{PERM}_n$ , where  $x^S$  is defined as follows: Select an integer vector  $x \in \mathbb{Z}^n$  with  $\{x_i : i \in S\} = \{1, \dots, \lfloor \frac{n}{2} \rfloor - 1\}$  and  $\lfloor \frac{n}{2} \rfloor - 1$  occurring twice, and  $\{x_i : i \notin S\} = \{\lfloor \frac{n}{2} \rfloor + 2, \dots, n\}$  and  $\lfloor \frac{n}{2} \rfloor + 2$  occurring twice.

## Proof.

$H \subseteq \text{aff}(\text{PERM}_n) \cap \mathbb{Z}^n \setminus \text{conv}(\text{PERM}_n)$ :

- $H$  only has values in  $\mathbb{Z}^n$ .
- $\text{conv}(\text{PERM}_n)$  has no common element with  $H$ , as demonstrated.
- $H \subseteq \text{aff}(\text{PERM}_n)$ , as demonstrated.

Let  $S_1, S_2 \in H$ . We will show that  $x := \frac{1}{2} \cdot (x^{S_1} + x^{S_2}) \in \text{conv}(\text{PERM}_n)$  holds. Since  $x$  satisfies all constraints that are satisfied by both  $x^{S_1}$  and  $x^{S_2}$ , it suffices to show that  $\sum_{i \in S_1} x_i \geq |S_1|(|S_1| + 1)/2$ ,  $\sum_{i \in S_1} x_i \geq |S_2|(|S_2| + 1)/2$  holds.

## Proof.

Let  $S_1, S_2 \in H$ . We will show that  $x := \frac{1}{2} \cdot (x^{S_1} + x^{S_2}) \in \text{conv}(\text{PERM}_n)$  holds. Since  $x$  satisfies all constraints that are satisfied by both  $x^{S_1}$  and  $x^{S_2}$ , it suffices to show that  $\sum_{i \in S_1} x_i \geq |S_1|(|S_1| + 1)/2$ ,  $\sum_{i \in S_2} x_i \geq |S_2|(|S_2| + 1)/2$  holds.

$$\begin{aligned} \sum_{i \in S_1} x_i &= \frac{1}{2} \sum_{i \in S_1} x_i^{S_1} + \frac{1}{2} \sum_{i \in S_1} x_i^{S_2} \\ &= \frac{1}{2} \left( \frac{m(m+1)}{2} - 1 \right) + \frac{1}{2} \frac{m(m+1)}{2} + 2 \\ &\geq \frac{m(m+1)}{2} \end{aligned}$$



## Rational relaxations

- What happens if we restrict ourselves to rational values in our description?
- We don't lose too much.

## Rational relaxations

- What happens if we restrict ourselves to rational values in our description?
- We don't lose too much.

## Theorem

Let  $X \subseteq \mathbb{Z}^d$  be finite and  $rc_Q(X)$  be the smallest number of facets of any rational relaxation for  $X$ . Then,  $rc_Q(X) \leq rc(X) + \dim(X) + 1$ .

If  $X \subseteq \{0, 1\}^d$  only, the coefficients of this rational relaxation  $R$  can be polynomially bounded in size by  $d$ .

## Questions

- Is it true that  $rc(X) \geq \dim(X) + 1$  holds for all polyhedral (or at least finite) sets  $X \subseteq \mathbb{R}^d$ ? (holds for  $X \subset \{0, 1\}^d$ ).
- Is there any polyhedral (or even finite) set  $X \subset \mathbb{Z}^d$  such that  $rc(X) < rc_Q(X)$ ?

## Weltge PHD thesis:

- In dimension two,  $rc(X)$  and  $rc_Q(X)$  coincide and are computable.
- $\dim(X) \geq k! \implies rc(X) \geq k$

## On the Size of Integer Programs with Bounded Coefficients or Sparse Constraints (2018)

- A simple algorithm to compute the maximum hiding set for  $X \subset S$  where  $|S| < |\mathbb{N}|$  is given.
- Given a nonempty set  $X \subset \{0, 1\}^n$ , build a graph  $G = (V, E)$  with  $V = \{0, 1\}^n$ . The edge set  $E$  of  $G$  is defined as  $E = \{\{x, y\} : x, y \in V, \text{conv}(x, y) \cap \text{conv}(X) \neq \emptyset\}$ .
- Any clique in  $G$  is a hiding set by definition. By computing the size of a maximum clique in  $G$ , we get the maximum size of any hiding set  $H \subseteq \{0, 1\}^n$  for  $X$ .

## Complexity of linear relaxations in integer programming (2020)

Main results: ( $X$  assumed a finite lattice-convex set)

- ①  $\text{rc}(X) = \text{rc}_Q(X)$  when:
  - ①  $X$  is at most four-dimensional,
  - ②  $X$  represents every residue class in  $(\mathbb{Z}/2\mathbb{Z})^d$
  - ③ the convex hull of  $X$  contains an interior integer point,
  - ④ the lattice-width of  $X$  is above a certain threshold.
- ②  $\text{rc}(X)$  can be algorithmically computed when
  - ①  $X$  is at most three-dimensional, or
  - ②  $X$  satisfies one of the last three conditions above.
- ③ An improved lower bound on  $\text{rc}(X)$  in terms of the dimension of  $X$  is obtained.  $\text{rc}(X) > \log(\dim(X)) - \log\log(\dim(X))$ .



## Strong IP formulations need large coefficients (2021)

Generalization of hiding sets.

- A set  $H \subseteq (\text{aff}(X) \cap \mathbb{Z}^n)$  is called a hiding set for  $X$  if  $\text{conv}\{x_1, x_2\} \cap \text{conv}(X) \neq \emptyset$  for each pair of distinct points  $x_1, x_2 \in H$ .
- A set  $H \subseteq (\text{aff}(X) \cap F)$  is called an  $F$ -hiding set for  $X$  if  $\text{conv}\{x_1, x_2\} \cap \text{conv}(X) \neq \emptyset$  for each pair of distinct points  $x_1, x_2 \in H$ , where  $F \subseteq \mathbb{R}^n$ .
- Let  $X \subseteq \mathbb{Z}^n$  be polyhedral, let  $F \subseteq \mathbb{R}^n$ , and let  $H$  be an  $F$ -hiding set for  $X$ . Then, every family of inequalities separating  $F \cap \text{conv}(X)$  and  $F \setminus \text{conv}(X)$  has size at least  $|H|$ .


## Computational Aspects of Relaxation Complexity: Possibilities and Limitations (2021)

- Tight and computable upper bounds on  $rc_Q(X)$ .
- $rc(X)$  can be computed in polynomial time if  $X$  is 2-dimensional.
- Explicit formula for  $rc(X)$  for specific classes of sets  $X$ .
- First practically applicable approach to compute  $rc(X)$  for specific classes of sets  $X$ .

## Efficient MIP Techniques for Computing the Relaxation Complexity (2022)

- Techniques to efficiently compute a numerically more robust variant of the relaxation complexity,  $rc_\varepsilon(X)$ , where  $\text{conv}(X) \cap \mathbb{Z}^d = \{x \in \mathbb{Z}^d \mid Ax \leq b + \varepsilon\}$  instead of  $\{x \in \mathbb{Z}^d \mid Ax \leq b\}$ .

# Ευχαριστούμε!



- ① R. D. Carr, L. Fleischer, V. J. Leung, and C. A. Phillips, Strengthening integrality gaps for capacitated network design and covering problems, in Proceedings of the ACM-SIAM Symposium on Discrete Algorithms, 2000, pp. 106–115.
- ② N. Bansal and K. Pruhs, Weighted geometric set multi-cover via quasi-uniform sampling, in Proceedings of the European Symposium on Algorithms, 2012, pp. 145–156.