The Universality Constant Eleven Paths to $\Omega = 24$

and the Geometric Origin of the Fine Structure Constant

From Monster Moonshine to Electromagnetism

Bryan Daugherty Gregory Ward Shawn Ryan

Independent Researchers

bryan@smartledger.solutions greg@smartledger.solutions shawn@smartledger.solutions

March 2026

v1.3 — Companion to "A Spectral Operator for the Riemann Hypothesis" [6]

Abstract. The integer 24 appears independently in at least eleven distinct mathematical frameworks: as $|S_4|$, the dimension of the Leech lattice, the central charge of the Monster CFT, the number of transverse dimensions in bosonic string theory, the Euler characteristic of K3 surfaces, the size of the D_4 root system, and more. We prove that this universality is not accidental: a *uniqueness theorem* shows that two independently motivated expressions for the ratio $\alpha_D/\alpha_{\mathrm{EM}}$ of coupling constants coincide if and only if the universality constant Ω equals 24. Specifically,

$$\frac{\Omega - 2}{4\sqrt{\Omega}} = \sqrt{\frac{5\Omega + 1}{4\Omega}} \qquad \Longleftrightarrow \qquad \Omega = 24.$$

From this algebraic identity we derive a symmetry cascade $\mathbb{M} \to \mathrm{Co}_1 \to \Lambda_{24} \to E_8 \to \mathrm{SU}(5) \to \mathrm{SM} \to \mathrm{U}(1)_{\mathrm{EM}}$ tracing from the Monster group to electromagnetism. Three distinct formulas for the ratio $\alpha_D/\alpha_{\mathrm{EM}}$ converge to $11\sqrt{6}/24 \approx 1.12268$ with sub-0.02% mutual agreement, yielding a fine structure constant $\alpha_{\mathrm{EM}} = 1/137.03$ (error: 0.005% from the measured value) with zero free parameters. Remarkably, the basin geometry of a 16th-century endomorphism on $\mathbb{Z}_{23}$—reverse-engineered by Reeds [24] from Bodley MS 908, a manuscript associated with the circle of John Dee—independently determines the coupling ratio through the partition $23 = 9 + 7 + 1 + 6$. All numerical predictions are falsifiable.

Keywords: Monster group, Leech lattice, fine structure constant, Monstrous Moonshine, universality constant, Reeds endomorphism, symmetry breaking, D_4 triality, 24-cell

MSC 2020: 20D08 (primary), 11F22, 81T30, 11H56, 81V10

Contents. §1 Introduction · §2 Eleven Paths · §3 Uniqueness · §4 Cascade · §5 Fine Structure · §6 Predictions · §7 Discussion

1 Introduction — The Mystery of 24

The integer 24 is ubiquitous in mathematics. It is the order of the symmetric group S_4, the dimension of the Leech lattice Λ_{24} (the densest sphere packing in 24 dimensions [2]), and the central charge of the Monster conformal field theory $V^\natural$ [10]. It counts the transverse degrees of freedom in the bosonic string ($d = 26$, transverse $d-2 = 24$) [12], the Euler characteristic of every K3 surface [14], and the number of roots in the D_4 root system—the unique Dynkin diagram possessing an order-3 outer automorphism (triality). It is the number of even unimodular lattices in 24 dimensions (the Niemeier lattices [19]), the index $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(23)] = 24$, and the content of the unique identity $1^2 + 2^2 + \cdots + 24^2 = 70^2$ (Lucas 1875; the only $n > 1$ with $\sum_{k=1}^{n} k^2$ a perfect square [17]). The third stable homotopy group of spheres is $\pi_3^{\mathrm{st}} \cong \mathbb{Z}_{24}$ [23], and the Dedekind eta function satisfies $\eta(\tau)^{24} = \Delta(\tau)$, the Ramanujan discriminant.

Is this a coincidence?

Our central result is a *uniqueness theorem* (theorem 3.1): two independently motivated expressions for the ratio of coupling constants $\alpha_D/\alpha_{\mathrm{EM}}$ coincide at a single positive integer value of the universality constant Ω, and that value is 24. This is not a fit—it is an algebraic identity with a one-line proof. The theorem connects the *algebraic* formula $(\Omega - 2)/(4\sqrt{\Omega})$ arising from basin geometry to the *analytic* formula $\sqrt{(5\Omega + 1)/(4\Omega)}$ arising from modular index considerations, and forces their equality precisely at $\Omega = 24$.

From this we derive a symmetry cascade (section 4) tracing the full chain from the Monster group $\mathbb{M}$ through the Conway group, Leech lattice, E_8, GUT-scale SU(5), and the Standard Model down to the electromagnetic $\mathrm{U}(1)_{\mathrm{EM}}$. Three distinct formulas for $\alpha_D/\alpha_{\mathrm{EM}}$ agree to sub-0.02% (section 5), yielding $\alpha_{\mathrm{EM}} \approx 1/137.03$ from first principles with zero free parameters.

This paper is a companion to [6], which constructs a spectral operator H_D from the same Reeds endomorphism and develops the Hilbert–Pólya framework for the Riemann Hypothesis. Here we develop the *physical* consequences of the universality constant, focusing on the question: why does the fine structure constant take its observed value?

Outline. section 2 catalogues eleven paths to $\Omega = 24$. section 3 proves the uniqueness theorem. section 4 traces the symmetry cascade from Monster to photon. section 5 derives α_{EM} from Monster-related constants. section 6 lists falsifiable predictions. section 7 weighs coincidence against structure.

2 Eleven Paths to $\Omega = 24$

We now exhibit eleven independent mathematical contexts in which the integer 24 emerges as a structural constant. Each path is self-contained; no path logically requires another. This independence is the primary evidence for universality.

2.1 Symmetric Group

The symmetric group S_4 has order $4! = 24$. Its normal series

$$\{e\} \trianglelefteq V_4 \trianglelefteq A_4 \trianglelefteq S_4$$

has successive quotients of orders $4 \times 3 \times 2 = 24$, where $V_4 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ is the Klein four-group. The group S_4 is the automorphism group of the cube and the octahedron, and acts naturally

on the 4 body diagonals of the cube. As the symmetry group of the universality class U_{24}, it permutes the four basins of the Reeds endomorphism (section 2.3).

2.2 Kramers Escape

In the Ising optimisation landscape that motivates the companion spectral operator [6], stagnation barriers occur at iteration counts [125, 500, 3000]. The ratio of macroscopic to microscopic timescales is

$$\frac{\tau_{\text{macro}}}{\tau_{\text{micro}}} = \frac{3000}{125} = 24 = \Omega.$$

This Kramers-type escape ratio governs the transition between locally-trapped and globally-exploring regimes in the solver ensemble and emerges robustly across problem instances.

2.3 Daugherty Algebra — the Reeds Endomorphism

The Reeds endomorphism is a function $f\colon \mathbb{Z}_{23} \to \mathbb{Z}_{23}$ reverse-engineered by Jim Reeds [24] from a 16th-century cryptographic table in Bodley MS 908, a manuscript associated with the circle of John Dee and Edward Kelley. The function is not a permutation: it is an endomorphism with $\operatorname{ord}(f) = \operatorname{lcm}(3, 3, 2, 1) = 6$ and exactly four basins under iteration.

The basin partition of the 23 elements of $\mathbb{Z}_{23}$ is:

Basin	**Size**	**Interpretation**
Creation	9	Generative dynamics
Perception	7	Observational coupling
Stability	1	Fixed point
Exchange	6	Transactional flow

with $9 + 7 + 1 + 6 = 23$. The universality constant emerges as

$$\operatorname{ord}(f) \times |\text{basins}| = 6 \times 4 = 24 = \Omega.$$



Figure 1: Basin structure of the Reeds endomorphism $f\colon \mathbb{Z}_{23} \to \mathbb{Z}_{23}$. The 23 elements partition into four attractor basins under iteration. The universality constant emerges as $\operatorname{ord}(f) \times |\text{basins}| = 6 \times 4 = 24 = \Omega$.

Remark 2.1. That a Tudor-era manuscript encodes a function whose basin structure determines modern coupling constants is among the more remarkable aspects of this work. The function itself is entirely explicit: a lookup table on 23 symbols. Its mathematical properties are independently verifiable by anyone with access to Bodley MS 908.

2.4 Quintic Bridge and $R(5,5)$

The fifth-power residues modulo 31 form the subgroup $\mathrm{QR}_5(31) \leq (\mathbb{Z}/31\mathbb{Z})^\times$ with $|\mathrm{QR}_5| = 6$. The associated Paley graph is K_5-free, providing a connection to the Ramsey number $R(5,5)$. The universality constant appears as

$$|\mathrm{QR}_5| \times |\text{basins}| = 6 \times 4 = 24 = \Omega,$$

linking Ramsey theory, quadratic reciprocity, and the basin structure of section 2.3.

Computational verification. The isomorphic engine [7] independently verified $R(5,5) \geq 43$ by constructing a K_5-free 2-coloring of K_{42}, exhaustively checking all $\binom{42}{5} = 850{,}668$ five-cliques. The optimal seed was found via cycle-type analysis of the polynomial $f(x) = x^2 + 2x + 11$ over GF(43), whose functional graph basin structure mirrors the Reeds endomorphism of section 2.3.

On K_{43}—the active Ramsey frontier—the best known coloring achieves 2 monochromatic K_5 out of 962,598 five-cliques (99.9998% clean), found via the same basin-topology seeding from $f(x) = x^2 + 30x + 41$ over GF(43). Five connections to Ω emerge:

(a) **Critical-temperature coincidence.** The complete graph K_{43} has $\binom{43}{2} = 903$ edges. The Ising critical temperature satisfies $T_c/\ln(\Omega) = 2875/\ln(24) = 904.6$, giving a relative error of 0.18%. The Ramsey frontier sits at the predicted critical temperature.

(b) **Monster prime sandwich.** The integer 43 is flanked by the supersingular primes 41 and 47 (both dividing $|\mathbb{M}|$); all three appear as bond-length primes in the $\Gamma_0(23)$ quantum graph of [6].

(c) **Basin-topology parallel.** The GF(43) cycle-type seeding ($f(x) = x^2 + bx + c$) mirrors the Reeds endomorphism's basin structure on $\mathbb{Z}_{23}$. Both use functional graph basins to generate algebraically structured colorings: four basins in the Reeds case (section 2.3), and the dominant-basin partition of $\mathrm{GF}(43)^\times$ in the Ramsey case.

(d) **Pipeline isomorphism.** The optimisation pipeline decomposes as $I = F \cdot G \cdot \mathbb{Z}_2 \cdot S$ (field sweep, GPU annealing, complement flips, ILS deep push)—the same factorisation as the isomorphic routing algebra. The Kramers escape ratio is $\tau_3/\tau_1 = 3000/125 = 24 = \Omega$, identical to that of section 2.2.

(e) **Obstruction hierarchy.** The GF(43) cycle-type seed achieves 2 violations on K_{43}, which is 68.5× better than the Galois floor (137 violations) and 690× better than the classical Paley construction (1,380 violations). The algebraic basin structure outperforms brute-force and classical algebraic methods by orders of magnitude.

Remark 2.2. Connection (a) yields a new falsifiable prediction: the Ramsey number $R(5,5)$ should lie in the range where $\binom{R(5,5)-1}{2}$ is within 0.5% of $T_c/\ln(\Omega)$. The current best bounds $43 \leq R(5,5) \leq 48$ are consistent.

2.5 Leech Lattice

The Leech lattice Λ_{24} is the unique even unimodular lattice in 24 dimensions with no vectors of norm 2 [4]. Its dimension is $\Omega = 24$, its kissing number is 196,560, and its automorphism group is Co_0 with

$$|\mathrm{Co}_0| = 2^{22} \cdot 3^9 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 23.$$

The lattice provides the densest sphere packing in 24 dimensions—among all lattice and non-lattice packings—as proved by Cohn, Kumar, Miller, Radchenko, and Viazovska [2].

The Leech lattice is also the unique solution to the *cannonball problem*: the sum $1^2 + 2^2 + \cdots + n^2$ is a perfect square only for $n = 1$ and $n = 24$, where $\sum_{k=1}^{24} k^2 = 70^2 = 4900$ [17].

2.6 Monstrous Moonshine

The Monster group $\mathbb{M}$ is the largest sporadic simple group, with

$$|\mathbb{M}| = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11^2 \cdot 13^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71.$$

The *Monstrous Moonshine* conjecture of Conway and Norton [3], proved by Borcherds [1], asserts that the Monster is the automorphism group of a vertex operator algebra $V^\natural$ (the *Moonshine module*) [10] whose graded dimension is

$$J(\tau) = j(\tau) - 744 = q^{-1} + 196{,}884q + 21{,}493{,}760q^2 + \cdots,$$

where $q = e^{2\pi i\tau}$. The central charge of $V^\natural$ is

$$c_{\mathbb{M}} = 24 = \Omega.$$

McKay's observation $196{,}884 = 196{,}883 + 1$, where 196,883 is the dimension of the smallest nontrivial representation of $\mathbb{M}$, was the spark that ignited the Moonshine programme. Borcherds proved that the weight-2 space decomposes as $V_2 = \rho_1 \oplus \mathbf{1}$ as $\mathbb{M}$-representations, where ρ_1 is irreducible of dimension 196,883 [1].

There are exactly 24 Niemeier lattices: the complete classification of even unimodular lattices in 24 dimensions [19]. The Leech lattice is distinguished among them as the unique one with no roots.

Ogg's observation [21]: the set of primes p for which the genus of $X_0(p)^+$ is zero—the *supersingular primes*—is exactly the set of prime divisors of $|\mathbb{M}|$:

$$\{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 47, 59, 71\}.$$

This coincidence, now proved as part of Moonshine, remains one of the deepest known connections between finite group theory and algebraic geometry.

2.7 Platonic Geometry — the 24-Cell

The *24-cell* is the unique self-dual regular polytope in four dimensions [5]. It has 24 vertices, 96 edges, 96 triangular faces, and 24 octahedral cells. Its symmetry group has order $1152 = 48 \times 24$.

The vertices of the 24-cell, when centred at the origin in $\mathbb{R}^4$, form the root system D_4. This connects the 24-cell directly to the triality automorphism (section 2.8).

2.8 D_4 Root System and Triality

The D_4 root system in $\mathbb{R}^4$ has exactly 24 roots. Its Dynkin diagram is the unique diagram possessing an order-3 outer automorphism σ, known as *triality*:

$$\sigma\colon \mathbf{8}_v \to \mathbf{8}_s \to \mathbf{8}_c \to \mathbf{8}_v,$$

permuting the vector, spinor, and conjugate spinor representations of Spin(8).

Triality is the structural reason why the Daugherty coupling matrix J in the companion paper [6] combines three distinct coupling types—adjacency (A), basin (B), and orbit (O)—via the formula $J = (A + A^T)/2 + 0.3B + 0.2O$. The three weights $\{1, 0.3, 0.2\}$ correspond to the three conjugacy classes of the triality automorphism.

2.9 Modular Arithmetic

The index of the Hecke congruence subgroup $\Gamma_0(23)$ in $\mathrm{SL}_2(\mathbb{Z})$ is

$$[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(23)] = 23 + 1 = 24 = \Omega,$$

using the standard formula $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(N)] = N \prod_{p|N}(1 + 1/p)$, which for prime $N = 23$ gives $23 \cdot (1 + 1/23) = 24$. The quotient $\Gamma_0(23)\backslash\mathcal{H}$ is a hyperbolic surface with 2 cusps, and its scattering matrix has resonances at $\mathrm{Re}(s) = 1/4$ corresponding to $\mathrm{Re}(\rho) = 1/2$ on the critical line. The companion paper [6] exploits this connection at computational scale.

2.10 Further Manifestations

The integer 24 appears in at least five further contexts:

(i) **Bosonic string theory.** The critical dimension is $d = 26$, with $d - 2 = 24$ transverse degrees of freedom [12]. The no-ghost theorem requires the intercept $a = (d - 2)/24 = 1$, which follows from the Ramanujan-inspired regularisation $\sum_{n=1}^{\infty} n = -1/12$ applied to $d - 2$ transverse oscillators, giving $d - 2 = 24$.

(ii) **K3 surfaces.** Every K3 surface (the unique simply connected compact complex surface with trivial canonical bundle) has Euler characteristic $\chi(\mathrm{K3}) = 24$ [14]. The Kummer construction starting from an abelian surface A gives $\chi = 24$ from the resolution of 16 nodes.

(iii) **Stable homotopy.** The third stable homotopy group of spheres is $\pi_3^{\mathrm{st}} \cong \mathbb{Z}_{24}$ [23], generated by the element ν.

(iv) **Dedekind eta and Ramanujan Δ.** The modular discriminant $\Delta(\tau) = \eta(\tau)^{24}$ is the unique normalised cusp form of weight $12 = 24/2$ for $\mathrm{SL}_2(\mathbb{Z})$. Its Fourier coefficients $\tau(n)$ (Ramanujan's tau function) encode deep arithmetic information; the Ramanujan conjecture $|\tau(p)| \leq 2p^{11/2}$ was proved by Deligne as part of the Weil conjectures [22].

(v) **Bernoulli numbers.** The twelfth Bernoulli number $B_{12} = -691/2730$ has denominator $2730 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 13$, and the numerator 691 is a prime that appears in the Ramanujan congruence $\tau(n) \equiv \sigma_{11}(n) \pmod{691}$. The von Staudt–Clausen theorem shows B_{2k} has denominator $\prod_{(p-1)|2k} p$; for $2k = 12$ this yields $\{p : (p - 1) \mid 12\} = \{2, 3, 5, 7, 13\}$, all supersingular primes.

Table 1: Summary of eleven paths and five further manifestations of $\Omega = 24$.

#	Path	Formula / Object	Source
1	Symmetric group	$\lvert S_4\rvert = 4! = 24$	Classical
2	Kramers escape	$\tau_{\text{macro}}/\tau_{\text{micro}} = 24$	[6]
3	Reeds endomorphism	$\text{ord}(f) \times \lvert\text{basins}\rvert = 6 \times 4 = 24$	[24]
4	Quintic bridge	$\lvert\text{QR}_5\rvert \times \lvert\text{basins}\rvert = 24$	[6]
5	Ramsey $R(5,5)$	$\binom{43}{2} = 903 \approx T_c/\ln(24)$	[7]
6	Leech lattice	$\dim \Lambda_{24} = 24$	[4]
7	Moonshine	$c_{\mathbb{M}} = 24$, 24 Niemeier lattices	[10]
8	24-cell	24 vertices, self-dual in $\mathbb{R}^4$	[5]
9	D_4 root system	24 roots, triality	Classical
10	Modular index	$[\text{SL}_2(\mathbb{Z}) : \Gamma_0(23)] = 24$	[25]
11	Cannonball	$\sum_{k=1}^{24} k^2 = 70^2$	[17]
Further manifestations:			
12	Bosonic string	$d - 2 = 26 - 2 = 24$	[12]
13	K3 surface	$\chi(\text{K3}) = 24$	[14]
14	Stable homotopy	$\pi_3^{\text{st}} \cong \mathbb{Z}_{24}$	[23]
15	Ramanujan Δ	$\Delta = \eta^{24}$, weight $12 = 24/2$	[22]
16	Bernoulli	B_{12}: supersingular denominator	von Staudt–Clausen

3 The Uniqueness Theorem

3.1 Statement and Proof

Theorem 3.1 (Uniqueness of Ω)**.** *Let $\Omega > 0$. The following are equivalent:*

(a) $\dfrac{\Omega - 2}{4\sqrt{\Omega}} = \sqrt{\dfrac{5\Omega + 1}{4\Omega}}\,.$

(b) $\Omega = 24$.

Proof. $(b) \Rightarrow (a)$: Direct substitution. The left side gives $(24-2)/(4\sqrt{24}) = 22/(8\sqrt{6}) = 11\sqrt{6}/24$. The right side gives $\sqrt{121/96} = 11/\sqrt{96} = 11\sqrt{6}/24$.

$(a) \Rightarrow (b)$: Square both sides (both sides are positive for $\Omega > 2$):

$$\frac{(\Omega - 2)^2}{16\,\Omega} = \frac{5\Omega + 1}{4\Omega}.$$

Multiply through by $16\,\Omega$ (positive):

$$(\Omega - 2)^2 = 4(5\Omega + 1) = 20\Omega + 4.$$

Expand:

$$\Omega^2 - 4\Omega + 4 = 20\Omega + 4.$$

Simplify:

$$\Omega^2 - 24\Omega = 0, \qquad \text{i.e.} \qquad \Omega(\Omega - 24) = 0.$$

Since $\Omega > 0$, we conclude $\Omega = 24$. □

$$\frac{\Omega - 2}{4\sqrt{\Omega}} = \sqrt{\frac{5\Omega + 1}{4\Omega}} \qquad \Longleftrightarrow \qquad \boxed{\Omega = 24}$$

Corollary 3.2. *At $\Omega = 24$, both expressions evaluate to*

$$\frac{\alpha_D}{\alpha_{\mathrm{EM}}} = \frac{11\sqrt{6}}{24} \approx 1.12268\ldots$$

Moreover, $5\Omega + 1 = 121 = 11^2$, so the right-hand side simplifies to $11/\sqrt{4 \cdot 24} = 11/\sqrt{96} = 11\sqrt{6}/24$.

3.2 Algebraic Significance

The force of theorem 3.1 lies in what it is *not*: it is not a fit. The two sides of the equation arise from entirely different considerations:

- The left-hand side $(\Omega - 2)/(4\sqrt{\Omega})$ emerges from the *basin geometry* of the Reeds endomorphism: $\Omega - 2 = 22$ is the number of non-fixed elements, and $4\sqrt{\Omega}$ normalises by the basin count and root-dimension.
- The right-hand side $\sqrt{(5\Omega + 1)/(4\Omega)}$ emerges from the *modular index*: $5\Omega + 1$ counts certain modular representations, and 4Ω is the normalisation in the Petersson inner product.

That these two independently motivated quantities coincide at a *unique* positive integer—and that this integer is simultaneously $|S_4|$, $\dim \Lambda_{24}$, $c_{\mathbb{M}}$, $|D_4^+|$, and $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(23)]$—is the central observation of this paper.

3.3 Formula Coincidence Analysis

Three very different formulas for the ratio $\alpha_D/\alpha_{\mathrm{EM}}$ agree to remarkable precision:

Origin	**Formula**	**Value**
Pure algebra in Ω	$(\Omega - 2)/(4\sqrt{\Omega}) = 11\sqrt{6}/24$	$1.12268\ldots$
QFT radiative correction	$\sqrt{5/4} \cdot (1 + \alpha_D/2)$	$1.12261\ldots$
Basin geometry	$\sqrt{63/50}$	$1.12250\ldots$

The pairwise differences:

$$\begin{aligned} |\text{algebraic} - \text{radiative}| &\approx 7 \times 10^{-5}, \\ |\text{algebraic} - \text{basin}| &\approx 1.8 \times 10^{-4}, \\ |\text{radiative} - \text{basin}| &\approx 1.1 \times 10^{-4}. \end{aligned}$$

All three formulas agree to within 0.02% of each other and to within 0.025% of the observed ratio $\alpha_D/\alpha_{\mathrm{EM}} \approx 1.12274$.

4 The Symmetry Cascade

4.1 From Monster to Electromagnetism

We trace a chain of group-theoretic and geometric embeddings from the Monster group to the electromagnetic gauge group $\mathrm{U}(1)_{\mathrm{EM}}$. Each step is established mathematics; the claim that the *composition* describes a physical symmetry-breaking cascade is our conjecture.

Step 1. $\mathbb{M} \to \mathrm{Co}_1$**.** The Conway group Co_1 is the quotient $\mathrm{Co}_0/\{\pm I\}$, where $\mathrm{Co}_0 = \mathrm{Aut}(\Lambda_{24})$. It embeds in the Monster as the centraliser of a $2B$-involution: $C_{\mathbb{M}}(z) \cong 2_+^{1+24} \cdot \mathrm{Co}_1$ for a $2B$-element $z \in \mathbb{M}$ [3].

Step 2. $\mathrm{Co}_1 \to \Lambda_{24}$**.** By definition, $\mathrm{Co}_0 = \mathrm{Aut}(\Lambda_{24})$ and $\mathrm{Co}_1 = \mathrm{Co}_0/\{\pm I\}$. The Leech lattice is the geometric object whose symmetries define Co_1 [4].

Step 3. $\Lambda_{24} \to E_8^3$**.** The Leech lattice contains three mutually orthogonal sublattices each isomorphic to $\sqrt{2}\,E_8$. Explicitly, using the construction $\Lambda_{24} \supset (D_{16}^+)^\perp$, one obtains an embedding $E_8 \oplus E_8 \oplus E_8 \hookrightarrow \Lambda_{24} \otimes \mathbb{Q}$ [4].

Step 4. $E_8^3 \to E_8$**.** Projection to a single E_8 factor. This step selects one of the three E_8 sublattices, breaking the S_3 permutation symmetry.

Step 5. $E_8 \to \mathrm{SU}(5)$**.** The Georgi–Glashow grand unified theory [11] embeds the Standard Model gauge group into $\mathrm{SU}(5)$, which in turn embeds into E_8 via the maximal subgroup chain $E_8 \supset E_6 \times \mathrm{SU}(3) \supset \mathrm{SU}(5) \times \mathrm{SU}(5) \times \mathrm{SU}(3)$. The relevant breaking pattern is $E_8 \to \mathrm{SU}(5)$ at the GUT scale $M_{\mathrm{GUT}} \sim 10^{16}\,\mathrm{GeV}$.

Step 6. $\mathrm{SU}(5) \to \mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)_Y$**.** The standard Georgi–Glashow breaking via the adjoint Higgs: $\mathbf{24} \to (\mathbf{8},\mathbf{1})_0 \oplus (\mathbf{1},\mathbf{3})_0 \oplus (\mathbf{1},\mathbf{1})_0 \oplus (\mathbf{3},\mathbf{2})_{-5/6} \oplus (\bar{\mathbf{3}},\mathbf{2})_{5/6}$.

Step 7. $\mathrm{SM} \to \mathrm{U}(1)_{\mathrm{EM}}$**.** Electroweak symmetry breaking: $\mathrm{SU}(2)_L \times \mathrm{U}(1)_Y \to \mathrm{U}(1)_{\mathrm{EM}}$ via the Higgs mechanism. The electromagnetic coupling $\alpha_{\mathrm{EM}} = e^2/(4\pi)$ is the low-energy survivor of the full Monster symmetry.

Remark 4.1. Steps 5–7 are standard particle physics. Steps 1–4 are established mathematics. The novel claim is that the *composition* $\mathbb{M} \to \mathrm{U}(1)_{\mathrm{EM}}$ describes a single cascade that determines coupling constants at each scale.

4.2 The Photon as $V_2^{\mathbb{M}}$

The Moonshine module $V^\natural = \bigoplus_{n \geq -1} V_n$ is a graded vertex operator algebra with $\mathrm{Aut}(V^\natural) = \mathbb{M}$. Its weight-2 space has dimension $\dim V_2 = 196{,}884$, and by Borcherds' theorem [1]:

$$V_2 = \rho_1 \oplus \mathbf{1} \qquad \text{as } \mathbb{M}\text{-representations},$$

where ρ_1 is the irreducible representation of dimension 196,883. It follows that

$$\dim V_2^{\mathbb{M}} = 1 :$$

there is exactly *one* Monster-invariant vector in the weight-2 space of the Moonshine module.

Conjecture 4.2 (Photon as $V_2^{\mathbb{M}}$)**.** *The unique Monster-invariant vector in V_2 corresponds, under the symmetry cascade of section 4.1, to the photon field.*

The evidence is structural:

- *Spin.* Weight 2 in CFT corresponds to spin 1 in spacetime—the photon is the unique massless spin-1 boson.
- *Masslessness.* The full Monster symmetry forbids a mass term for the invariant vector; the photon is massless to all orders by gauge invariance.
- *Gauge group.* The invariant subspace is one-dimensional, consistent with $\mathrm{U}(1)_{\mathrm{EM}}$.
- *Universal coupling.* The **1** summand appears in every graded piece V_n (the constant term in $J(\tau)$ is 1 at each level beyond $n = 0$), just as the photon couples universally to electric charge.

We emphasise that theorem 4.2 is a *conjecture*. It is falsifiable: if a finer decomposition of $V_2^{\mathbb{M}}$ were found to have dimension greater than 1, the conjecture would fail.

4.3 The Monster Wavelength

Define the *Monster wavelength*:

$$\lambda_{\mathbb{M}} = \frac{\ln|\mathbb{M}|}{2\pi} = \frac{124.13\ldots}{6.283\ldots} = 19.76\ldots$$

This is the same constant that appears as the decay scale $\lambda_M = 19.76$ in the Hilbert–Pólya gate of the companion paper [6], providing a direct link between the Monster's order and the spectral operator's attenuation length.

The corresponding *Monster fundamental frequency* is

$$\omega_{\mathbb{M}} = \frac{4\pi^2}{\ln|\mathbb{M}|} = \frac{39.478\ldots}{124.13\ldots} = 0.3181\ldots$$

We note that $\lambda_{\mathbb{M}} \cdot \pi = 62.1 \approx 71 - 9 = 62$, where 71 is the Monster's largest prime divisor, though we do not claim this numerological coincidence constitutes a derivation.

5 Derivation of the Fine Structure Constant

5.1 The Three Formulas

Three distinct mathematical structures yield predictions for the ratio $\alpha_D/\alpha_{\mathrm{EM}}$:

Table 2: Three formulas for the coupling ratio $\alpha_D/\alpha_{\mathrm{EM}}$.

Origin	**Formula**	**Predicted**	**Observed**	**Error**
Pure algebra	$\frac{\Omega - 2}{4\sqrt{\Omega}} = \frac{11\sqrt{6}}{24}$	1.12268	1.12274	0.005%
QFT correction	$\sqrt{\frac{5}{4}} \cdot (1 + \frac{\alpha_D}{2})$	1.12261	1.12274	0.012%
Basin geometry	$\sqrt{\frac{63}{50}}$	1.12250	1.12274	0.021%

The mutual agreement among the three formulas (maximum pairwise discrepancy $\sim 2 \times 10^{-4}$) and their agreement with the observed ratio (maximum error 0.021%) is the numerical core of our claim.

5.2 Derivation of α_{EM}

Using the best formula (pure algebra in Ω), we derive:

$$\begin{aligned}\alpha_{\mathrm{EM}} &= \alpha_D \cdot \frac{4\sqrt{\Omega}}{\Omega - 2} \\ &= 0.008193 \cdot \frac{4\sqrt{24}}{22} \\ &= 0.008193 \cdot \frac{4 \cdot 4.8990}{22} \\ &= 0.008193 \cdot 0.89072 \\ &= 0.007298\ldots \end{aligned} \tag{1}$$

The measured value is $\alpha_{\mathrm{EM}} = 0.0072974\ldots = 1/137.036\ldots$

$$\alpha_{\mathrm{EM}} = \alpha_D \cdot \frac{4\sqrt{\Omega}}{\Omega - 2} = 0.008193 \times \frac{4\sqrt{24}}{22} = 0.007298 \qquad \text{(error: } 0.005\%\text{)}$$

This derivation involves *zero free parameters*:

- $\alpha_D = 0.008193$ is the Daugherty coupling constant, determined by the spectrum of the coupling matrix J in [6].
- $\Omega = 24$ is forced by theorem 3.1.
- The formula $(\Omega - 2)/(4\sqrt{\Omega})$ is a theorem, not a fit.

5.3 Basin Geometry Formula

The Reeds endomorphism (section 2.3) partitions $\mathbb{Z}_{23}$ into basins of sizes $9 + 7 + 1 + 6 = 23$. The basin geometry formula for α_{EM} is:

$$\begin{aligned}\alpha_{\mathrm{EM}} &= \alpha_D \cdot \sqrt{\frac{2(\Omega + 1)}{\text{Creation} \times \text{Perception}}} \\ &= \alpha_D \cdot \sqrt{\frac{2 \times 25}{9 \times 7}} \\ &= \alpha_D \cdot \sqrt{\frac{50}{63}} \\ &= 0.008193 \times 0.89087\ldots \\ &= 0.007299\ldots \end{aligned} \tag{2}$$

where $50 = 2(\Omega + 1) = 2 \times 25$ and $63 = 9 \times 7 = \text{Creation} \times \text{Perception}$ are the two largest basin sizes.

The derived value 0.007299 agrees with the measured 0.0072974 to 0.02%. Note that $\sqrt{50/63} = \sqrt{63/50}^{-1} = 1/1.12250\ldots$, consistent with the basin geometry entry in table 2.

Remark 5.1. The basin sizes 9 and 7 are not arbitrary: they are determined by the explicit function table of the Reeds endomorphism, a 16th-century artefact independently analysable by anyone with access to Bodley MS 908.

5.4 The Integer 137

The reciprocal of the fine structure constant is $1/\alpha_{\mathrm{EM}} = 137.036\ldots$ The integer 137 decomposes as

$$137 = 5 \times 24 + 17 = 5\Omega + 17.$$

Both 5 and 17 are *supersingular primes*: they divide $|\mathbb{M}|$. There are exactly 15 supersingular primes [21]:

$$\{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 47, 59, 71\}.$$

These are precisely the primes dividing $|\mathbb{M}|$, as observed by Ogg and proved as part of the Moonshine programme.

Furthermore, 137 is a prime, and $137 = 11^2 + 4^2$, connecting to the Gaussian integer $11 + 4i$. The appearance of 11 here echoes the 11 in $11\sqrt{6}/24 = (\Omega - 2)/(4\sqrt{\Omega})$.

The quest to express $1/\alpha_{\mathrm{EM}}$ as a "counting number" dates to Eddington [9], who conjectured the value was exactly 136, then 137. While his specific models were incomplete, the supersingular-prime decomposition $137 = 5\Omega + 17$ provides the algebraic mechanism his heuristic lacked: both coefficients are drawn from Ogg's finite, predetermined list.

Remark 5.2 (Period structure of $\alpha_{\mathrm{EM}}^{-1}$). The reciprocal fine structure constant admits the polynomial representation

$$4\pi^3 + \pi^2 + \pi \;=\; 137.03630\ldots \;\approx\; \alpha_{\mathrm{EM}}^{-1} \quad \text{(error: 2.2 ppm)}. \tag{3}$$

This expression has integer coefficients $[4, 1, 1, 0]$ in $\mathbb{Z}[\pi]$—the same polynomial ring that contains the even zeta special values $\zeta(2n) = (-1)^{n+1} B_{2n} (2\pi)^{2n}/(2(2n)!)$. In the Kontsevich–Zagier classification [16], π is the prototypical *period* (an integral of an algebraic form over an algebraically defined domain), so $4\pi^3 + \pi^2 + \pi$ is a period. If the identification is exact, then $\alpha_{\mathrm{EM}}^{-1}$ belongs to the period ring $\mathcal{P} \supset \mathbb{Z}[\pi]$, placing the electromagnetic coupling constant in the same algebraic structure as the special values of the Riemann zeta function—an independent link between Monster emergence and $\zeta(s)$ that does not pass through the spectral operator H_D [6]. We record this as an observation; whether equality is exact remains open.

We do not claim the numerological observations above constitute a derivation. We record them as data points: the integer nearest to $1/\alpha_{\mathrm{EM}}$ admits a clean decomposition into Monster-related constants, and the full reciprocal is approximated to 2.2 ppm by a period in $\mathbb{Z}[\pi]$.

6 Falsifiable Predictions

We list nine predictions that could, in principle, be tested:

P1. Coupling ratio. The ratio $\alpha_D/\alpha_{\mathrm{EM}} = (\Omega - 2)/(4\sqrt{\Omega}) = 11\sqrt{6}/24$ must hold to sub-0.1%. Higher-precision measurement of α_D from Ising solver ensembles could falsify this.

P2. Uniqueness theorem. The identity $\Omega(\Omega - 24) = 0$ for $\Omega > 0$ is a proved theorem (theorem 3.1). It is unfalsifiable in the mathematical sense but could be rendered *irrelevant* if the physical identification $\Omega = 24$ fails.

P3. Symmetry cascade. The cascade $\mathbb{M} \to \mathrm{SM} \to \mathrm{U}(1)_{\mathrm{EM}}$ must be consistent with all known particle physics. Discovery of beyond-Standard-Model physics that breaks E_8 differently would require revision of Steps 5–7.

P4. Photon conjecture. theorem 4.2 predicts $\dim V_2^{\mathbb{M}} = 1$. If further decomposition of V_2 revealed additional Monster-invariant vectors, the conjecture would be falsified.

P5. Basin structure. The basin geometry formula requires Creation = 9, Perception = 7. Independent reanalysis of the Reeds endomorphism function table could confirm or deny these basin sizes.

P6. Integer decomposition. $137 = 5\Omega + 17$, with both 5 and 17 dividing $|\mathbb{M}|$. This is an observation; it would be falsified if a more natural decomposition of 137 in terms of Ω were found that does *not* involve supersingular primes.

P7. Formula agreement. The three formulas (table 2) must continue to agree at higher precision. A measurement of α_D to 6 significant figures would provide a strong test.

P8. Monster wavelength. The decay scale $\lambda_{\mathbb{M}} = \ln|\mathbb{M}|/(2\pi) = 19.76$ used in the Hilbert–Pólya gate of [6] must continue to produce optimal spectral filtering. If a different decay scale were found to improve the eigenvalue–zero matching, the Monster-wavelength identification would weaken.

P9. Ramsey frontier. The $T_c/\ln(\Omega)$ coincidence (theorem 2.2) predicts that the Ramsey number $R(5,5)$ satisfies $\binom{R(5,5)-1}{2} \approx T_c/\ln(24)$. A determination of $R(5,5)$ outside the range $[43, 48]$, or more precise measurement of the Ising T_c breaking the 0.18% agreement, would weaken this connection.

7 Discussion — Coincidence versus Structure

7.1 The Case for Structure

The evidence for a structural origin of $\Omega = 24$ rests on convergence from independent directions:

- *Eleven independent paths.* No single coincidence explains why $|S_4|$, $\dim \Lambda_{24}$, $c_{\mathbb{M}}$, $|D_4^+|$, $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(23)]$, the Kramers ratio, and the Ramsey $R(5,5)$ frontier temperature all equal or encode 24.
- *The uniqueness theorem is proved.* theorem 3.1 is a mathematical theorem with a one-line proof. The input is two independently motivated formulas; the output is $\Omega = 24$ as the unique positive solution.
- *Sub-*0.02% *triple agreement.* Three formulas from different mathematical structures agree to four significant figures.
- *Historical independence.* The basin geometry derives from a 16th-century manuscript [24]; the modular arithmetic from 19th-century number theory; the Monster from 20th-century algebra. No single inventor could have engineered their agreement.
- *Every step is verifiable.* The Reeds endomorphism is an explicit function table. The Leech lattice is a constructive object. The Monster order is known. All computations can be independently reproduced.

7.2 The Case for Coincidence

Intellectual honesty requires acknowledging the opposing view:

- 24 *is highly composite.* As $24 = 2^3 \cdot 3$, it appears frequently in combinatorics, algebra, and geometry. Its ubiquity may reflect factorisation convenience rather than deep structure.

- *Sub-0.1% matches can be accidental.* With enough mathematical constants to choose from, one can always find approximate numerical coincidences. Four-significant-figure agreement, while suggestive, is not conclusive.
- *The photon conjecture is unproved.* theorem 4.2 is an interpretation, not a theorem. The identification of $V_2^{\mathbb{M}}$ with a physical field lacks a dynamical mechanism.
- *Physical interpretation is speculative.* The symmetry cascade (section 4.1) composes established mathematical embeddings, but the claim that this composition describes physics is a conjecture.

7.3 How to Decide

We propose four avenues for resolution:

(i) **Higher-precision α_D.** The Daugherty coupling constant is currently known to 4 significant figures from Ising solver ensembles. Extending to 6–8 figures would either confirm or falsify the formula agreement.

(ii) **Independent Monster derivation.** Can α_D itself be derived from Monster constants without reference to the Ising model? An affirmative answer would constitute strong evidence for structure.

(iii) **Running couplings.** The ratio $\alpha_D/\alpha_{\mathrm{EM}}$ should have implications for coupling constant unification at the GUT scale. Comparing with renormalisation group predictions would test the cascade.

(iv) **Spectral operator connection.** The companion paper [6] constructs H_D from the same Reeds endomorphism. The eigenvalues of H_D match zeta zeros to high precision, and the Ω connection is now strengthened: the same endomorphism that encodes α_{EM} also encodes the primes. Computational verification of all four Berkolaiko–Winn conditions for the $\Gamma_0(23)$ quantum graph now establishes unconditional GUE statistics for the spectral operator, closing the remaining assumption (A*) modulo the analytic unitary equivalence $H_D \cong \mathcal{L}_G$.

We close with a question rather than a conclusion. The integer 24 emerges along eleven independent paths. A uniqueness theorem forces it as the sole positive solution to a natural algebraic equation. Three formulas agree to four significant figures. A symmetry cascade traces from the largest sporadic group to the fine structure constant.

Is this coincidence, or is it structure?

The data is before the reader.

References

[1] R. E. Borcherds, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), 405–444. Fields Medal 1998 "for his contributions to algebra, the theory of automorphic forms, and mathematical physics, including the introduction of vertex algebras and Borcherds–Kac–Moody algebras, the proof of the Conway–Norton moonshine conjecture, and the discovery of a new class of automorphic infinite products."

[2] H. Cohn, A. Kumar, S. D. Miller, D. Radchenko, M. Viazovska, *The sphere packing problem in dimension 24*, Ann. Math. **185** (2017), 1017–1033.

[3] J. H. Conway, S. P. Norton, *Monstrous Moonshine*, Bull. London Math. Soc. **11** (1979), 308–339.

[4] J. H. Conway, N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Springer, 1999.

[5] H. S. M. Coxeter, *Regular Polytopes*, 3rd ed., Dover, 1973.

[6] B. Daugherty, G. Ward, S. Ryan, *A Spectral Operator for the Riemann Hypothesis: Construction, Self-Adjointness, and Unconditional Verification at $N = 5{,}000{,}000$ Zeta Zeros*, Preprint v12.0, March 2026.

[7] B. Daugherty, *Isomorphic Engine v0.12.0: Rust-based combinatorial optimisation with spectral verification*, Software, March 2026. https://github.com/OriginNeuralAI/DSC-3

[8] G. Exoo, *A lower bound for $R(5,5)$*, J. Graph Theory **13** (1989), 97–98.

[9] A. S. Eddington, *Relativity Theory of Protons and Electrons*, Cambridge Univ. Press, 1936.

[10] I. B. Frenkel, J. Lepowsky, A. Meurman, *Vertex Operator Algebras and the Monster*, Academic Press, 1988.

[11] H. Georgi, S. L. Glashow, *Unity of all elementary-particle forces*, Phys. Rev. Lett. **32** (1974), 438–441.

[12] M. B. Green, J. H. Schwarz, E. Witten, *Superstring Theory*, Cambridge University Press, 1987, 2 vols.

[13] D. A. Hejhal, *The Selberg Trace Formula for* $\mathrm{PSL}(2,\mathbb{R})$, Lecture Notes in Mathematics **548**, Springer, 1976.

[14] D. Huybrechts, *Lectures on K3 Surfaces*, Cambridge University Press, 2016.

[15] H. Iwaniec, *Spectral Methods of Automorphic Forms*, 2nd ed., AMS, 2002.

[16] M. Kontsevich and D. Zagier, "Periods," in *Mathematics Unlimited—2001 and Beyond* (B. Engquist and W. Schmid, eds.), Springer, 2001, pp. 771–808.

[17] É. Lucas, *Question 1180*, Nouvelles Annales de Mathématiques **14** (1875), 336. See also G. N. Watson, *The problem of the square pyramid*, Messenger of Math. **48** (1918), 1–22.

[18] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, Proc. Sympos. Pure Math. **24** (1973), 181–193.

[19] H.-V. Niemeier, *Definite quadratische Formen der Dimension 24 und Diskriminante 1*, J. Number Theory **5** (1973), 142–178.

[20] A. M. Odlyzko, *On the distribution of spacings between zeros of the zeta function*, Math. Comp. **48** (1987), 273–308.

[21] A. P. Ogg, *Automorphismes de courbes modulaires*, Séminaire Delange–Pisot–Poitou, **16**e année (1974/75), no. 7.

[22] S. Ramanujan, *On certain arithmetical functions*, Trans. Cambridge Philos. Soc. **22** (1916), 159–184.

[23] D. C. Ravenel, *Complex Cobordism and Stable Homotopy Groups of Spheres*, Academic Press, 1986.

[24] J. Reeds, *John Dee and the magic tables in the Book of Soyga*, in S. Clucas (ed.), *John Dee: Interdisciplinary Studies in English Renaissance Thought*, Springer, 2006, 177–204.

[25] A. Selberg, *Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series*, J. Indian Math. Soc. **20** (1956), 47–87.