




Characterizing word processing pathway using frequency tagging

Kishen Senziani¹, Berk Gercek¹, Nina Kazanina¹; ¹University of Geneva, Faculty of Medicine, Department of Basic Neurosciences

A **6 / 7.06 Hz flicker tag** tracks the visual input, not wordness; the **word-locked response** (250–600 ms) separates words from consonant strings, and in anterior temporal cortex the difference is sustained across words.



Update vs. submitted abstract: with fixation baselines and equal trial counts, the tag shows no word-specific effect and the source effect is Word > Non-word.

Background



- AIM 1**
Do MEG sensors and sources recover the V1 → vOT → temporal sequence of the literature?
- AIM 2**
Does the tag response (SSVEF) differ between words and consonant strings?

Methods



Pitfalls we hit — and how we fixed them

- Tags off by one FFT bin.** The display ran ~0.5 % slow (photodiode: 5.965 / 7.024 Hz): read the SNR at the measured bin.
- Unequal trial counts faked an effect.** A time-resolved source effect vanished once counts were equalized (the whole-epoch effect remains).
- No gap between words.** The pre-stimulus window holds the previous word: baseline on fixation instead.
- Check the noise-covariance rank.** After SSS, 3 subjects kept a ~1e-38 eigenvalue (dSPM × 10⁶, distorted maps): fixed with `mne.compute_rank(tol='auto')`.



Code · supplementary figures S1–S4 · cohort
github.com/Orion212-lab/snl2026-ft-word-pathway

Cohort 30 recorded; each panel states its N. Excluded: no flicker on the photodiode, malformed blocks, all epochs rejected; sources: no T1 MRI, artifact (sub-20/24). Kept with trigger anomalies: sub-10–14 (main results unchanged without them). List via QR.


A priori Word vs Non-word at the tag and in the evoked cascade windows (80–130, 150–225, 250–600 ms), left-temporal sensors. Not preregistered; everything else is exploratory.

Limits consonant strings differ in orthography, pronounceability, visual complexity, difficulty · condition-pure blocks · late vOT fragile · localizer, catch-query accuracy not yet analysed

Next pseudowords · item-rate (FPVS) tagging instead of contrast flicker · interleaved conditions · two-word phrase task (intermodulation), recorded, not yet analysed


Results

1 · Sensor space: the word-locked response separates words from consonant strings (Aim 1)




One Word > Non-word cluster over left posterior and temporal sensors (N = 26, 84 sensors, $p < .001$), robust to blinks, saccades and artifacts (clean epochs: $dz = 1.31$, 20/21 subjects). Tag × Wordness: n.s.

2 · Sources: Word > Non-word, maximal and sustained in anterior temporal cortex (Aim 1)





3 · The tag follows the visual input, not wordness (Aim 2)



Tag (harmonics 1–4): occipital Non-words > Words ($dz = -.78$, $p = .006$), matching their extra visual complexity (+10 % ink); left-temporal Words = Non-words ($dz = -.14$, $BF01 = 3.5$). **Evoked** 250–600 ms, the reverse: left-temporal Words > Non-words ($dz = .93$, $p = .002$; Tag × Wordness n.s.), occipital n.s.

4 · Direct test: the word-locked response beats the tag, ROI by ROI (Aim 2)



Aim 1 — partly: temporal stages replicated, vOT only late and fragile, latency order n.s. **Aim 2 — no:** the tag indexes visual input, not wordness; the word-locked response does.

References

- Salmelin, R. 2007. “Clinical Neurophysiology of Language: The MEG Approach.” *Clinical Neurophysiology* 118, no. 2: 237–254. <https://doi.org/10.1016/j.clinph.2006.07.316>.
- Samoylov, A., Semenova, N., Myachikov, A., & Shtyrov, Y. (2025). Patterns of cortical activity in a silent single-word reading task depend on word frequency and age-related differences: An MEG study. *Psychophysiology*, 62(3), e70192. <https://doi.org/10.1111/psyp.70192>.
- Thesen, T., McDonald, C. R., Carlson, C., Doyle, W., Cash, S., Sherley, J., Felsovalyi, O., Girard, H., Barr, W., Devinsky, O., Kuzniecky, R., & Halgren, E. (2012). Sequential then interactive processing of letters and words in the left fusiform gyrus. *Nature Communications*, 3, 1284. <https://doi.org/10.1038/ncomms2220>.
- Cornelissen, P. L., Kringelbach, M. L., Ellis, A. W., Whitney, C., Holliday, I. E., & Hansen, P. C. (2020). Activation of the left inferior frontal gyrus in the first 200 ms of reading: Evidence from magnetoencephalography (MEG). *PLoS ONE*, 15(4), e023559. <https://doi.org/10.1371/journal.pone.0235559>.
- Norcia, A. M., Appelbaum, L. G., Ales, J. M., Cottareau, B. R., & Rossion, B. (2015). The steady-state visual evoked potential in vision research: A review. *Journal of Vision*, 15(6), 4. <https://doi.org/10.1167/15.6.4>.
- Figueira, J. S. B., Kutlu, E., Scott, L. S., & Keil, A. (2022). The FreqTag toolbox: A principled approach to analyzing electrophysiological time series in frequency tagging paradigms. *Developmental Cognitive Neuroscience*, 54, 101066.
- Kabdebon, C., Fló, A., De Heering, A., & Aslin, R. (2022). The power of rhythms: how steady-state evoked responses reveal early neurocognitive development. *NeuroImage*, 254, 119150.
- Lochy, A., Van Belle, G., & Rossion, B. (2015). A robust index of lexical representation in the left occipito-temporal cortex as evidenced by EEG responses to fast periodic visual stimulation. *Neuropsychologia*, 66, 18–31. <https://doi.org/10.1016/j.neuropsychologia.2014.11.007>.
- Wang, F., Kaneshiro, B., Strauber, C. B., Hasak, L., Nguyen, Q. T. H., Yakovleva, A., ... & McCandless, B. D. (2021). Distinct neural sources underlying visual word form processing as revealed by steady state visual evoked potentials (SSVEP). *Scientific reports*, 11(1), 18229.