

The SIRS Model for Viral Infection

Brianne Gill

May 16, 2008

Dividing and Describing a Population

When describing a population in relation to disease, you can divide all members of the population into three categories:

- ▶ Susceptible Individuals

Dividing and Describing a Population

When describing a population in relation to disease, you can divide all members of the population into three categories:

- ▶ Susceptible Individuals
- ▶ Infected Individuals

Dividing and Describing a Population

When describing a population in relation to disease, you can divide all members of the population into three categories:

- ▶ Susceptible Individuals
- ▶ Infected Individuals
- ▶ Recovered Individuals

Population Rates of Change

The rate of change of the Susceptible Population is given by,

- ▶ Amount of Increase, or gain from Recovered Population = γR ,

Population Rates of Change

The rate of change of the Susceptible Population is given by,

- ▶ Amount of Increase, or gain from Recovered Population = γR ,
- ▶ Amount of Decrease, or loss from Infection = $-\beta IS$.

Population Rates of Change

The rate of change of the Susceptible Population is given by,

- ▶ Amount of Increase, or gain from Recovered Population = γR ,
- ▶ Amount of Decrease, or loss from Infection = $-\beta IS$.
- ▶ Thus, the differential equation for the change in the Susceptible Population is

$$\frac{dS}{dt} = -\beta IS + \gamma R,$$

The Infected Population

The rate of change of the Infected Population is given by,

- ▶ Amount of Increase, or gain from Susceptible Population = βIS ,

The Infected Population

The rate of change of the Infected Population is given by,

- ▶ Amount of Increase, or gain from Susceptible Population = βIS ,
- ▶ Amount of Decrease, or loss from Recovery = $-\nu I$.

The Infected Population

The rate of change of the Infected Population is given by,

- ▶ Amount of Increase, or gain from Susceptible Population = βIS ,
- ▶ Amount of Decrease, or loss from Recovery = $-\nu I$.
- ▶ Therefore, the rate of change in the Infected Population is given by

$$\frac{dI}{dt} = \beta IS - \nu I.$$

The Recovered Population

- ▶ Amount of Increase, or gain from Infected Population = νI ,

The Recovered Population

- ▶ Amount of Increase, or gain from Infected Population = νI ,
- ▶ Amount of Decrease, or loss from immunity loss = $-\gamma R$.

The Recovered Population

- ▶ Amount of Increase, or gain from Infected Population = νI ,
- ▶ Amount of Decrease, or loss from immunity loss = $-\gamma R$.
- ▶ Consequently, the equation for the rate of change of the Recovered Population is

$$\frac{dR}{dt} = \nu I - \gamma R.$$


Summary of Our Equations

$$\begin{aligned}\frac{dS}{dt} &= -\beta IS + \gamma R, \\ \frac{dI}{dt} &= \beta IS - \nu I, \\ \frac{dR}{dt} &= \nu I - \gamma R.\end{aligned}$$



Figure: A diagram of the flow between the susceptible, infected, and recovered populations.

Three Related Populations



Reducing the SIRS System

► $N = S + I + R$

Reducing the SIRS System

- ▶ $N=S+I+R$
- ▶ $\frac{dS}{dt} = -\beta IS + \gamma(N - I - S),$

Reducing the SIRS System

- ▶ $N=S+I+R$
- ▶ $\frac{dS}{dt} = -\beta IS + \gamma(N - I - S),$
- ▶ $\frac{dI}{dt} = \beta IS - \nu I,$

Finding the Nullclines

► $\beta IS + \gamma S = \gamma(N - I),$

Finding the Nullclines

- ▶ $\beta IS + \gamma S = \gamma(N - I),$
- ▶ $S = \frac{\gamma}{\beta},$

Finding the Nullclines

- ▶ $\beta IS + \gamma S = \gamma(N - I),$
- ▶ $S = \frac{\gamma}{\beta},$
- ▶ $I = 0,$

The Nullclines of S' and I'

$$\begin{aligned} S' &= -BIS + y(N - I - S) & N &= 500 & v &= 25 \\ I' &= BIS - vI & B &= .3 & y &= .8 \end{aligned}$$




Figure: The Nullclines of I' and S' .

The Jacobian

$$J = \begin{pmatrix} -\beta I - \gamma & -\beta S - \gamma \\ \beta I & \beta S - \nu \end{pmatrix}.$$

Finding the Equilibrium Points

$$\blacktriangleright I = \frac{\gamma(N - S)}{BS + \gamma}$$

Finding the Equilibrium Points

- ▶ $I = \frac{\gamma(N - S)}{BS + \gamma}$
- ▶ $I = 12.9199$

Finding the Equilibrium Points

- ▶ $I = \frac{\gamma(N - S)}{BS + \gamma}$
- ▶ $I = 12.9199$
- ▶ $S = \frac{v}{B}$

Finding the Equilibrium Points

- ▶ $I = \frac{\gamma(N - S)}{BS + \gamma}$
- ▶ $I = 12.9199$
- ▶ $S = \frac{v}{B}$
- ▶ $S = 83.3333$

A Phase Plane Graph of S and I




Figure: An Equilibrium point at $(\frac{\nu}{\beta}, \beta IS = \gamma(N - I - S))$.

Finding the Equilibrium Points

► $I = 0$

Finding the Equilibrium Points

- ▶ $I = 0$
- ▶ $S = \frac{\gamma(N - I)}{BI + \gamma}$

Finding the Equilibrium Points

- ▶ $I = 0$
- ▶ $S = \frac{\gamma(N - I)}{BI + \gamma}$
- ▶ $S = 500$




Figure: An Equilibrium Point at $(N, 0)$.

The Threshold Effect

$$\blacktriangleright I = \frac{\gamma(N - S)}{\beta S - \gamma},$$

The Threshold Effect

- ▶ $I = \frac{\gamma(N - S)}{\beta S - \gamma},$
- ▶ $S = \frac{\nu}{B},$

The Threshold Effect

- ▶ $I = \frac{\gamma(N - S)}{\beta S - \gamma},$
- ▶ $S = \frac{\nu}{B},$
- ▶ $N > \frac{\nu}{B}.$




Figure: The plot of I versus S where $N < \frac{v}{\beta}$.