



Integrated Pest Management in the Tropics; pp. 219-247
© 2016, Editor, Dharam P. Abrol
New India Publishing Agency, New Delhi (India)

CHAPTER - 8

Integrated Pest Management in Tropical Vegetable Crops

R. Srinivasan

Entomologist, AVRDC – The World Vegetable Center, Shanhua, Tainan, Taiwan
email: srini.ramasamy@worldveg.org

8.1 Introduction

Vegetables are cultivated over an area of about 19 million ha globally, with an annual production of 268 million tonnes; Asia and Africa account for more than 90% of the hectareage and production (FAO, 2011). Vegetable crops contribute to human and environmental health. Most vegetables can supply plant proteins, vitamins (especially vitamins A and C), folic acid, minerals such as iron and calcium, and dietary fiber to the human diet (Fowler, 2011). Vegetable legumes also fix atmospheric nitrogen in the soil, thus improving soil fertility. Their haulms can be used as high quality livestock fodder. Most vegetables are high-value, repeat-cycle crops that can help lift small-scale farmers out of poverty in developing tropical countries (Genova *et al.*, 2010). However, vegetable production is often constrained by several severe biotic- and abiotic factors that reduce yields and profits.

Insect pests and diseases are one of the major production constraints in tropical vegetable production. For example, the eggplant fruit and shoot borer (EFSB), *Leucinodes orbonalis* Guenée (Lepidoptera: Pyralidae) can reduce yield by as much as 90% (Parimi and Zehr, 2009). Up to 80% yield losses have been reported in tomato due to damage by its fruit borer, *Helicoverpa armigera* Hubner (Lepidoptera: Noctuidae) (Tewari and Krishnamoorthy, 1984). Tomato

yellow leaf curl virus disease transmitted by the whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) can reach up to 100% incidence with yield losses often exceeding 90% (Saikia and Muniyappa, 1989). A loss of 30 to 100% can be caused in various cucurbits by the melon fly, *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae) (Dhillon *et al.*, 2005). As a major pest of vegetable brassicas, diamondback moth, *Plutella xylostella* (Linnaeus) (Lepidoptera: Plutellidae) causes up to 50% yield losses (Sandur, 2004). Significant yield losses of up to 50% have been attributed to onion thrips, *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) in bulb alliums (Mote, 1976).

Owing to the significant loss of marketable yield due to pests and diseases in the tropics, farmers tend to apply extremely high quantities of chemical pesticides to vegetable crops, often in an indiscriminating fashion. For example, farmers may spray pesticides more than 50 times on their tomato crops in a given season in India (Nagaraju *et al.*, 2002). Cambodian farmers spray up to 20 times per season using up to five different pesticides on major vegetable crops (Sodavy *et al.*, 2000). Cabbage producers in Benin applied approximately 46.8 l/ha of pesticide concentrate on the crop using 19 applications (Rosendahl *et al.*, 2008). Weekly spraying of pesticides in vegetable production is routine in Cameroon (Abang *et al.*, 2013).

With few vegetable pesticides available in sub-Saharan Africa, farmers apply pesticides registered for other crops, often in incorrect dosages; some may even use banned pesticides (Williamson, 2003; Amoah *et al.*, 2006). Pesticide misuse and/or overuse are thus not uncommon in the tropics, and lead to the development of resistance in target insects. Pesticides are often applied to vegetables as the time for harvest approaches in many countries (Middleton *et al.*, 2004; Hoi, 2010), which leaves residue on the fruit that cannot be reduced to levels suitable for human consumption (Neufeld *et al.*, 2010). The adverse effects of these harmful pesticides on the health of consumers are considered to be high (Brink *et al.*, 2003).

It is likely that integrated pest management (IPM) strategies can achieve a sufficient level of pest control to be acceptable to vegetable producers. Various IPM strategies have been developed and are being promoted among farmers in the tropics. This chapter aims to compile simple, affordable, and environmentally sound IPM strategies, developed mostly by AVRDC – The World Vegetable Center and/or its collaborators, for the management of key pests in selected tropical vegetable crops.

8.2 Eggplant, *Solanum melongena* L. (Solanales: Solanaceae)

The major pests of eggplant include the fruit and shoot borer (*L. orbonalis*), leafhopper [*Amrasca devastans* Distant (Hemiptera: Cicadellidae)], whitefly

(*B. tabaci*), thrips [*Thrips palmi* Karny (Thysanoptera: Thripidae)] and aphid [*Aphis gossypii* Glover (Hemiptera: Aphididae)].

The sucking pests such as whitefly, aphid, thrips and leafhopper first appear on the crop either in the seedling nursery or in the main field soon after transplanting. Depending on management practices, they may be present in the crop throughout the cropping season. Both the adults and nymphs of *B. tabaci* suck the plant sap and reduce the vigor of the plant. In severe infestations, the leaves turn yellow and drop off. Whiteflies secrete large quantities of honeydew, which favors the growth of sooty mould on leaf surfaces and reduces the photosynthetic efficiency of the plants. *A. gossypii* occurs in large numbers on the tender shoots and lower leaf surfaces, and sucks the plant sap. Slightly infested leaves exhibit yellowing. Severe aphid infestations cause young leaves to curl and become deformed. Like whitefly, aphids also produce honeydew, which leads to the development of sooty mould.

Both nymphs and adults of *A. devastans* suck the sap from the leaves, which leads to yellowing, bronzing and drying, or “hopper burn”. *T. palmi* prefers to feed mostly on foliage, but occasionally on fruit. Slightly infested leaves exhibit silvery feeding scars on the lower leaf surfaces, especially along the mid-rib and veins. In severe infestations, the leaves turn yellow or brown and dry on the lower leaf surfaces. Infested fruit is scarred and deformed.

During the early vegetative stage of the crop, the *L. orbonalis* larva feeds inside the tender shoots, which results in wilting of the young shoots, followed by drying and drop-off. The pest prefers to feed on the fruit during the fruiting stage of the crop; damaged fruit also exhibits boreholes on the surface that often are sealed with excreta, rendering the fruit unfit for marketing and consumption.

Integrated Pest Management

- 8.2.1. Crop rotation with non-solanaceous crops is an effective way to reduce the population build-up of *L. orbonalis*. However, careful consideration in choosing the crops for rotation is imperative, because the crops selected (*e.g.*, cotton or okra) may also support other common pests such as whitefly, leafhopper and aphid.
- 8.2.2. Planting resistant or moderately resistant cultivars can reduce pest damage and thus allow greater yields. For example, accessions or varieties such as EG058, Pusa Purple Long, Pusa Purple Cluster, Pusa Purple Round, etc. have been reported to be tolerant or resistant to *L. orbonalis* (Parker *et al.*, 1995; Alam *et al.*, 2003; Shivalingaswamy and Satpathy, 2007). Except for EG058, which is an AVRDC accession,

most of the varieties noted originated in India. Indian cultivars such as 'Manjari Gota,' 'Vaishali,' 'Mukta Kesi,' 'Round Green,' and 'Kalyanipur T3' and the Bangladeshi variety, Bagun 6, are reported to be less susceptible or tolerant to *A. devastans* damage (Parker *et al.*, 1995; Rashid *et al.*, 2003). Three cultivated eggplant accessions (S130, S145 and S158) are reported to be less susceptible to *A. devastans* and *A. gossypii* (AVRDC, 1999). However, no reported resistant or tolerant eggplant varieties are available for *B. tabaci* and *T. palmi*.

- 8.2.3. Eggplant seedlings should not be grown near fields with previous or existing crops, or near dried eggplant heaps (Alam *et al.*, 2003). If seedlings must be grown in such areas, the seedling beds should be covered with 30-mesh nylon net to prevent the entry of *L. orbonalis* moths and thus egg laying. If *B. tabaci* and *T. palmi* are common in the region, use of 50–64 mesh nylon net to cover the seedling beds is useful preventive measure. If non insect-proof nets are used, they should be treated with insecticides or neem oil to prevent the entry of sucking insects into the protective structures (Cerkauskas, 2004).
- 8.2.4. Yellow (565-590 nm) sticky traps can be used to monitor and/or mass-trap *B. tabaci* and *A. gossypii* in the field (Durairaj *et al.*, 2007). Similarly, blue (H⁷460-475 nm) sticky traps can be used to monitor and/or mass-trap *T. palmi* (Chen *et al.*, 2004; Naojee *et al.*, 2010).
- 8.2.5. Prompt removal and destruction of infested shoots and fruit at regular intervals until final harvest throughout a community substantially reduces *L. orbonalis* damage (Alam *et al.*, 2003). Pruning does not adversely affect the plant growth or yield (Srinivasan and Huang, 2009).
- 8.2.6. *L. orbonalis* sex pheromone lures in traps can be installed at the rate of 100 traps per hectare (Cork *et al.*, 2003) to reduce fruit damage and increase yield. Traps should be placed either at canopy level or at slightly above the canopy level for effective attraction (Alam *et al.*, 2003).
- 8.2.7. Generalist predators such as ladybird beetles and green lacewings are highly efficient in preying on sucking pests. Parasitoids such as *Anagrus flaveolus* Waterhouse and *Stethynium triclavatum* Enock are effective against *A. devastans* (Subba Rao 1968; Parker *et al.*, 1995). Parasitoids such as *Trathala flavoorbitalis* (Cameron), *Eriborus sinicus* Holmgren, and *Pristomerus testaceus* Morley are commonly found occurring on *L. orbonalis* larvae (AVRDC, 1996a;

Alam *et al.*, 2003). In addition, weekly releases of the egg parasitoid *Trichogramma chilonis* Ishii @ 1 g parasitized eggs/ha/week, and larval parasitoid *Bracon hebetor* Say @ 800-1000 adults/ha/week are effective in reducing *L. orbonalis* (Alam *et al.*, 2006). Broad-spectrum pesticides should not be used, as they may harm these and other natural enemies.

- 8.2.8. Microbial pesticides and botanical pesticides can be used to manage *L. orbonalis* and other sucking pests (Rao *et al.*, 1999; Islam *et al.*, 2011; Mathur *et al.*, 2012). Need based application of appropriately selective pesticides, when recommended by the local extension service, is also advisable.

8.3 Tomato, *Solanum lycopersicum* L. (Solanales: Solanaceae)

Several kinds of insect and mite pests attack tomato leaves, flower buds, and fruit during tomato cultivation. The most commonly occurring pests on tomato are fruit worm (*H. armigera*), common armyworm [*Spodoptera litura* Fabricius (Lepidoptera: Noctuidae)], whitefly (*B. tabaci*), leaf miner [*Liriomyza* spp. (Diptera: Agromyzidae)] and spider mites [*Tetranychus urticae* Koch., *T. cinnabarinus* (Boisd.) and *T. evansi* Baker & Pritchard (Acarina: Tetranychidae)]. Whitefly and fruit worm are the major insect pests in most of the tropical tomato-producing regions of the world, as the whitefly transmits leaf-curl virus disease and fruit worm causes severe damage to the fruit, reducing the marketable yield of tomato.

Liriomyza spp. will appear on the crop in the seedling stage as well as in the main field soon after transplanting. Their larval feeding causes irregular mines on leaf surfaces. In severe infestations, several mines are formed on the same leaf, which will drastically reduce photosynthesis and thus reduce yield. Sometimes, death of the whole plant occurs. Although *B. tabaci* feeding can cause direct damage to the growing plants, it also acts as a vector for tomato yellow leaf-curl virus (TYLCV) disease. Plants infected by TYLCV show stunted growth with erect shoots. Leaflets curl upwards and inwards, are reduced in size, are stiff, thicker than normal, and leathery in texture. Leaves have yellowing along the margins and have interveinal chlorosis (yellowing). The flowers wither and droop, and fruit set is reduced or may be nil. Fruit, if produced, is small and unmarketable.

The neonate larvae of *S. litura* feed on leaf surfaces. Mature larvae feed on the whole leaves, leaving only the main veins. Rarely, they feed on the immature growth stages of tomato fruit. However, *S. litura* does not bore into fruit like *H. armigera*. Sometimes, the larvae may cut the seedlings or young plants at soil level. *H. armigera* is a polyphagous and highly mobile insect. The

older larvae prefer to feed on flowers and young fruits. The larvae make holes and then feed by thrusting their heads inside. The holes are circular and often surrounded by fecal pellets.

Spider mites have emerged recently as serious pests of tomato in South Asia, Southeast Asia, and Africa. Low relative humidity favors their multiplication and hence they pose a serious production threat during the dry season. Spider mite feeding leads to the formation of several white or yellow speckles on the leaves. In severe infestations, leaves will completely desiccate and drop-off. The mites also produce webbing on the leaf surfaces; under severe conditions, the whole plant is wrapped in webs. Under high population densities, the mites move to the tip of the leaf or top of the plant and congregate to form a ball-like mass, which will be carried by the wind to newer leaves or plants.

Integrated Pest Management

- 8.3.1. Crop rotation with non-host crops is effective in managing tomato pests. However, selection of an appropriate non-host crop is challenging, because most of the tomato pests are highly polyphagous and feed on several agricultural and horticultural crops as well as weed plants.
- 8.3.2. Although using resistant tomato cultivars can reduce pest damage, commercial cultivars with appreciable levels of resistance to major tomato pests are not yet available. Germplasm screening at AVRDC – The World Vegetable Center revealed the presence of high levels of *H. armigera* resistance only in wild *Solanum* species, particularly *S. habrochaites* and *S. pennellii*. Efforts to introgress resistance from wild species into cultivated tomato resulted in small-fruited resistant accessions (Talekar *et al.*, 2006). Although resistant sources, mostly in wild tomatoes, are available for *B. tabaci* (Fancelli and Vendramim, 2002; Toscano *et al.*, 2002; Muigai *et al.*, 2003; Srinivasan and Uthamasamy, 2004; Baldin *et al.*, 2005; Firdaus *et al.*, 2012; 2013), cultivation of commercially available TYLCV resistant tomato varieties is currently the suggested practice. For instance, varieties from Southern India such as ‘Sankranthi,’ ‘Nandi,’ and ‘Vybhav’ have been reported to be effectively resistant (Muniyappa *et al.*, 2002).
- 8.3.3. Healthy seedling production as explained in the previous eggplant section might also prevent or reduce the incidences of early season sucking pests, including whitefly.
- 8.3.4. African marigold (*Tagetes erecta* L.) can be planted as a trap crop to reduce the incidence of *H. armigera* (Srinivasan *et al.*, 1994). It is

important to synchronize transplanting of both crops so that flowering coincides, which attracts *H. armigera* female adults. Tropical soda apple (*Solanum viarum* Dunal) also can be used as an effective trap crop to manage *H. armigera* (AVRDC, 2000; 2001; Srinivasan *et al.*, 2013).

- 8.3.5. Yellow is the most attractive color to the adults of *Liriomyza* sp. (Parrella, 1987). Hence, yellow sticky traps can be used to monitor and/or mass-trap the adults of *Liriomyza* spp., besides *B. tabaci*.
- 8.3.6. Sex pheromone traps can be used to monitor, mass-trap, or disrupt the mating of male moths of *H. armigera* and *S. litura*. Although sex pheromone traps baited with pheromone lures can be used to trap more males, this method is less effective for polyphagous insects like *H. armigera* and *S. litura*. Polyphagous populations are always higher due to the availability of multiple host plants in the tropics. Two population parameters viz., density and polymorphism, are important factors that affect the trap catches of males (Kumar and Shivakumara, 2003). Placing a high concentration of sex pheromone (all or one component of the multi-component pheromone) in a slow-release formulation on a 5- and 10-m grid in the field results in a drastic reduction in male moths, which adversely affects mating in *H. armigera* (AVRDC, 1988). Significant trapping of hundreds of male moths has also been achieved in *S. litura* with sex pheromone lures (Dharma Putra *et al.*, 2013; Srinivasan *et al.*, 2013). Hence, mass-trapping is possible when the traps are baited with effective pheromone lures/formulations.
- 8.3.7. Egg parasitoids (*Trichogramma pretiosum* Riley and *T. chilonis*) and larval parasitoids (*Campoletis chlorideae* Uchida) can be conserved and/or released in tomato fields at regular intervals to check the build-up of *H. armigera* and *S. litura* (Ballal and Singh, 2003; Gupta *et al.*, 2004; Krishnamoorthy, 2012). In addition, commercially available biopesticides based on *Bacillus thuringiensis* Berliner, *Helicoverpa armigera* nucleopolyhedrovirus (HaNPV), *Spodoptera litura* nucleopolyhedrovirus (SINPV), *Beauveria bassiana* (Bals.-Criv.) Vuill. and neem (*Azadirachta indica* A. Juss.) can be used either alone or in combinations against *H. armigera* and *S. litura* (Nathan and Kalaivani, 2006; Mohan *et al.*, 2007; Senthilkumar *et al.*, 2008). Application of *B. bassiana* in the soil has substantially contributed to the mortality of *H. armigera* pupae (AVRDC, 1992). Bio-pesticide formulations based on *B. thuringiensis* subsp. *aizawai*

and *B. thuringiensis* subsp. *kurstaki* are highly toxic to the larvae of *H. armigera* (AVRDC, 1995).

- 8.3.8. Use of broad-spectrum chemical pesticides should be avoided, as these pesticides harm the parasitoid and predator complex of *Liriomyza* spp. and *Tetranychus* spp. Leaf miners have several parasitoids. For instance, *Gronotoma micromorpha* Perkins (larval-pupal parasitoid), *Chrysocharis pentheus* Walker, *Neochrysocharis formosa* (Westwood) and *Diglyphus isaea* Walker (larval parasitoids) and *Halticoptera circulus* Walker and *Opius phaseoli* Fischer (pupal parasitoids) are known to occur in Asia (Lee *et al.*, 1990; Sivapragasam and Syed, 1999; Niranjana *et al.*, 2005; Abe, 2006;), and help to keep the *Liriomyza* spp. population in check. Similarly, several predators of *Tetranychus* spp. occur in most countries. For instance, *Stethorus* spp., *Oligota* spp., *Anthrocnodax occidentalis* Felt, *Feltiella minuta* Felt, etc. are known to occur in Taiwan (Ho, 2000). A Brazilian strain of the predatory mite, *Phytoseiulus longipes* Evans, and the pathogenic fungus, *Neozygites floridana* (Weiser & Muma), have shown promising results against *T. evansi* (Furtado *et al.*, 2007; Wekesa *et al.*, 2007). Oil-based formulations of entomopathogenic fungi such as *B. bassiana* isolate GPK and *Metarhizium anisopliae* (Metchnikoff) Sorokin isolate ICIPE78 also reduced the population density of *T. evansi* substantially (Wekesa *et al.*, 2005). The predatory mite and biopesticides can be used to manage *Tetranychus* spp.
- 8.3.9. Chemical pesticides are widely used against tomato pests. Pesticide spraying should be scheduled soon after noticing the eggs or during the early larval stages of *H. armigera* and *S. litura*. Selective and systemic pesticides can be used for the sucking insects. It is advisable to follow a proper pesticide rotation. Before application, it is important to check the effectiveness of chemical pesticides in the region and their registration status for tomato.

8.4. Peppers, *Capsicum* spp. (Solanales: Solanaceae)

The major pests on peppers are thrips [*T. palmi*, *Scirtothrips dorsalis* Hood, and *T. parvispinus* Karny (Thysanoptera: Thripidae)], aphids [*A. gossypii* and *Myzus persicae* Sulzer (Hemiptera: Aphididae)], fruit borer (*H. armigera* and *S. litura*), whitefly (*B. tabaci*) and broad mite [*Polyphagotarsonemus latus* (Banks) (Acarina: Tarsonemidae)].

Thrips feed mostly on foliage, especially on the undersides. Slightly infested leaves exhibit silvery to brown feeding scars, especially along the mid-rib and

veins. In severe infestations, the leaves curl upward like the shell of a boat, and are reduced in size. Infested fruit are scarred. Thrips also transmit *Tomato spotted wilt virus* and *Capsicum chlorosis virus* (Adkins *et al.*, 2010). Aphids feed on the tender shoots and lower leaf surfaces. The infested leaves are distorted, stunted and curled. The upper leaf surfaces develop sooty mould due to the production of honeydew. Aphids transmit *Chilli veinal mottle virus* and *Cucumber mosaic virus*. Broad mite feeding is usually confined to the undersides of leaves, where areas between veins are brownish and dried out and brittle in severe cases. Young leaves are curled downward and narrower than normal (Parker *et al.*, 1995). Damage by *H. armigera* and *S. litura* during flowering and fruit formation can lead to 90% flower and fruit drop (Reddy and Reddy, 1999).

Integrated Pest Management

- 8.4.1. Healthy seedling production practices should be followed to reduce the incidence of early season sucking pests.
- 8.4.2. Selected *Capsicum annum* L. and *C. baccatum* L. accessions such as PBC018, PBC030, PBC081, PBC151, PBC272, PBC681, PBC726, PBC732, PBC785 and PBC880 were reported to be moderately resistant or resistant to *A. gossypii* (AVRDC, 1993a; 1998). Accessions such as PBC725, PBC901 and PBC1398 were reported to be resistant to *P. latus* (AVRDC, 1998). Accessions such as C00069 and PBC145 were found to be resistant to both *M. persicae* and *T. palmi*. Cultivation of these or similar locally available resistant accessions could substantially reduce incidences of thrips, aphids and broad mite in the tropics.
- 8.4.3. Castor (*Ricinis communis* L.) can be planted as a trap crop to divert the incidence of *S. litura* on peppers (Reddy and Rosaiah. 1987).
- 8.4.4. Yellow sticky traps can be used to monitor and/or mass-trap the adults of *B. tabaci*. Blue sticky traps and kairomonal cues baited traps can be used to trap thrips (Nielsen *et al.*, 2010). A semiochemical-baited autoinoculation device treated with *M. anisopliae* could also be adapted to manage thrips on peppers (Niassy *et al.*, 2012).
- 8.4.5. Sex pheromone traps, biopesticides and parasitoids as described in the tomato section can also be used to manage *H. armigera* and *S. litura* larvae.
- 8.4.6. Chemical pesticides are widely used against pepper pests. Pesticide spraying should be scheduled soon after noticing eggs or during the

early larval stages of *H. armigera* and *S. litura*. A new group of insecticides, the diamides, offer potential control of *H. armigera* and *S. litura* (Tatagar *et al.*, 2009). However, it is advisable to follow a proper pesticide rotation to avoid the development of resistance. A combination of organic compounds containing natural alkaloids, lactone and isoflavones, seaweed extract and chavicol, and neem with avermectins can substantially reduce the leaf curl caused by thrips and broad mite (Mondal and Mondal, 2012).

8.5. Vegetable Brassicas, *Brassica* spp. (Brassicales: Brassicaceae)

Diamondback moth, *P. xylostella*, is a cosmopolitan and destructive pest of vegetable brassicas. The larvae feed and create holes in the leaves; however, severe damage is caused when the larvae tunnel into the heads of crops like cabbage, sometimes causing almost 100% crop loss. Cabbage head caterpillar, *Crocidolomia binotalis* (Lepidoptera: Pyralidae) is another serious pest of brassicas. The early instar larvae migrate toward the growing center of vegetable brassicas, where they conceal themselves in webbing. Subsequent feeding can damage the plant's apical meristem and thus destroy the entire crop (Smyth *et al.*, 2003). Cabbage webworm, *Hellula undalis* (Lepidoptera: Pyralidae) is also a major brassica pest in lowland production systems (Sivapragasam and Chua, 1997), especially in the summer season (AVRDC, 1978). It prefers to feed on the young terminal bud and the unopened leaves (Sivapragasam and Chua, 1997), leading to yield losses of up to 100% in Asia, Africa and the Pacific (Kalbfleisch, 2006). Imported cabbage worm, *Pieris rapae* L. (Lepidoptera: Pieridae) occasionally causes serious damage to vegetable brassicas. The larvae bore directly into the growing heads or cause irregular feeding holes on the leaves. Aphids [*M. persicae*, *Lipaphis erysimi* (Kaltenbach) and *Brevicoryne brassicae* L. (Hemiptera: Aphididae)] also cause significant damage to brassica plants in all growth stages. Aphid infestation leads to leaf curling and wrinkling. With severe infestation, the leaves wilt and the entire plant dies (Parker *et al.*, 1995). The striped flea beetle, *Phyllotreta striolata* Fab. (Coleoptera: Chrysomelidae) causes serious damage to leafy brassicas and radish in Southeast Asia. Infested young plants have numerous small holes in the cotyledons and leaves.

Integrated Pest Management

- 8.5.1. Growing resistant accessions can avert pest damage. Four non-heading Chinese cabbage accessions (B464, B487, B488 and B490) were found to be tolerant to aphids (AVRDC, 1977). Five Chinese cabbage accessions (B159, B186, B197, B488 and B501) were least susceptible

to *H. undalis* (AVRDC, 1978; 1987). Two *Brassica campestris* ssp. *chinensis* var. *parachinensis* accessions (B095 and B583) that are compatible with Chinese cabbage for breeding, were least susceptible to *P. xylostella* (AVRDC, 1981). Cultivation of these, or similar locally available resistant accessions, could substantially reduce the pest incidences on vegetable brassicas in the tropics.

- 8.5.2. Brassica seedling production can be carried out under 32-mesh nylon net to reduce *P. xylostella* infestation early in the season. Growers can thus reduce the amount of chemical pesticides used at the beginning of the crop cycle, which will help to protect natural enemies. In peri-urban areas where the brassicas are grown on smaller farms, plots can be confined within 32-mesh nylon net barriers on all four sides using a 2-2.5 m high net barrier to manage *P. xylostella* and *C. binotalis* (AVRDC, 1999; 2000; 2001).
- 8.5.3. Clean cultivation is recommended for *H. undalis* because it feeds and develops on certain weeds such as *Cleome rutidosperma* and *C. viscosa*, which act as alternate hosts during the off-season in South and Southeast Asia (Sivapragasam and Chua, 1997; Kalbfleisch, 2006).
- 8.5.4. Sequential trap cropping with Indian mustard (*Brassica juncea* (L.) Czern.), at the rate of two rows of mustard between every 25 rows of brassica, is an effective way of reducing *P. xylostella* and *C. binotalis* damage. One row of mustard should be sown 15 days before brassica planting, and a second sowing is recommended on the adjacent ridge on the 25th day after planting brassica (Srinivasan and Krishnamoorthy, 1992). The trap crop should be sprayed with chemical pesticides. Yellow rocket, *Barbarea vulgaris* var. *arcuata*, can be used as a dead-end trap crop for *P. xylostella* (Shelton and Badenes-Pérez, 2006). Yellow rocket is highly attractive to *P. xylostella*, but its offspring cannot survive on it. Dead-end trap crops do not require any pesticide treatment to prevent pest populations from moving onto the main crop.
- 8.5.5. Intercropping of common cabbage with safflower or garlic can substantially reduce the infestation of *P. xylostella* (AVRDC, 1987).
- 8.5.6. Traps baited with female sex pheromone lures can be placed at the rate of 50 traps per hectare randomly in the field, and replaced twice during the season (AVRDC, 1991) to manage *P. xylostella*. A recent study has shown that sex pheromone traps installed at the rate of 30-45 traps/ha were effective for controlling *P. xylostella* in the field, and reduced insecticide use by 30% per planting season in

southwestern China (Zhao *et al.*, 2011). It must be noted that different *P. xylostella* populations may react differently to the same sex pheromone lure (Feng *et al.*, 2011). Hence, the pheromone lures should be validated in a new region before using them as a pest management tool. Traps baited with a 10:1 mixture of (Z)-11-hexadecenyl acetate and (Z)-9-tetradecenyl acetate attracted substantially high numbers of male *C. binotalis* moths in vegetable brassica fields in Indonesia (Adati *et al.*, 2007). Wing traps baited with female sex pheromone (E,E-11,13-hexadecadienal @ 10 µg per trap) can be placed at a height of 0.5 m above ground at a distance of 15 m, and replaced every two to six weeks depending on the trap catches to manage *H. undalis* (Kalbfleisch, 2006).

- 8.5.7. Traps baited with aggregation pheromones plus host plant volatiles could trap substantially large numbers of *P. striolata* beetles (Beran *et al.*, 2011; Srinivasan, 2012).
- 8.5.8. Release of a larval parasitoid (*Diadegma semiclausum* Hellen) and the pupal parasitoid [*Diadromus collaris* (Gravenhorst)] substantially reduced damage from *P. xylostella* in highland areas. The larval parasitoid *Cotesia plutellae* Kurdjumov is well-adapted to manage *P. xylostella* in lowland areas. Unlike the egg parasitoid, augmentative releases for the larval and pupal parasitoids at regular intervals are not necessary. These larval and pupal parasitoids already have been introduced into several countries in Asia (AVRDC, 1993b; 1996b) and Africa (Nyambo and Lohr, 2005). A generalist egg parasitoid, *T. chilonis*, was found to efficiently parasitize egg masses of *C. binotalis* in Samoa (Uelese *et al.*, 2011). Although the polyphagous pentatomid bug, *Eocanthecona furcellata* Wolff, predate more than 70% of the third instar *C. binotalis* larvae, the searching ability in the field decreases (AVRDC, 1996a). However, there is no species-specific efficient parasitoid available for *C. binotalis* hitherto. Release of an egg parasitoid, *Trichogrammatoidea bactrae* Nagaraja once a week for five weeks starting within a week after transplanting brassicas is recommended to manage *H. undalis* (Sivapragasam, 1996). In addition, a few larval parasitoids such as *Bassus* sp., *T. flavoorbitalis*, *Chelonus* sp. and *Phanerotoma* sp. have been reported to infest *H. undalis* in Malaysia. *Apanteles glomeratus* L. is the widely occurring larval parasitoid against *P. rapae*.
- 8.5.9. *P. xylostella* and *C. binotalis* are susceptible to *B. thuringiensis* toxins, for example, Cry1Ac, Cry1Ab, Cry1Aa, Cry1Ba and Cry1Ca (Srinivasan and Hsu, 2008; Shelton *et al.*, 2009). Hence,

B. thuringiensis formulations based on the most sensitive Bt toxins would offer effective control (Ooi, 1980; Sastrosiswojo and Setiawati, 1992; Malathi and Sriramulu, 2000). As *B. thuringiensis* biopesticides are not harmful to the parasitoids, both these components synergistically reduce *P. xylostella* and *C. binotalis* damage. Strains of entomopathogenic fungi such as *M. anisopliae* were also effective against *P. xylostella* (AVRDC, 1999). Improved formulations of *B. bassiana* showed promising prolonged impact in suppressing *P. xylostella* populations in field conditions (Ghosh *et al.*, 2011). *H. undalis* is highly sensitive to Cry1Ba and Cry1Ca, but less susceptible to Cry1A toxins (Srinivasan and Hsu, 2008; Shelton *et al.*, 2009). This is also confirmed by its susceptibility to *B. thuringiensis* subsp. *aizawai* based formulations (Srinivasan and Hsu, 2008), but not for the *B. thuringiensis* subsp. *kurstaki* based formulations (AVRDC, 1987), although the latter was shown to be effective in the Philippines (Ulrichs and Mewis, 2003) and India (Singh *et al.*, 2000). Hence, after local validation, *B. thuringiensis* formulations can be used to manage *H. undalis*. Similarly, *P. rapae* is also sensitive to Cry1Ba and Cry1Ca toxins (Shelton *et al.*, 2009), and most formulations of *B. thuringiensis* are effective against this pest. A granulosis virus specifically infecting *P. rapae* is also found in Taiwan (Parker *et al.*, 1995).

- 8.5.10. Chemical pesticides still occupy the predominant role in managing *P. xylostella*. Due to extensive and inappropriate pesticide use, it has developed resistance to almost every class of insecticide that has been used against it, including novel insecticides such as chlorantraniliprole and flubendiamide (Sukonthabhirom *et al.*, 2011) and biopesticides such as *B. thuringiensis* (Tabashnik *et al.*, 1990). In addition, use of broad-spectrum chemical pesticides early in the season often disrupts the control exerted by parasitoids (Talekar and Shelton, 1993). The rotation and windows approach for pesticide application can effectively manage *P. xylostella* on vegetable brassicas (Baker, 2011; Ridland and Endersby, 2011). Proper selection and judicious use of chemical pesticides targeting *P. xylostella* should also control *C. binotalis* and *H. undalis* on vegetable brassicas; *C. binotalis* is found to be susceptible to newer pesticides such as emamectin benzoate, spinosad and indoxacarb (Kannan *et al.*, 2011). An improved mode of pesticide application via brassica shoots is suggested for *H. undalis* (Sivapragasam, 1996) as this method minimizes the adverse effects of chemical pesticides on the natural enemies present in brassica production systems.

8.6 Cucurbits: *Cucurbita* spp., *Citrullus* spp., *Cucumis* spp., *Momordica* spp. (Cucurbitales: Cucurbitaceae)

Melon fly, *Bactrocera cucurbitae* (Diptera: Tephritidae) has been recorded as a predominant pest on several host plants in the Cucurbitaceae. The extent of losses varies between 30 to 100%, depending on the plant species and the season (Dhillon *et al.*, 2005). Maggots feed inside the fruit, leading to distortion or rotting. The pumpkin caterpillar, *Diaphania indica* Saund. (Lepidoptera: Pyralidae) also causes major yield losses in several cucurbits. It widely occurs in most tropical countries. The young larvae fold the leaves together and feed on the foliage. They also feed on the young fruits, especially on the fruit skin. Aphids (*M. persicae* and *A. gossypii*) also cause significant damage to cucurbits as a direct pest, as well as vectors of viruses such as *Zucchini yellow mosaic virus*, *Watermelon mosaic virus*, *Cucumber mosaic virus* and *Cucurbit aphid-borne yellows virus* (Biswas and Varma, 2006; Bunwaree, 2001; Dahal *et al.*, 1997; Haque *et al.*, 2004; Shaifullah *et al.*, 2005; van der Meer and Garnett, 1987; Wakman *et al.*, 2002). The viruses can cause yield losses of up to 100% (Dahal *et al.*, 1997; Gaba *et al.*, 2004; Provvidenti, 1996). The whitefly (*B. tabaci*) transmits *Squash leaf curl virus* in cucurbits. Symptoms include leaf curling, mottling and severe stunting of the whole plant, and fruit may be small and distorted (Duffus and Stenger, 1998). Up to 100% disease incidence (with 100% yield losses) has been reported (Bananej *et al.*, 2002; Isakeit and Robertson, 1994).

Integrated Pest Management

- 8.6.1. Sources of resistance to melon fly are available in pumpkin, bitter gourd, bottle gourd, sponge gourd, ridge gourd and round melon (Dhillon *et al.*, 2005; Gogi *et al.*, 2010). However, the resistance may vary across locations. For instance, bitter gourd varieties such as Faisalabad-long and Col-II, reported as the most resistant in Pakistan (Gogi *et al.*, 2010) were highly susceptible to melon fly in Taiwan. Resistance and/or tolerance to *A. gossypii* in *Cucumis melo* L. germplasm has been reported (Bohn *et al.*, 1996; Dhillon *et al.*, 2007; Martin *et al.*, 1996). Resistance in melon accessions to *A. gossypii* has also decreased viral transmission (Lecoq *et al.*, 1979). Aphid resistance has been introduced into several commercial melon cultivars (Lecoq *et al.*, 1998). Three Indian *C. melo* accessions (PI414723, PI164723 and 90625) and a Korean accession (PI161375) had field resistance to *B. tabaci* in some Caribbean islands (Boissot *et al.*, 2003). Thus availability and cultivation of similar locally available resistant accessions could substantially reduce pest incidences on cucurbits.

- 8.6.2. Healthy seedling production as described in the eggplant section might also prevent or reduce the incidence of early season sucking pests including aphids and whitefly, and also reduce the transmission of viruses in susceptible varieties.
- 8.6.3. If field sanitation (collection and destruction or burying of infested fruit 15–45 cm deep in the soil) is adopted throughout a community, it will substantially reduce the melon fly population build-up (Klungness *et al.*, 2005; Kumar *et al.*, 2011). Bagging the fruits soon after formation prevents melon fly damage substantially (Akhtaruzzaman *et al.*, 1999; Kumar *et al.*, 2011).
- 8.6.4. Yellow sticky traps and biopesticides can be used to manage whitefly and aphids.
- 8.6.5. Kairomonal attractant cue-lure [4-(p-acetoxyphenyl)-2-butanone] traps are highly effective for monitoring and mass-trapping of *B. cucurbitae* in bitter melon and other crops (Pawar *et al.*, 1991; Vargas *et al.*, 2000). Similarly, protein baits are highly attractive to female melon flies (Kumar *et al.*, 2011). Sex pheromone lures are effective against *D. indica* moths (Wakamura *et al.*, 1998) and they are now commercially available.
- 8.6.6. An egg – pupal parasitoid, *Fopius arisanus* (Sonan) is believed to have high potential in reducing melon fly damage and it has been introduced into Hawaii and Mexico (Bautista *et al.* 2004; Zenil *et al.*, 2004). *Apanteles taragamae* Viereck, *T. flavoorbitalis* and *Elasmus indicus* Rohwer were found to parasitize *D. indica*, although they are not species-specific parasitoids (Ganehiarachchi, 1997; Muniappan *et al.*, 2012).
- 8.6.7. Soil application of *M. anisopliae* formulations has reduced the emergence of fruit flies (Ekesi *et al.*, 2005). This strategy can be easily adapted for melon fly. *B. thuringiensis* is effective against *D. indica* (Schreiner, 1991).
- 8.6.8. Since application of chemical pesticides against melon fly may not be effective, attractants can be combined with small quantities of pesticides to develop an ‘attract and kill’ system. For example, protein bait sprays mixed with chemical pesticides can be spot-sprayed on the roosting host plants of melon fly. Crops like maize, cassava, sorghum and castor should be planted around the main crop (bitter melon, water melon, etc). These roosting host plants can be spot-sprayed with protein bait sprays once a week (or more often during the wet season) to kill the newly emerged flies before they mature (Vargas *et al.*, 2008; Mcquate, 2011). Since both melon fly and *D. indica* can develop insecticide resistance rapidly (Chen, 2002), chemical pesticides should be selected and applying with care.

8.7 Okra, *Abelmoschus* spp. (Malvales: Malvaceae)

Cotton aphid (*A. gossypii*) and whitefly (*B. tabaci*) are the major pests of okra in its early growth stages. In addition to causing direct feeding damage, *B. tabaci* also transmits yellow vein mosaic virus disease. Early symptoms include alternate green and yellow patches, vein clearing, and vein chlorosis of leaves. The yellow network of veins becomes very conspicuous with thickened veins and veinlets in the later stages. Cotton leafhopper (*A. devastans*) is another major pest of okra. Feeding by the nymphs and adults leads to hopper burn symptoms. Flea beetles [*Podagrica uniformis* (Jac.) and *P. sjostedti* (Jac.) (Coleoptera: Chrysomelidae)] are two serious pests of okra in Africa. Their infestation commences from germination through all stages of plant growth. Although they are mainly leaf eaters (Fasunwon and Banjo, 2010), they also are vectors for *Okra mosaic virus* (Alegbejo *et al.*, 2008). Pod borers [*H. armigera*, *Earias vitella* Fab., *E. insulana* Boisd., *E. biplaga* Walker and *E. cupreoviridis* Walker (Lepidoptera: Noctuidae)] are the major pests of okra during the vegetative as well as fruiting stage. *Earias* spp. bore into the terminal shoots of young plants causing death of the growing tips and leading to the development of lateral branches. During the reproductive stage, the larvae bore and feed into the flower buds, flowers and pods (Varela and Seif, 2004).

Integrated Pest Management

- 8.7.1. Although sources of resistance to whitefly are not available, varieties that are resistant or moderately resistant to yellow vein mosaic virus disease are available (Shetty *et al.*, 2013). Four *A. gossypii* resistant okra accessions (VI033805, VI036213, VI051114 and VI033824) and two leafhopper resistant okra accessions (VI041230 and VI033809) have been recently identified by AVRDC. Thus availability and cultivation of similar locally available resistant accessions could substantially reduce pest incidences on okra. Other control methods mentioned in previous sections for aphid, whitefly and leafhopper can be adapted for okra.
- 8.7.2. Sex pheromone traps can be used to monitor the population of *Earias* spp. (Hall *et al.*, 1980; Cork *et al.*, 1985; 1988; Hamed and Nadeem, 2010).
- 8.7.3. Biopesticides including *B. thuringiensis* and neem, and novel pesticides such as spinosad are highly effective in reducing the damage caused by *Earias* spp. (Mirmoayedi *et al.*, 2010; Tripathi and Maurya, 2011).
- 8.7.4. Insecticide applications may be necessary if large flea beetle populations are present at the early stage of crop growth. Foliar

insecticides are recommended for quick control of large populations attacking vulnerable seedlings (Varela and Seif, 2004). Since development of insecticide resistance has been reported in *Earias* spp. (Satpute *et al.*, 2003), proper pesticide rotation and windows should be followed to minimize the risk.

8.8. Alliums, *Allium* spp. (Asparagales: Amaryllidaceae)

Onions and garlic are the two most important alliums in the developing world. Onion thrips, *T. tabaci*, is the key pests in most onion production regions of the world. Immature and adult thrips feed on onion foliage by rasping and sucking the juice. The removal of chlorophyll leads to the development of white to silvery patches or streaks, and eventually the damaged leaf area dries up. Thrips feeding can reduce bulb yields by 30-50% and losses can be substantially high if thrips transmit *Iris yellow spot virus*, or create damage that permits other pathogens to infect the crop (Nault *et al.*, 2012). Beet armyworm (*Spodoptera exigua* Hubner) and common armyworm (*S. litura*) are serious defoliators of onions and shallot in Asia. The larvae either feed on the leaf surface or in most cases bore holes in the leaf cuticle and feed from inside the tubular leaves.

Integrated Pest Management

- 8.8.1. *Allium cepa* L. and *A. fistulosum* L. accessions (AC 570, AF 204, AF 218 and TA 246) are resistant to *S. exigua* (AVRDC, 1997; 1998). Five other onion accessions (AC 521, AC 525, AC 584, TA 189 and TA 243) are resistant to thrips (AVRDC, 1998). Cultivation of similar locally available resistant accessions could substantially reduce pest incidence on onions.
- 8.8.2. Growing castor as a trap crop on the borders of onion plots reduces the infestation of *S. litura* on onions. Castor should be sown one month before transplanting onions (AVRDC, 1996a). Intercropping of onion with subterranean clover substantially reduces thrips damage. Onions should be planted 30 days after sowing the subterranean clover (AVRDC, 1997).
- 8.8.3. Onion thrips prefer older plants and infestation is critical during the bulb enlargement stage. Keeping soil moist during this stage helps to reduce onion thrips damage. More irrigation combined with straw mulching reduces thrips damage in onions (AVRDC, 1999).
- 8.8.4. Sex pheromone traps can be used to monitor and mass-trap *S. litura* and *S. exigua* adults. Installing pheromone traps baited with only one [(Z, E)-9, 12-tetradecadienyl acetate] of the four components of

S. exigua sex pheromone results in mating disruption of both *S. exigua* and *S. litura* (AVRDC, 1997). Traps baited with kairomonal attractants (*e.g.*, p-anisaldehyde and salicylaldehyde) show considerable influence in attracting onion thrips, and could be used as a monitoring and/or mass-trapping tool (AVRDC, 1996a).

- 8.8.5. Biopesticides and bio-control options noted for *S. litura* above can be adapted for both *S. exigua* and *S. litura* on onions.
- 8.8.6. Since onion thrips can rapidly evolve resistance to insecticides, chemical pesticides should be applied only when needed (Nault *et al.*, 2012). For instance, severe thrips infestation during the bulb enlargement period (the period of 10 weeks after transplanting) can lead to substantial yield losses. It is advisable to initiate spraying two weeks before the initiation of bulb enlargement (AVRDC, 2002). This period may vary slightly depending upon latitude, temperature and onion variety used.

8.9 Conclusion

Cultural practices such as crop rotation, clean cultivation, intercropping, trap cropping, mulching and irrigation are effective in reducing the population build-up of major insect and mite pests in tropical vegetable production systems. Healthy seedling production practices should be adopted to postpone the incidence of early season sucking pests, leaf and stem feeders, and delay the transmission of virus diseases by insect vectors. If insect-resistant varieties with desirable horticultural traits are available, farmers should adopt them to avert severe pest damage and substantial yield loss. Colored sticky traps, traps baited with kairomonal attractants or sex pheromone lures should be installed in fields to monitor and/or mass-trap insect pests. Application of chemical pesticides, especially broad-spectrum insecticides, should be avoided early in the growing season to encourage the proliferation of natural enemies, and to avoid the risk of resurgence of secondary pests. Inoculative releases of parasitoids should be made in newer regions where they are not present. Inundative releases of selected egg parasitoids also provide sufficient control of lepidopteran pests on vegetables. The use of biopesticides including *B. thuringiensis*, *M. anisopliae*, *B. bassiana*, NPVs and neem should be increased. Chemical pesticides should be properly integrated with other components of pest management in vegetable crops; for example, it is important to choose selective pesticides, and follow a mode of action rotation with a window approach to effectively curtail development of resistance in the target insects. Novel pesticide deployment strategies such as seed treatment, seedling dips, soil drenching or spot application will reduce the direct contact of toxic chemical pesticides on biocontrol agents and pollinators.

References

- [AVRDC] AVRDC – The World Vegetable Center. 1977. Progress for 1977. Asian Vegetable Research and Development Center, Shanhua, Taiwan, ROC. 90 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1978. Cabbage webworm. In: Progress Report of the Asian Vegetable Research and Development Center, Shanhua, Taiwan. pp. 34-35.
- [AVRDC] AVRDC – The World Vegetable Center. 1981. Progress Report for 1979 (AVRDC 81-148). Asian Vegetable Research and Development Center, Shanhua, Taiwan, Republic of China. 107 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1987. 1984 Progress Report. Asian Vegetable Research and Development Center, Shanhua, Taiwan. 480 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1988. 1986 progress report. Asian Vegetable Research and Development Center, Shanhua, Taiwan. 540 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1991. 1990 Progress Report. Asian Vegetable Research and Development Center, Shanhua, Taiwan. 312 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1992. 1991 Progress Report. Asian Vegetable Research and Development Center, Shanhua, Taiwan. 410 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1993a. AVRDC 1992 Progress Report. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan (ROC). 523 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1993b. Vegetable Research and Development in Southeast Asia: The AVNET Final Report. AVRDC, P.O. Box 205, Taipei 10099. 50 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1995. AVRDC 1994 Progress Report. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan (ROC). 520 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1996a. AVRDC 1995 Report. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan (ROC). 187 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1996b. Vegetable research networking in South Asia: SAVERNET Phase I final report; AVRDC, P.O. Box 42, Shanhua, Tainan, Taiwan 741, ROC. 76 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1997. AVRDC 1996 Report. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan. 172 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1998. AVRDC Report 1997. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan. 191 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 1999. AVRDC Report 1998. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan. vii+148 pp.
- [AVRDC] AVRDC – The World Vegetable Center, 2000. AVRDC Progress Report, 1999. Asian Vegetable Research and Development Center, Shanhua, Taiwan. pp.152
- [AVRDC] AVRDC – The World Vegetable Center. 2001. AVRDC Report 2000. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan. vii + 152 pp.
- [AVRDC] AVRDC – The World Vegetable Center. 2002. AVRDC Report 2001. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan. vii+151 pp.
- FAO, Food and Agriculture Organization. 2011. FAOSTAT. <http://faostat.fao.org> [accessed 3 June 2011].
- Abang AF, Kouame CM, Abang M, Hannah R, Fotso AK. 2013. Vegetable growers perception of pesticide use practices, cost, and health effects in the tropical region of Cameroon. *International Journal of Agronomy and Plant Production* 4(5): 873-883.

- Abe Y. 2006. Exploitation of the serpentine leafminer *Liriomyza trifolii* and tomato leafminer, *L. bryoniae* (Diptera: Agromyzidae) by the parasitoid *Gronotoma micromorpha* (Hymenoptera: Eucoilidae). *European Journal of Entomology* 103: 55–59.
- Adati J, Toriumi W, Sudiarta P, Susila W, Sumiarta K, Suastika G, Purwanto H. 2007. Effect of synthetic sex pheromone traps on the cabbage cluster caterpillar, *Crociodolomia pavonana* and its host preference in the vegetable yield of Indonesia. *Journal of ISSAAS, International Society for Southeast Asian Agricultural Sciences*, 13(2): 105
- Adkins S, Karthikeyan G, Damayanthi T, Kodetham G, Riley DJ, Naidu RA. 2010. IPM CRSP project on tospoviruses and thrips vectors in South and Southeast Asia. *Journal of Insect Science*. <http://www.insectscience.org/10.166/abstract01.html>
- Akhtaruzzaman M, Alam MZ, Ali-Sardar MM. 1999. Suppressing fruit fly infestation by bagging cucumber at different days after anthesis. *Bangladesh Journal of Entomology* 9: 103–112.
- Alam SN, Rashid MA, Rouf FMA, Jhala RC, Patel JR, Satpathy S, Shivalingaswamy TM, Rai S, Wahundeniya I, Cork A, Ammaranan C, Talekar NS. 2003. Development of an integrated pest management strategy for eggplant fruit and shoot borer in South Asia, Technical Bulletin TB28, AVRDC – The World Vegetable Center, Shanhua, Taiwan. 66 pp.
- Alam SN, Hossain MI, Rouf FMA, Jhala RC, Patel MG, Rath LK, Sengupta A, Baral K, Shylesha AN, Satpathy S, Shivalingaswamy TM, Cork A, Talekar NS. 2006. Implementation and promotion of an IPM strategy for control of eggplant fruit and shoot borer in South Asia. Technical Bulletin No. 36. AVRDC publication number 06-672. AVRDC – The World Vegetable Center, Shanhua, Taiwan. 74 pp.
- Alegbejo M, Ogunlana M, Banwo O. 2008. Survey for incidence of *Okra mosaic virus* in northern Nigeria and evidence for its transmission by beetles. *Spanish Journal of Agricultural Research* 6(3): 408–411.
- Amoah P, Drechel P, Abaidoo RC, Ntow WJ. 2006. Pesticide and pathogen contamination of vegetables in Ghana's urban markets. *Arch. Environ. Contam. Toxicol.* 50: 1–6.
- Baker GJ. 2011. Crucifer vegetable insecticide resistance management strategies and issues in Australia. pp. 229–235. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21–25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.
- Baldin ELL, Vendramim JD, Lourencao AL. 2005. Resistance of tomato genotypes to the whitefly *Bemisia tabaci* (Gennadius) biotype B (Hemiptera: Aleyrodidae). *Neotrop Entomol* 34(3): 435–441.
- Ballal CR, Singh SP. 2003. The Effectiveness of *Trichogramma chilonis*, *Trichogramma pretiosum* and *Trichogramma brasiliense* (Hymenoptera: Trichogrammatidae) as Parasitoids of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on Sunflower (*Helianthus annuus*) and Redgram (*Cajanus cajan*). *Biocontrol Science and Technology*, 13: 231–240.
- Bananej K, Ahoonmanesh A, Kheyr-Pour A. 2002. Host Range of an Iranian Isolate of *Watermelon Chlorotic Stunt Virus* as Determined by Whitefly-mediated Inoculation and Agroinfection, and its Geographical Distribution. *Journal of Phytopathology* 150(8-9): 423–430.
- Bautista R, Harris E, Vargas R, Jang E. 2004. Parasitization of melon fly (Diptera: Tephritidae) by *Fopius arisanus* and *Psytalia fletcheri* (Hymenoptera: Braconidae) and the effect of fruit substrates on host preference by parasitoids. *ARS-Research*. http://www.ars.usda.gov/research/publications/Publications.htm?seq_no_115=155470 (26 July 2004).

- Beran F, Mewis I, Srinivasan R, Reinecke A, Hansson BS, Ulrichs C. 2011. Semiochemicals mediating aggregations of *Phyllotreta striolata* (F.) (Coleoptera: Chrysomelidae). pp. 5. In: AVRDC – The World Vegetable Center. Abstracts from the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC Publication No. 11-746. AVRDC – The World Vegetable Center, Taiwan. 101 pp.
- Biswas C, Varma A. 2006. Characterization of virus from pumpkin as an isolate of PRSV-W. *Indian Phytopathology* 59(1): 101-104.
- Bohn GW, Kishaba AN, McCreight JD. 1996. A survey of tolerance to *Aphis gossypii* in part of the world collection of *Cucumis melo* L. In: Cucurbitaceae '96 – Plant Diseases and Disease Resistances. <http://cuke.hort.ncsu.edu/cgc/conferences/cuc96dise.html>
- Boissot N, Lafortune D, Pavis C, Sauvion N. 2003. Field resistance to *Bemisia tabaci* in *Cucumis melo*. *HortScience* 38: 77–80.
- Brink PJ van den, Sureshkumar N, Daam MA, Domingues I, Milwain GK, Beltman WHJ, Perera MWP, Satapornvanit K. 2003. Environmental and human risks of pesticide use in Thailand and Sri Lanka; results of a preliminary risk assessment. Wageningen : Alterra, 2003 (MAMAS Report Series 3/2003). 88 pp.
- Bunwaree AG. 2001. Status of major viruses affecting cucurbitaceous crops in Mauritius. *Revue Agricole et Sucriere de l' Ile Maurice* 80/81(3-1/3): 98-103.
- Cerkauskas R. 2004. Tomato Yellow Leaf Curl Virus (TYLCV). AVRDC Publication 04-610. <http://www.hazera.com/sa/marketing/TYLCV1.pdf>
- Chen TY, Chu CC, Fitzgerald G, Natwick ET, Henneberry TJ. 2004. Trap Evaluations for Thrips (Thysanoptera: Thripidae) and Hoverflies (Diptera: Syrphidae). *Environ. Entomol.* 33(5): 1416-1420.
- Chen WM. 2002. Preliminary Studies on the Insecticide-Resistance and Management of Indian Cabbage Moth. *Diaphania indica* (Saunders). Master's thesis, South China University of Tropical Agriculture.
- Cork A, Alam SN, Rouf FMA, Talekar NS. 2003. Female pheromone of brinjal fruit and shoot borer, *Leucinodes orbonalis*: trap optimization and preliminary mass trapping trials. *Bulletin of Entomological Research*, 93: 107–113.
- Cork A, Beevor PS, Hall DR, Nesbitt BF, Campion DG. 1985. A sex attractant for the spotted bollworm *Earias vittella*. *Trop. Pest Manag.* 31: 156-158.
- Cork A, Chamberlain DJ, Beevor PS, Hall DR, Nesbitt BF, Campion DG, Attique MR. 1988. Components of female sex pheromone of spotted bollworm, *Earias vittella* F. (Lepidoptera: Noctuidae): identification and field evaluation in Pakistan. *J. Chem. Ecol.* 14: 929-945.
- Dahal G, Lecoq H, Albrechtsen SE. 1997. Occurrence of papaya ringspot potyvirus and cucurbit viruses in Nepal. *Annals of Applied Biology* 130(3): 491-502.
- Dharma Putra GNG, Sudiarta IP, Dharma IP, Sumiarta K, Srinivasan R. 2013. Monitoring of imago of *Spodoptera litura* and *Helicoverpa armigera* using sex pheromone trap. *E-Jurnal Agroekoteknologi Tropika* 2(1): 56-61. (<http://ojs.unud.ac.id/index.php/JAT/article/view/4579/3506>)
- Dhillon MK, Singh R, Naresh JS, Sharma HC. 2005. The melon fruit fly, *Bactrocera cucurbitae*: A review of its biology and management. *J. Insect Sci.* 5: 1–16.
- Dhillon NPS, Ranjana R, Kuldeep S, Eduardo I, Monforte AJ, Pitrat M, Dhillon NK, Singh PP. 2007. Diversity among landraces of Indian snapmelon (*Cucumis melo* var. *momordica*). *Genetic Resources and Crop Evolution* 54: 1267–1283.

- Duffus JE, Stenger DC. 1998. *Squash leaf curl virus*. CMI/AAB Descriptions of Plant Viruses No. 358. Kew, UK: Commonwealth Mycological Institute. <http://www.dpvweb.net/dpv/showdpv.php?dpvno=358> (accessed on June 18, 2013).
- Durairaj C, Shobanadevi R, Suresh S, Natarajan S. 2007. A non-chemical method for the management of leaf miner and whitefly in brinjal. *Acta Horticulturae* 752: 527-529.
- Ekesi S, Maniania NK, Mohamed SA, Lux SA. 2005. Effect of soil application of different formulations of *Metarhizium anisopliae* on African tephritid fruit flies and their associated endoparasitoids. *Biological Control* 35(1): 83-91.
- Fancelli M, Vendramim JD. 2002. Development of *Bemisia tabaci* (Gennadius, 1889) biotype B on *Lycopersicon* spp. genotypes. *Scientia Agricola* 59: 665-669.
- Fasunwon BT, Banjo AD. 2010. Seasonal population fluctuations of *Podagrica species* on okra plant (*Abelmoschus esculentus*). *Research Journal of Agriculture and Biological Sciences* 6(3): 283-288.
- Feng X, Li ZY, Wu QJ, Chen AD, Wu YD, Hou YM, He YR, Li JH, Xie SH, Zhang JM, Wie F, Ma CS. 2011. Rising to the challenge: a national project on sustainable control of DBM in China. pp. 13-16. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.
- Firdaus S, van Heusden AW, Hidayati N, Supena EDJ, Mumm R, de Vos RCH, Visser RGF, Vosman B. 2013. Identification and QTL mapping of whitefly resistance components in *Solanum galapagense*. *Theor Appl Genet*. 126: 1487-1501.
- Firdaus S, van Heusden AW, Hidayati N, Supena EDJ, Visser RGF, Vosman B. 2012. Resistance to *Bemisia tabaci* in tomato wild relatives. *Euphytica* 187: 31-45.
- Fowler D. 2011. Nutritional Value of Common Fruits & Vegetables Grown in Cambodia. MCC Prey Veng, Cambodia. p. 8. <http://thehomeaway.files.wordpress.com/2011/09/nutritional-value-of-cambodian-crops.pdf>
- Furtado IP, de Moraes GJ, Kreiter S, Garcin M-S, Knapp M. 2007. Potential of a Brazilian population of the predatory mite *Phytoseiulus longipes* as a biological control agent of *Tetranychus evansi* (Acari: Phytoseiidae, Tetranychidae). *Biol Control* 42:139-147.
- Gaba V, Zelcer A, Gal-On A. 2004. Invited Review: Cucurbit biotechnology – The importance of virus resistance, *In Vitro Cellular & Developmental Biology – Plant* 40: 346-358.
- Ganehilarachchi GASM. 1997. Aspects of the biology of *Diaphania indica* (Lepidoptera : Pyralidae). *J. Natn. Sci. Coun. Sri Lanka* 25(4): 203-209.
- Genova C, Weinberger K, Acedo AL Jr. 2010. An assessment of market opportunities for leafy vegetables in the upland areas of Cambodia, Laos, and Vietnam. Final project report, RETA 6376. AVRDC Publication No. 10-734. AVRDC – The World Vegetable Center, Taiwan. 47 pp.
- Ghosh SK, Chaudhary M, Kumar P. 2011. Myco Jaal® – A novel formulation of *Beauveria bassiana* for managing diamondback moth (*Plutella xylostella*) in sub tropical and tropical crucifer production system. pp. 143-148. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.

- Gogi MD, Ashfaq M, Arif MJ, Sarfraz RM, Nawab NN. 2010. Investigating phenotypic structures and allelochemical compounds of the fruits of *Momordica charantia* L. genotypes as sources of resistance against *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae). *Crop Protection*, 29: 884-890.
- Gupta RK, Desh Raj, Nirmala Devi. 2004. Biological and Impact Assessment Studies on *Campoletis chloridae* Uchida: A Promising Solitary Larval Endoparasitoid of *Helicoverpa armigera* (Hübner). *Journal of Asia Pacific Entomology*. 7: 239 – 247.
- Hall DR, Beevor PS, Lester R, Nesbitt BF. 1980. (E,E)-10,12-hexadecadienal: a component of the female sex pheromone of the spiny bollworm, *Earias insulana* (Boisd.) (Lepidoptera: Noctuidae). *Experientia* 36: 152-154.
- Hamed M, Nadeem S. 2010. Forecasting of spotted bollworm (*Earias vitella* (Fab.) (Lepidoptera: Noctuidae) Occurrence in Cotton. *Pakistan J. Zool.* 42(5): 575-580.
- Haque MN, Kundu AK, Akanda AM, Bhuiyan MKA. 2004. Comparative biological and serological properties of Watermelon mosaic virus 2 and Zucchini yellow mosaic virus. *Bangladesh Journal of Plant Pathology* 20(1/2): 71-76.
- Ho CC. 2000. Spider-mite problems and control in Taiwan. *Experimental and Applied Acarology* 24: 453-462.
- Hoi PV. 2010. Governing pesticide use in vegetable production in Vietnam. PhD Dissertation, Wageningen University, The Netherlands. pp. 181.
- Isakeit T, Robertson NL. 1994. First Report of Squash Leaf Curl Virus on Watermelon in Texas. *Plant Disease* 78: 1010.
- Islam MT, Omar D, Latif MA, Morshed MM. 2011. The integrated use of entomopathogenic fungus, *Beauveria bassiana* with botanical insecticide, neem against *Bemisia tabaci* on eggplant. *African Journal of Microbiology Research* 5(21): 3409-3413.
- Kalbfleisch S. 2006. Integrated Pest Management of *Hellula undalis* on Crucifers in Central Luzon, Philippines, with E,E-11,13-Hexadecadienal as Synthetic Sex Pheromone', unpublished dissertation, Technische Universität München.
- Kannan M, Vijayaraghavan C, Jayaprakash SA, Uthamsamy S. 2011. Studies on the biology and toxicity of newer insecticide molecules on cabbage head caterpillar, *Crociodolomia binotalis* (Zeller) (Lepidoptera: Pyralidae) in India. pp. 26-32. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.
- Klungness LM, Jang EB, Mau RFL, Vargas RI, Sugano JS, Fujitani E. 2005. New approaches to sanitation in a cropping system susceptible to tephritid fruit flies (Diptera: Tephritidae) in Hawaii. *Journal of Applied Science and Environmental Management* 9: 5-15.
- Krishnamoorthy A. 2012. Exploitation of egg parasitoids for control of potential pests in vegetable ecosystems in India. *Comunicata Scientiae* 3(1): 1-15.
- Kumar ARV, Shivakumara B. 2003. Variable response of male *Helicoverpa armigera* moth to sex pheromone blends: A case of behavioural polymorphism? *Current Science* 84: 705-709.
- Kumar P, Alma Linda A, Ketelaar JW, Shanmugam V. 2011. Field Exercise Guides on Fruit Flies Integrated Pest Management. Pp. 63. http://ipm.ait.asia/wp-content/uploads/2010/11/Field_Exercise_Guide_on_Fruit_Fly_IPM-N.pdf
- Lecoq H, Cohen S, Pitrat M, Labonne G. 1979. Resistance to cucumber mosaic virus transmission by aphids in *Cucumis melo*. *Phytopathology* 69: 1223–1225.

- Lecoq H, Wisler G, Pitrat M. 1998. Cucurbit viruses: the classics and emerging. In: McCreight JD (Ed.), *Cucurbitaceae '98, Evaluation and Enhancement of Cucurbit Germplasm*, pp. 126–142. ASHS Press, Alexandria, VA 22314–2562, USA.
- Lee HS, Lu FM, Wen HC. 1990. Effects of temperature on the development of leafminer, *Liriomyza bryoniae* (Kaltenbach) (Diptera: Agromyzidae) in Taiwan. *Chinese Journal of Entomology* 10: 143–150.
- Malathi S, Sriramulu M. 2000. Laboratory Efficacy of Biotic Insecticides against Lepidopterous Pests Fed on Treated Cabbage Leaves. *Shashpa*, 7: 63–66.
- Martin B, Alvarez JI, Fereres A. 1996. Feeding behaviour of *Aphis gossypii* during transmission of cucumber mosaic virus on resistance and susceptible *Cucumis melo* lines. In: *Cucurbitaceae '96 – Plant Diseases and Disease Resistances*. <http://cuke.hort.ncsu.edu/cgc/conferences/cuc96dise.html>
- Mathur A, Singh S, Singh NP, Meena M. 2012. Field evaluation of plant products and microbial formulations against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee under semi-arid conditions of Rajasthan. *J. Biopest.* 5(1): 71–74.
- Mcquate GT. 2011. Assessment of attractiveness of cassava as a roosting plant for melon fly, *Bactrocera cucurbitae*, and oriental fruit fly, *B. dorsalis*. *Journal of Insect Science* 11: 1536–2442.
- Middleton C, Makarady K, Koma YS. 2004. Dangerous pesticide dealing in Cambodia. *Pesticides News*, 65: 4–5.
- Mirmoayedi A, Maniee M, Yoghutipoor A. 2010. Control of Cotton Spiny Bollworm, *Earias insulana* Boisduval, Using Three Bio-Insecticides, Bt, Spinosad and Neem-Azal. *Journal of Entomology* 7(2): 89–94.
- Mohan MC, Reddy NP, Devi UK, Ramesh Kongara, Sharma HC. 2007. Growth and insect assays of *Beauveria bassiana* with neem to test their compatibility and synergism. *Biocontrol Science and Technology*, 17: 1059–1069.
- Mote UN. 1976. Control of onion thrips (*Thrips tabaci* Lin.). *Pesticide* 10: 42–43.
- Muigai SG, Bassett MJ, Schuster DJ, Scott JW. 2003. Greenhouse and field screening of wild *Lycopersicon* germplasm for resistance to the whitefly *Bemisia argentifolii*. *Phytoparasitica* 31(1): 27–38.
- Muniappan R, Shepard BM, Carner GR, Ooi PAC. 2012. *Arthropod pests of horticultural crops in tropical Asia*. Wallingford, Oxfordshire, CABI. 168 pp.
- Muniyappa V, Padmaja AS, Venkatesh HM, Sharma A, Chandrashekar S, Kulkarni RS, Hanson PM, Chen JT, Green SK, Colvin J. 2002. Tomato leaf curl virus resistant tomato lines TLB111, TLB130 and TLB182. *HortScience* 37: 603–606.
- Nagaraju N, Venkatesh HM, Warburton H, Muniyappa V, Chancellor TCB, Colvin J. 2002. Farmers' perceptions and practices for managing tomato leaf curl virus disease in southern India. *International Journal of Pest Management* 48: 333–338.
- Naojee K, Abeeluck D, Rawananshah T, Facknath S. 2010. The use of blue sticky traps in management of Thrips palmi (*Karny*) (*Thysanoptera: Thripidae*) in eggplant cultivation in Mauritius. In: *Sustainable Green Intelligent Innovative Island (Research Week 2009–2010)*, 15 – 19 February 2010, University of Mauritius. <http://vcampus.uom.ac.mu/researchweek/rw0910/abstracts/Agri4.pdf>
- Nathan SS, Kalaivani K. 2006. Combined effects of azadirachtin and nucleopolyhedrovirus (SpVtNPV) on *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae) larvae. *Biological Control* 39: 96–104.

- Nault BA, Shelton AM, Hsu CL, Hoepting CA. 2012. How to win the battle against onion thrips, 6 pgs. In: Proceedings of the 2012 Empire State Fruit & Vegetable EXPO. January 24-26, 2012. Syracuse, NY. <http://www.hort.cornell.edu/expo/proceedings/2012/Onions/onion%20Nault.pdf>
- Neufeld DSG, Savoeun H, Phoeurk C, Glick A, Hernandez C. 2010. Prevalence and Persistence of Organophosphate and Carbamate Pesticides in Cambodian Market Vegetables. *Asian Journal of Water, Environment and Pollution*, 7(4): 89-98.
- Niassy S, Maniania NK, Subramanian S, Gitonga LM, Ekesi S. 2012. Performance of a semiochemical-baited autoinoculation device treated with *Metarhizium anisopliae* for control of *Frankliniella occidentalis* on French bean in field cages. *Entomologia Experimentalis et Applicata* 142: 97-103.
- Nielsen MC, Worner SP, Chapman RB, de Kogel WJ, Perry NB, Sansom CM, Murai T, Muvea AM, Sevgan S, Teulon DAJ. 2010. Optimising the use of allelochemicals for thrips pest management. Paper presented at the International Society of Chemical Ecology Conference, 26th Annual Meeting, Tours, France.
- Niranjana RF, Wijeyagunsekara, HNP, Raveendranath S. 2005. Parasitoids of *Liriomyza sativae* in farmer fields in the Batticaloa district. *Tropical Agricultural Research* 17: 214-220.
- Nyambo B, Lohr B. 2005. The role and significance of farmer participation in biocontrol-based IPM for Brassica crops in East Africa. In: Second International Symposium on Biological Control of Arthropods, September 12-16, 2005. Davos, Switzerland. pp. 290-301.
- Ooi PAC. 1980. The Pathogenicity of *Bacillus thuringiensis* for *Crocidolomia binotalis*. MAPPS Newsletter, 3: 4.
- Parimi S, Zehr UB. 2009. Insect resistant crops are an integral part of IPM programs: Bt brinjal. In: Sol 2009: the 6th Solanaceae Genome Workshop, 2009, New Delhi, India, Book of Abstracts, p. 145.
- Parker BL, Talekar NS, Skinner M. 1995. Field guide: Insect pests of selected vegetables in tropical and subtropical Asia. Asian Vegetable Research and Development Center, Shanhua, Tainan, Taiwan, ROC. Publication no. 94-427. 170 pp.
- Parrella MP. 1987. Biology of *Liriomyza*. *Annual Review of Entomology* 32: 201-224.
- Pawar DB, Mote UN, Lawande KE. 1991. Monitoring of fruit fly population in bitter melon crop with the help of lure trap. *Journal of Research, Maharashtra Agricultural Universities* 16: 281.
- Provvidenti R. 1996. Diseases caused by viruses. In: Zitter TA, Hopkins DL, Thomas CE (Eds.). Compendium of cucurbit diseases, pp. 37-45. APS Press, St. Paul, Minnesota.
- Rao NGV, Majumdar A, Mandaokar AD, Nimbalkar SA, Kumar PA. 1999. Susceptibility of brinjal shoot and fruit borer to the δ -endotoxins of *Bacillus thuringiensis*. *Current Science* 77(3): 1-2.
- Rashid MA, Rahman MA, Ahmad S, Alam SN, Rezaul Karim ANM, Luther G, Miller S. 2003. Varietal screening of eggplant for resistance to bacterial wilt, fruit and shoot borer, jassid and root-knot. Tenth Annual Report, IPM CRSP, Virginia Tech. USA, pp. 125-128.
- Reddy AS, Rosaiah B. 1987. Insect pest management. In: Veerabhadra Rao M, Sithanatham S (eds.), Plant protection in field crops, Publ. 4, Plant Protection Association of India, Hyderabad, pp. 293-299.
- Reddy MRS, Reddy GS. 1999. An eco-friendly method to combat *Helicoverpa armigera* (Hub.). *Insect Environ.* 4: 143-144.

- Ridland PM, Endersby NM. 2011. Some Australian populations of diamondback moth, *Plutella xylostella* (L.) show reduced susceptibility to fipronil. pp. 195-203. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.
- Rosendahl I, Laabs V, James BD, Atcha-Ahowe C, Agbotse SK, Kone D, Kogo A, Salawu R, Glitho IA. 2008. Living with Pesticides: A Vegetable Case Study, SP-IPM, IITA, Nigeria. http://www.spipm.cgiar.org/c/document_library/get_file?p_l_id=17828&folderId=18430&name=DLFE-77.pdf
- Saikia AK, Muniyappa V. 1989. Epidemiology and control of tomato leaf curl virus in Southern India. *Trop Agric (Trinidad)* 66: 350–354.
- Sandur S. 2004. Implications of diamondback moth control for Indian farmers. Consultant Report for the Centre for Environmental Stress and Adaptation Research. La Trobe University, Victoria, Australia, 31 pp.
- Sastrosiwojo S, Setiawati W. 1992. Biology and Control of *Crociodolomia binotalis* in Indonesia. In: Management of Diamondback Moth and Other Crucifer Pests: Proceedings of the Second International Workshop, ed. N.S. Talekar, Shanhuai, Taiwan: Asian Vegetable Research and Development Center, pp. 81-87.
- Satpute NS, Deshmukh SD, Rao NGV, Tikar SN, Moharil MP, Nimbalkar SA. 2003. Native insecticide resistance in spotted bollworm, *Earias vitella* (Fabricius) in Western Vidarbha region of India. *Resistant Pest Management Newsletter* 13(1): 9-11.
- Schreiner IH. 1991. Damage threshold for *Diaphania indica* Saunders (Lepidoptera: Pyralidae) on cucumbers in Guam. *Tropical Pest Management* 37(1): 17-20.
- Senthilkumar N, Murugan K, Zhang W. 2008. Additive interaction of *Helicoverpa armigera* Nucleopolyhedrovirus and Azadirachtin. *BioControl.*, 53: 869–880.
- Shaifullah SMK, Kundu AK, Akanda AM, Haque MN. 2005. Detection and characterization of Cucumber mosaic virus naturally infecting pumpkin plants in Southern Bangladesh. *Bangladesh Journal of Plant Pathology* 21(1/2): 11-17.
- Shelton AM, Badenes-Pérez FR. 2006. Concepts and applications of trap cropping in pest management. *Annual Review of Entomology* 51: 285-309.
- Shelton AM, Gujar GT, Chen M, Rauf A, Srinivasan R, Kalia V, Mittal A, Kumari A, Ramesh K, Borkakatti A, Zhao JZ, Endersby N, Russell D, Wu YD, Uijtewaal B. 2009. Assessing the susceptibility of cruciferous Lepidoptera to Cry1Ba2 and Cry1Ca4 for future transgenic cruciferous vegetables. *Journal of Economic Entomology*, 102(6): 2217-2223.
- Shetty AA, Singh JP, Singh D. 2013. Resistance to yellow vein mosaic virus in okra: a review. *Biological Agriculture & Horticulture*. DOI: 10.1080/01448765.2013.793165
- Shivalingaswamy TM, Satpathy, S. 2007. Integrated pest management in vegetable crops. In: Jain PC, Bhargava MC (eds.), *Entomology: Novel Approaches*, New India Publishing Agency, New Delhi, India. pp. 353-375.
- Singh IP, Seema K, Singh R. 2000. Bioefficacy of Insecticides against Cabbage Head Borer (*Hellula undalis* Fabr.) and Dissipation of Chlorpyrifos in Cabbage. *Journal of Applied Biology*, 10: 194-198.
- Sivapragasam A. 1996. Integrated management of cabbage with special reference to the control of *Hellula undalis* (Fabr.). In: Hokyo N, Norton G (eds.), *Proc. Int. Workshop on the Pest Management Strategies in Asian Monsoon Agroecosystems*. pp. 223-234.

- Sivapragasam A, Chua TH. 1997. Preference for sites within plant by larvae of the cabbage webworm, *Hellula undalis* (Fabr.) (Lep., Pyralidae). *Journal of Applied Entomology* 121: 361-365.
- Sivapragasam A, Syed AR. 1999. The problem and management of agromyzid leafminers on vegetables in Malaysia. In Proceedings of a Workshop on Leafminers of Vegetables in Southeast Asia, Lim GS, Soetikno SS, Loke WH (eds.), Serdang, Malaysia, CAB International, Southeast Asia Regional Centre, pp. 36-41.
- Smyth RR, Hoffmann MP, Shelton AM. 2003. Larval performance in relation to labile oviposition preference of *Crociodolomia pavonana* [F.] (Lepidoptera: Pyralidae) among phenological stages of cabbage. *Environmental Entomology* 32: 765-770.
- Sodavy P, Sitha M, Nugent R, Murphy H. 2000. Situation Analysis on Farmers' Awareness and Perceptions of the Effect of Pesticides on their Health. FAO COMMUNITY IPM PROGRAMME, Field document. pp.16. <http://www.communityipm.org/toxictrail/Documents/Cambodia-SituationAnalysis.pdf>
- Srinivasan K, Krishna Moorthy PN, Raviprasad TN. 1994. African marigold as a trap crop for the management of the fruit borer, *Helicoverpa armigera* on tomato. *International Journal of Pest Management*, 40: 56-63.
- Srinivasan K, Krishnamoorthy P. 1992. Development and adoption of integrated pest management for major pests of cabbage using Indian Mustard as trap crop. In: Management of Diamondback moth and other crucifer pests. Proceedings of the second International Workshop. (Ed.) Talekar, NS. Asian Vegetable Research and Development Center, Shanhua, Taiwan. 603 pp.
- Srinivasan R, Hsu YC. 2008. Susceptibility of major lepidopterans to δ -endotoxins and a formulation of *Bacillus thuringiensis* (B.t.) on vegetable brassicas in Taiwan. *Biocontrol Science and Technology*. 18(9): 935 – 939.
- Srinivasan R, Huang CC. 2009. The effect of simulated borer infested shoot pruning on yield parameters of eggplant. *Journal of Asia Pacific Entomology*. 12: 41 – 43.
- Srinivasan R, Su FC, Huang CC. 2013. Oviposition dynamics and larval development of *Helicoverpa armigera* on a highly preferred unsuitable host plant, *Solanum viarum*. *Entomologia Experimentalis et Applicata* 147: 217-224.
- Srinivasan R, Uthamasamy S. 2004. Rapid Plant Screening Technique – a novel tool for the evaluation of antixenosis resistance mechanism of tomato accessions to Whitefly in laboratory condition. *Resistant Pest Management Newsletter* 14(1): 4-5.
- Srinivasan R. 2012. Integrating biopesticides in pest management strategies for tropical vegetable production. *Journal of Biopesticides* 5: 36 – 45.
- Srinivasan R, Lin M-Y, Sopana Yule, Chuanpit Khumsuwan, Thanh Hien, Vu Manh Hai, Lê Đức Khánh, Bhanu KRM. 2013. Use of insect pheromones in vegetable pest management: Successes and struggles. In: New Horizons in Insect Science. Chakravarthy AK, Ashok Kumar CT, Abraham Verghese, Thyagaraj NE (eds.), International Conference on Insect Science, 14-17 Feb 2013, Bangalore, India. pp. 6.
- Subba Rao BR, Parshad B, Ram A, Singh RP, Srivastava ML. 1968. Distribution of *Empoasca devastans* and its egg parasites in the Indian Union. *Entomologia Experimentalis et Applicata* 11(2): 250-254.

- Sukonthabhirom S, Dumrongsak D, Jumroon S, Saroch T, Chaweng A, Tanaka T. 2011. Update on DBM diamide resistance from Thailand: causal factors and learnings. Pp. 190-194. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.
- Tabashnik BE, Cushing NL, Finson N, Johnson MW. 1990. Field development of resistance to *Bacillus thuringiensis* in diamondback moth (Lepidoptera: Plutellidae). *J. Econ. Entomol.*, 83: 1671-1676.
- Talekar NS, Opana RT, Hanson P. 2006. *Helicoverpa armigera* management: a review of AVRDC's research on host plant resistance in tomato. *Crop Protection* 25(5): 461-467.
- Talekar NS, Shelton AM. 1993. Biology, ecology, and management of the diamondback moth. *Annu. Rev. Entomol.* 38: 275-301.
- Tatagar MH, Mohankumar HD, Shivaprasad M, Mesta RK. 2009. Bio-efficacy of flubendiamide 20 WG against chilli fruit borers, *Helicoverpa armigera* (Hub.) and *Spodoptera litura* (Fb.). *Karnataka J. Agric. Sci.* 22(3): 579-581.
- Tewari GC, Krishnamoorthy PN. 1984. Yield loss in tomato caused by fruit borer. *Indian J. Agric. Sci.* 54: 341-343.
- Toscano LC, Boic, a JAL, Maruyama WI. 2002. Nonpreference of whitefly for oviposition in tomato genotypes. *Scientia Agricola* 59: 677-681.
- Tripathi MK, Maurya TN. 2011. Efficacy of synthetic, botanical and biopesticides for management of *Earias vitella* in okra. *Journal of Experimental Zoology* 14(2): 595-596.
- Uelese A. 2011. The potential of *Trichogramma chilonis* Ishii (Hymenoptera: Trichogrammatidae) as a biological control agent for *Crociodolomia pavonana* F. (Lepidoptera: Crambidae). pp. 50-51. In: AVRDC – The World Vegetable Center. 2011. Abstracts from the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21-25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC Publication No. 11-746. AVRDC – The World Vegetable Center, Taiwan. 101 pp.
- Ulrichs C, Mewis I. 2003. Evaluation and Risk-Assessment of Bio-Rational Pesticides for Controlling Crucifer Pests in Urban Horticulture in the Philippines. In: Balder H, Strauch KH, Backhaus GF. (eds.), Proceedings of the Second International Symposium on Plant Health in Urban Horticulture. pp. 211-221.
- van der Meer FW, Garnett HM. 1987. Purification and identification of a South African isolate of watermelon mosaic virus - Morocco. *Journal of Phytopathology* 120(3): 255-270.
- Varela AM, Seif A. 2004. A Guide to IPM and hygiene standards in okra production in Kenya. The International Centre of Insect Physiology and Ecology, Nairobi, Kenya. p. 116.
- Vargas RI, Stark JD, Kido MH, Ketter HM, Whitehand LC. 2000. Methyl eugenol and cue-lure traps for suppression of male oriental fruit flies and melon flies (Diptera: Tephritidae) in Hawaii: effects of lure mixtures and weathering. *J Econ Entomol.* 93(1): 81-87.
- Vargas RI, Mau RFL, Jang EB, Faust RM, Wong L. 2008. The Hawaii Fruit Fly Area-wide Pest Management Programme. *Publications from USDA-ARS / UNL Faculty*. Paper 656. <http://digitalcommons.unl.edu/usdaarsfacpub/656>
- Wakamura S, Arakaki N, Kinjo K, Yasuda T. 1998. Sex pheromone of the cotton caterpillar, *Diaphania indica* (Saunders) (Lepidoptera: Pyralidae): Identification and field attraction. *Appl. Entomol. Zool.* 33: 429-434.

- Wakman W, Kontong MS, Teakle DS, Persley DM. 2002. Watermelon mosaic virus of pumpkin (*Cucurbita maxima*) from Sulawesi: identification, transmission, and host range. *Indonesian Journal of Agricultural Science* 3(1): 33-36.
- Wekesa VW, Maniania NK, Knapp M, Boga HI. 2005. Pathogenicity of *Beauveria bassiana* and *Metarhizium anisopliae* to the tobacco spider mite *Tetranychus evansi*. *Exp Appl Acarol* 36: 41–50.
- Wekesa WV, de Moraes GJ, Knapp M, Delalibera I Jr. 2007. Interactions of two natural enemies of *Tetranychus evansi*, the fungal pathogen *Neozygites Xoridana* (Zygomycetes: Entomophthorales) and the predatory mite, *Phytoseiulus longipes* (Acari: Phytoseiidae). *Biol Control* 41:408–414.
- Williamson S. 2003. The dependency syndrome: Pesticide use by African smallholders, Pesticide Action Network UK, London.
- Zenil M, Liedo P, Williams T, Valle J, Cancino J, Montoya P. 2004. Reproductive biology of *Fopius arisanus* (Hymenoptera: Braconidae) on *Ceratitis capitata* and *Anastrepha* spp. (Diptera: Tephritidae). *Biological Control* 29: 169–178.
- Zhao XQ, Li XY, Yin YQ, Chen AD. 2011. Occurrence and control of *Plutella xylostella* (Lepidoptera: Plutellidae) in Yunnan, China. pp. 273-277. In: Srinivasan R, Shelton AM, Collins HL, eds. 2011. Proceedings of the Sixth International Workshop on Management of the Diamondback Moth and Other Crucifer Insect Pests, 21–25 March 2011, Kasetsart University, Nakhon Pathom, Thailand. AVRDC – The World Vegetable Center, Publication No. 11 755. AVRDC – The World Vegetable Center, Taiwan. 321 pp.

