The Internet Gaming Disorder Test (IGD-20 Test) (Pontes et al., 2014)

Instructions: These questions relate to your gaming activity during the past year (i.e., 12 months). By gaming activity we mean any gaming-related activity that was played on either a computer/laptop, gaming console and/or any other kind of device online and/or offline.

	Strongly disagree	Disagree	Neither agree or disagree	Agree	Strongly agree
1. I often lose sleep because of long gaming sessions.	○	○	○	○	○
2R. I never play games in order to feel better.	○	○	○	○	○
3. I have significantly increased the amount of time I play games over last year.	○	○	○	○	○
4. When I am not gaming I feel more irritable.	○	○	○	○	○
5. I have lost interest in other hobbies because of my gaming.	○	○	○	○	○
6. I would like to cut down my gaming time but it's difficult to do.	○	○	○	○	○
7. I usually think about my next gaming session when I am not playing.	○	○	○	○	○
8. I play games to help me cope with any bad feelings I might have.	○	○	○	○	○
9. I need to spend increasing amounts of time engaged in playing games.	○	○	○	○	○
10. I feel sad if I am not able to play games.	○	○	○	○	○
11. I have lied to my family members because the amount of gaming I do.	○	○	○	○	○
12. I do not think I could stop gaming.	○	○	○	○	○
13. I think gaming has become the most time consuming activity in my life.	○	○	○	○	○
14. I play games to forget about whatever's bothering me.	○	○	○	○	○
15. I often think that a whole day is not enough to do everything I need to do in-game.	○	○	○	○	○
16. I tend to get anxious if I can't play games for any reason.	○	○	○	○	○
17. I think my gaming has jeopardized the relationship with my partner.	○	○	○	○	○
18. I often try to play games less but find I cannot.	○	○	○	○	○
19R. I know my main daily activity (i.e., occupation, education, homemaker, etc.) has not been negatively affected by my gaming.	○	○	○	○	○
20. I believe my gaming is negatively impacting on important areas of my life.	○	○	○	○	○

Scoring information:

In order to obtain participants' total scores, all 20 items must be summed up. However, to compute factor-specific scores, the items representing each factor must be averaged due to the imbalanced number of indicators of the factors. As reported by Pontes et al., (2014), the suggested cut-off point for the IGD-20 Test is 71 points.

- **Salience** (Factor 1): items: 1, 7, and 13.
- **Mood Modification** (Factor 2): items: 2R, 8, and 14.
- **Tolerance** (Factor 3): items: 3, 9, and 15.
- **Withdrawal Symptoms** (Factor 4): items: 4, 10, and 16.
- **Conflict** (Factor 5)**:** items: 5, 11, 17, 19R, and 20.
- **Relapse** (Factor 6): items: 6, 12, and 18.

The following items are reversely scored: 2R and 19R.

References:

Pontes, H. M., & Griffiths, M. D. (2017). The development and psychometric evaluation of the Internet Disorder Scale (IDS-15). *Addictive Behaviors, 64*, 261-268. doi: 10.1016/j.addbeh.2015.09.003

Fuster, H., Carbonell, X., Pontes, H. M., & Griffiths, M. D. (2016). Spanish validation of the Internet Gaming Disorder-20 (IGD-20) Test. *Computers in Human Behavior, 56*, 215-224. doi:10.1016/j.chb.2015.11.050

Hawi, N. S., & Samaha, M. (2017). Validation of the Arabic Version of the Internet Gaming Disorder-20 Test. *Cyberpsychology, Behavior, and Social Networking, 20*(4), 268-272. doi:10.1089/cyber.2016.0493