

7 Proving Non-Conditional Statements

Prove the following statements.

2. Suppose $x \in \mathbb{Z}$. Then x is odd if and only if $3x + 6$ is odd.

Proof. First we prove that x being odd implies $3x + 6$ is odd. Suppose x is odd. Then, by definition of an odd number, $x = 2k + 1$ for some integer k . Thus $3x + 6 = 3(2k + 1) + 6 = 6k + 9 = 2(3k + 4) + 1$. Therefore $3x + 6$ is odd by definition of an odd number.

Conversely, we need to prove that $3x + 6$ being odd implies x is odd. We use contrapositive proof. Suppose x is not odd. Then x is even, so $x = 2k$ for some integer k . Thus $3x + 6 = 3(2k) + 6 = 6k + 6 = 2(3k + 3)$. Therefore $3x + 6$ is even by definition of an even number. Thus $3x + 6$ is not odd. Therefore by contrapositive we have proved that $3x + 6$ being odd implies x is odd. $\square$

4. Given an integer a , then $a^3 + a^2 + a$ is even if and only if a is even.

Proof. First we prove that $a^3 + a^2 + a$ being even implies that a is even. We use contrapositive proof. Suppose a is not even, then a is odd. Thus $a = 2k + 1$ for some integer k . Hence, $a^3 + a^2 + a = (2k + 1)^3 + (2k + 1)^2 + 2k + 1$. After expanding we get $16k^3 + 24k^2 + 14k + 2 + 1 = 2(8k^3 + 12k^2 + 7k + 1) + 1$. Therefore, $a^3 + a^2 + a$ is odd by definition of an odd number.

Conversely, we need to prove that a being even implies that $a^3 + a^2 + a$ is even. Suppose a is even. Then $a = 2k$ for some integer k . Then $a^3 + a^2 + a = (2k)^3 + (2k)^2 + (2k) = 8k^3 + 4k^2 + 2k = 2(4k^3 + 2k^2 + k)$. Therefore, $a^3 + a^2 + a$ is even by definition of an even number, $\square$

6. Suppose $x, y \in \mathbb{R}$. Then $x^3 + x^2y = y^2 + xy$ if and only if $y = x^2$ or $y = -x$.

Proof. First we prove that if $x^3 + x^2y = y^2 + xy$ then $y = x^2$ or $y = -x$. Assume $x^3 + x^2y = y^2 + xy$. From this we get $x^2(x + y) = y(x + y)$. Dividing both sides by $(x + y)$ we see that $y = x^2$.

Conversely, we need to prove that if $y = x^2$ or $y = -x$ then $x^3 + x^2y = y^2 + xy$, we look at both cases.

Case 1. Assume $y = x^2$. By substituting this value for y in the equation we get $x^3 + x^4 = x^4 + x^3$.

Case 2. Assume $y = -x$. By substituting this value for y in the equation we get $x^3 - x^2 = x^2 - x^2$.

Therefore $x^3 + x^2y = y^2 + xy$ both when $y = x^2$ and $y = -x$. □

8. Suppose $a, b \in \mathbb{Z}$. Prove that $a \equiv b \pmod{10}$ if and only if $a \equiv b \pmod{2}$ and $a \equiv b \pmod{5}$.

Proof. First we prove that if $a \equiv b \pmod{10}$ then $a \equiv b \pmod{2}$ and $a \equiv b \pmod{5}$. Assume $a \equiv b \pmod{10}$. Then $(a - b) = 10k$ for some integer k . Hence $(a - b) = 2(5k)$ and $(a - b) = 5(2k)$. In other words, $a \equiv b \pmod{2}$ and $a \equiv b \pmod{5}$.

Conversely, we prove that if $a \equiv b \pmod{2}$ and $a \equiv b \pmod{5}$ then $a \equiv b \pmod{10}$. We prove this by contrapositive. Suppose $a \not\equiv b \pmod{10}$. Then $(a - b) \neq 10k$ for some integer k . Hence $(a - b) \neq 2(5k)$ and $(a - b) \neq 5(2k)$. In other words, $a \not\equiv b \pmod{2}$ and $a \not\equiv b \pmod{5}$. □