

| Clean semileptonic flavor-changing neutral currents that directly test the same Z -coupling distortions many RS variants generate | $C_{9,10}^{(\prime)}$, $C_7^{(\prime)}$, modified Zbs | Nonlocal charm effects and form factors in exclusive modes | LHCb Upgrade and Belle II will sharpen the angular/LFU picture, even though R_K and R_{K^*} are now SM-like. (hflav.web.cern.ch) | | 10 | $B_{s,d} \rightarrow \mu^+ \mu^-$ and related rare leptonic B decays | Very clean helicity-suppressed decays with direct sensitivity to semileptonic/scalar structures | Modified Z couplings, scalar/pseudoscalar operators | Experiment dominates B_d , while B_s is already precision territory | HL-LHC and LHCb Upgrade should make these significantly sharper. (hflav.web.cern.ch) | | 11 | D -mixing and charm CPV | Becomes central in up-aligned warped models, where kaons can be engineered safe but up-sector FCNCs remain | Up-sector KK-gluon/ Z FCNCs, up-sector CP phases, chromomagnetic dipoles | Long-distance charm dynamics still complicate interpretation | LHCb Upgrade, Belle II, and future charm facilities are highly relevant. (hflav-eos.web.cern.ch) | | 12 | Top FCNCs $t \rightarrow cZ$, $t \rightarrow cH$, $t \rightarrow c\gamma$ and production analogues | Clean null tests of the same up-sector flavor misalignment that drives D -physics in many RS variants | Tree-level tqZ/tqH , dipoles, contact terms | Present bounds are still $\mathcal{O}(10^{-4})$ or weaker in BR terms | HL-LHC will be very powerful, especially for tqH and tqZ . (cms-results.web.cern.ch) | | 13 | Direct KK-gluon, VLQ, and custodial-partner searches | These test the flavor model at its dynamical origin rather than through low-energy EFT tails | Direct production of KK gluons, KK electroweak bosons, and vector-like partners | Mass limits are benchmark- and width-dependent | HL-LHC extends reach further; future lepton colliders would reshape the EW side. (annualreviews.org) | | 14 | $B \rightarrow \pi K$ and penguin-dominated CP asymmetries | Good diagnostics of CP phases and electroweak penguins, but weaker as a headline bound | EW penguins, CP phases, RH currents | Hadronic amplitudes remain the bottleneck | Worth keeping in the fit as a correlated diagnostic, not as the main mass setter. (hflav.web.cern.ch) | | 15 | CKM/W-coupling universality, m_W , S , T , U , W , Y | Especially important for custodial/non-custodial architecture decisions and light-fermion compositeness | Gauge-sector mixing, oblique corrections, nonuniversal charged currents | Global-fit correlations matter more than any single number | FCC-ee/CEPC/ILC would dramatically strengthen this whole block. (arxiv.org) |

For clarity, the complete ranking table is reproduced in finished form below.

Rank	Observable	Why it is powerful	Main RS effect	Current limiting uncertainty	Likely future impact
1	ε_K plus operator-level kaon $\Delta S = 2$ bounds	Historically the hardest RS flavor problem; LR operators get RG and chiral enhancement	KK-gluon LR $\Delta S = 2$, new CP phases	Lattice and CKM inputs are under control; the constraint is genuinely severe	Will remain decisive; theory steadily improving via lattice/global fits. (arxiv.org)

Rank	Observable	Why it is powerful	Main RS effect	Current limiting uncertainty	Likely future impact
2	ε'/ε	Unique direct-CPV handle on $\Delta S = 1$ penguins and dipoles, precisely where warped models often generate effects	Z -penguins, EWP, chromomagnetic dipoles	Hadronic matrix elements and isospin breaking	Still likely top-tier; lattice and kaon-theory improvements matter a lot. (pdg.lbl.gov)
3	$B \rightarrow X_s \gamma$ and photon chirality observables	Clean probe of dipoles, which are a natural RS pain point	C_7, C_7' from KK fermion/Higgs/gauge loops	Inclusive rate theory already precise; exclusive polarization limited by hadronic effects	Belle II and LHCb will sharpen chirality tests substantially. (hflav.web.cern.ch)
4	$K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$	Golden FCNC kaon modes with direct sensitivity to modified Z -type currents	Tree-level FCNC $Z/KK - Z/Z'$ exchange	Experiment still dominates, not theory	NA62, KOTO, and eventually KLEVER can become transformational for RS fits. (arxiv.org)
5	$\phi_s, \Delta m_s$ and correlated B_d mixing phases	Strong test of whether alignment suppresses kaons but leaves b -mixing phases	$\Delta B = 2$ from KK gluons and Z -like exchange	Lattice uncertainties smaller than in the past; phase measurements very good	LHCb Upgrade I/II will keep improving this block. (hflav.web.cern.ch)
6	$Z \rightarrow b\bar{b}$ and the EW fit	Classic warped-extra-dimension constraint; directly tests fermion compositeness and custodial protection	Vertex corrections from gauge/fermion KK mixing	Experimental precision already superb; theory fit mature	FCC-ee/CEPC/ILC would be devastatingly powerful, especially for custodial variants. (arxiv.org)

Rank	Observable	Why it is powerful	Main RS effect	Current limiting uncertainty	Likely future impact
7	EDMs (d_e, d_n, d_{Hg})	Best probes of the RS CP problem when anarchic phases are unsuppressed	Quark/electron EDMs, qCEDMs, Weinberg operator	Hadronic/nuclear matching matters, but current limits are already brutal	Next-generation EDM programmes could exclude large regions even if flavor is aligned. (colorado.edu)
8	$\mu N \rightarrow e N$, $\mu \rightarrow e \gamma$, $\mu \rightarrow 3e$	First-line discriminator for any warped model with bulk leptons	LFV dipoles and contact terms from KK exchange	Mostly experimental, not theory	Mu2e, COMET, Mu3e, and MEG II are the cleanest future LB stress test. (arxiv.org)
9	$B \rightarrow K^* \ell \ell$ angular observables and inclusive $B \rightarrow X_s \ell \ell$	Clean semileptonic flavor-changing neutral currents that directly test the same Z -coupling distortions many RS variants generate	$C_{9,10}^{(\ell)}, C_7^{(\ell)}$, modified Zbs	Nonlocal charm effects and form factors in exclusive modes	LHCb Upgrade and Belle II will sharpen the angular/LFU picture, even though R_K and R_{K^*} are now SM-like. (hflav.web.cern.ch)
10	$B_{s,d} \rightarrow \mu^+ \mu^-$ and related rare leptonic B decays	Very clean helicity-suppressed decays with direct sensitivity to semileptonic/scalar structures	Modified Z couplings, scalar/pseudoscalar operators	Experiment dominates B_d , while B_s is already precision territory	HL-LHC and LHCb Upgrade should make these significantly sharper. (hflav.web.cern.ch)

Rank	Observable	Why it is powerful	Main RS effect	Current limiting uncertainty	Likely future impact
11	D -mixing and charm CPV	Becomes central in up-aligned warped models, where kaons can be engineered safe but up-sector FCNCs remain	Up-sector KK-gluon/ Z FCNCs, up-sector CP phases, chromomagnetic dipoles	Long-distance charm dynamics still complicate interpretation	LHCb Upgrade, Belle II, and future charm facilities are highly relevant. (hflav-eos.web.cern.ch)
12	Top FCNCs $t \rightarrow cZ$, $t \rightarrow cH$, $t \rightarrow c\gamma$ and production analogues	Clean null tests of the same up-sector flavor misalignment that drives D -physics in many RS variants	Tree-level tqZ/tqH , dipoles, contact terms	Present bounds are still $\mathcal{O}(10^{-4})$ or weaker in BR terms	HL-LHC will be very powerful, especially for tqH and tqZ . (cms-results.web.cern.ch)
13	Direct KK-gluon, VLQ, and custodial-partner searches	These test the flavor model at its dynamical origin rather than through low-energy EFT tails	Direct production of KK gluons, KK electroweak bosons, and vector-like partners	Mass limits are benchmark- and width-dependent	HL-LHC extends reach further; future lepton colliders would reshape the EW side. (annualreviews.org)
14	$B \rightarrow \pi K$ and penguin-dominated CP asymmetries	Good diagnostics of CP phases and electroweak penguins, but weaker as a headline bound	EW penguins, CP phases, RH currents	Hadronic amplitudes remain the bottleneck	Worth keeping in the fit as a correlated diagnostic, not as the main mass setter. (hflav.web.cern.ch)

Rank	Observable	Why it is powerful	Main RS effect	Current limiting uncertainty	Likely future impact
15	CKM/W-coupling universality, m_W, S, T, U, W, Y	Especially important for custodial/non-custodial architecture decisions and light-fermion compositeness	Gauge-sector mixing, oblique corrections, nonuniversal charged currents	Global-fit correlations matter more than any single number	FCC-ee/CEPC/ILC would dramatically strengthen this whole block. (arxiv.org)

Seed-list audit

Seed item	Verdict	Modern notation / correction	Closely related observables that should also be included
ε'/ε in $K \rightarrow \pi\pi$	Yes, absolutely relevant	No notation change needed	$K \rightarrow \pi\nu\bar{\nu}, K_{L,S} \rightarrow \pi^0\ell\bar{\ell}, K_S \rightarrow \mu^+\mu^-,$ radiative kaon modes. (pdg.lbl.gov)
$K^+ \rightarrow \pi^+\nu\bar{\nu}$	Yes, critical	No notation change needed	$K_L \rightarrow \pi^0\nu\bar{\nu},$ future KLEVER, CKM-clean kaon NP fits. (arxiv.org)
$K_L \rightarrow \pi^0\nu\bar{\nu}$	Yes, critical	No notation change needed	$K^+ \rightarrow \pi^+\nu\bar{\nu}, K_L \rightarrow \pi^0 X^0$ invisible modes in KOTO analyses. (arxiv.org)
$K_L \rightarrow \mu^+\mu^-$	Relevant, but not clean enough to stand alone	No notation change needed	$K_L \rightarrow e^+e^-, K_{L,S} \rightarrow \pi^0\ell^+\ell^-, K_S \rightarrow \mu^+\mu^-.$ (pdg.lbl.gov)
$K_L \rightarrow \pi^0\ell^+\ell^-$	Yes	Keep separate e and μ channels	Include the $K_S \rightarrow \pi^0\ell^+\ell^-$ control modes because they feed the interpretation. (pdg.lbl.gov)
$b \rightarrow s\gamma$	Yes, critical	Use the full modern block: $B \rightarrow X_s\gamma, B \rightarrow K^*\gamma, B_s \rightarrow \phi\gamma,$ CP and polarization observables	Add $b \rightarrow d\gamma, C_7, C_7'$ fits, and time-dependent radiative CP asymmetries. (hflav.web.cern.ch)
$B \rightarrow X_s\ell^+\ell^-$	Yes	Use inclusive low- q^2 and high- q^2 bins separately	Correlate with $B \rightarrow K^{(*)}\ell\bar{\ell}, B_s \rightarrow \phi\ell\bar{\ell},$ LFU ratios. (hflav.web.cern.ch)

Seed item	Verdict	Modern notation / correction	Closely related observables that should also be included
$B \rightarrow K^* \ell^+ \ell^-$	Yes, critical	Modern fit language is angular observables $P_i^{(\prime)}, F_L, A_{\text{FB}}, S_i, A_i$ in standard q^2 bins	Add $B_s \rightarrow \phi \ell \ell, \Lambda_b \rightarrow \Lambda \ell \ell$, and inclusive $X_s \ell \ell$. (hflav.web.cern.ch)
$B^+ \rightarrow \tau^+ \nu_\tau$	Relevant but not first-tier in generic anarchic RS	Usually fit together with $B^+ \rightarrow \mu^+ \nu, R(D), R(D^*),$ and $B_c \rightarrow \tau \nu$ consistency	Most useful if your RS setup alters W couplings. (hflav.web.cern.ch)
$B \rightarrow \pi K$	Relevant as a diagnostic	Use full πK rate and CP-asymmetry system, not one number	Also include penguin CP-eigenstates such as $\phi K_S, \eta' K_S, \pi^0 K_S$. (hflav.web.cern.ch)
$B \rightarrow \phi K_S / B_s \rightarrow \phi \phi$	Yes	Use modern notation $B^0 \rightarrow \phi K_S^0; B_s \rightarrow \phi \phi$ is fine	Add $B_s \rightarrow K^{*0} \bar{K}^{*0}$ and the broader penguin S_f set. (hflav-eos.web.cern.ch)
$t \rightarrow c Z$	Yes, especially for up alignment	Modern fits usually bundle $t \rightarrow q Z$ with single-top FCNC production	Add $t \rightarrow q H, t \rightarrow q \gamma, tqg$ production. (cms-results.web.cern.ch)
$Z \rightarrow b \bar{b}$	Yes, classic RS observable	Use the complete set $R_b, A_b, A_{\text{FB}}^b, \delta g_{Lb}, \delta g_{Rb}$ rather than only one rate	Also include $Z \rightarrow c \bar{c}, \Gamma_Z, S, T, U, W, Y$, and m_W in any custodial analysis. (arxiv.org)

A notation point worth flagging explicitly is your request about older “ $K_1 \rightarrow \pi \gamma$ ” language. In modern phenomenology that is **not** a standard standalone fit observable. Old K_1 notation refers to CP eigenstates in the CP limit, roughly corresponding to K_S and K_L , while the modern radiative kaon fit basis is built from channels such as $K \rightarrow \pi \ell \ell, K \rightarrow \pi \pi \gamma, K \rightarrow \pi \pi e^+ e^-,$ and $K_{L,S} \rightarrow \pi^0 \gamma \gamma$. In other words, treat “ $K_1 \rightarrow \pi \gamma$ ” as legacy shorthand to be translated into the modern radiative/virtual-photon kaon modes, not as an observable by itself. (pdg.lbl.gov)

Potentially missing observables

Overlooked observable	Why it matters specifically for RS
$K_S \rightarrow \mu^+ \mu^-$	It is one of the cleanest “forgotten” kaon probes of short-distance $s \rightarrow d \ell \ell$ structure and CP composition, and it is directly relevant when modified $Z d s$ couplings are present. (pdg.lbl.gov)

Overlooked observable	Why it matters specifically for RS
Semileptonic asymmetries $a_{\text{sl}}^{d,s}$	They are not the headline constraints, but they probe phases in $\Delta B = 2$ that are not identical to those seen in $S_{\psi K_S}$ or ϕ_s . In aligned or CP-protected models they are excellent cross-checks. (hflav.web.cern.ch)
Photon polarization in radiative B decay	RS often produces sizeable right-handed dipoles , so the polarization observables are more diagnostic than the branching ratio alone. (eos.web.cern.ch)
$\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ and baryonic LFU ratios	These give complementary helicity information in $b \rightarrow s \ell \ell$ and can help disentangle operator structure if the mesonic modes stay theory-limited in some bins. (hflav.web.cern.ch)
CKM first-row unitarity plus $\pi/K/\tau/W/Z$ universality	These are easy to miss if one thinks of RS only as a FCNC model, but they matter whenever fermion localization induces nonuniversal charged-current couplings . (hflav.web.cern.ch)
$h \rightarrow t\bar{c}$ and $t \rightarrow cH$ as a correlated pair	In up-aligned or partially protected up-sector scenarios, these are among the few genuinely clean tree-level FCNC LHC probes. (cms-results.web.cern.ch)
$B_s \rightarrow \tau^+ \tau^-$ and $B \rightarrow K^{(*)} \tau^+ \tau^-$	These are not yet leading constraints, but they become important if your warped setup features nontrivial third-generation or bulk-lepton structure. (hflav.web.cern.ch)
High- p_T dilepton/dijet tails	Once the KK spectrum moves high enough that resonances are broad or inaccessible on-shell, contact-interaction tails become the cleaner direct collider handle on the same underlying physics. (cms-results.web.cern.ch)
EDMs in a nominally “flavor” fit	Warped models with anarchic phases often fail first in CP-odd dipoles , not in a traditional flavor branching ratio. Leaving EDMs out can give a very misleading sense of viability. (colorado.edu)

Not useful, theory-limited, or redundant

Observable or class	Why it should not drive the fit
Δm_K as a standalone “measured minus SM” bound	The measurement is exquisite, but the SM long-distance contribution is not clean enough to turn Δm_K alone into the anchor NP constraint. Use the operator-level $\Delta S = 2$ bounds instead. (arxiv.org)
y_D by itself as a new-physics observable	y_D is dominated by long-distance SM physics, so it is not the cleanest NP discriminator. In warped fits, x_D , $ q/p $, ϕ_D , and direct charm CPV are usually more informative. (hflav-eos.web.cern.ch)
$K_L \rightarrow \mu^+ \mu^-$ or $K_L \rightarrow e^+ e^-$ as primary constraints	They are relevant, but long-distance two-photon physics makes them poor headline bounds compared with $K \rightarrow \pi \nu \bar{\nu}$ or ε'/ε . (pdg.lbl.gov)

Observable or class	Why it should not drive the fit
$B \rightarrow \pi K, B \rightarrow \pi\pi, B \rightarrow KK$ as hard mass-bound setters	They are excellent diagnostics of weak phases and electroweak penguins, but hadronic amplitudes remain too uncertain for them to dominate a serious RS exclusion. (hflav.web.cern.ch)
Charm rare decays integrated over all phase space	Modes like $D \rightarrow \pi\ell\ell$ or $D \rightarrow V\gamma$ are often long-distance dominated, so they are most useful only in carefully chosen short-distance windows or as null tests. (cds.cern.ch)
$h \rightarrow bs, bd, sd$ as present-day practical constraints	They are theoretically very relevant to warped Higgs flavor violation, but the current direct experimental reach is still far from what low-energy flavor and top-Higgs FCNC already constrain indirectly. (inspirehep.net)
Old narrow-benchmark resonance limits used without reinterpretation	RS direct-search bounds depend strongly on widths, couplings, and whether the state is a KK gluon, KK graviton, radion, or custodial fermion. Use modern benchmark-specific or EFT-style reinterpretations, not a single historical mass number. (annualreviews.org)
R_K and R_{K^*} treated as an “anomaly prior”	They remain important observables, but in a 2026 RS fit they should be treated as precision semileptonic inputs that are now broadly SM-compatible, not as a standing discrepancy. (hflav.web.cern.ch)

Open questions and limitations

This catalog is deliberately **family-complete** rather than **per-bin exhaustive**. For giant observable blocks — the canonical example is the standard suite of q^2 -binned $B \rightarrow K^*\mu^+\mu^-$ angular observables — the table identifies the block, its present experimental status, the RS operator mapping, and the source from which the full bin-by-bin numerical content should be imported into a later machine-readable fit. That is the right granularity for a modern RS analysis plan, even though it is not a substitute for the final covariance-matrix implementation. (hflav.web.cern.ch)

A second limitation is that some **secondary electroweak and direct-search entries** are inherently benchmark dependent. For instance, there is no single universal “KK-gluon mass limit” independent of width, localization pattern, or decay branching fractions; similarly, the usefulness of CLFV, LFU, Higgs-LFV, and Z -LFV observables depends strongly on whether the model includes warped bulk leptons. I have therefore marked these rows accordingly and emphasized where they become first-tier only in specific RS variants. (annualreviews.org)

Bibliography

Heavy-flavor averages and fit infrastructure

Heavy Flavor Averaging Group, *Averages of b -hadron, c -hadron, and tau-lepton properties as of 2023* and the continuously updated online result tables, including the 2026 executive summary, the Winter 2024 UT-angle page, the CKM25 charm-mixing tables, and the 2023 τ -LFV combinations. These are the primary sources for $\Delta m_{d,s}, \phi_s, R(D^{(*)}), B \rightarrow X_s\gamma, R_K, R_{K^*}$, charm mixing, direct charm CPV, τ -universality, and τ -LFV averages. (hflav.web.cern.ch)

Y. Aoki et al., FLAG, *Review of lattice results concerning low-energy particle physics*, 2024 update. This is the key lattice source for kaon and B -mixing bag parameters, decay constants, and semileptonic inputs needed to translate RS Wilson coefficients into observables. ([arxiv.org](https://arxiv.org/abs/2403.17867))

M. Bona et al., UTFit-style model-independent $\Delta F = 2$ new-physics analyses, especially the 2022 update on neutral-meson mixing. Use these for the operator-by-operator bounds that are more meaningful in RS than simply quoting a single meson-mixing observable. ([arxiv.org](https://arxiv.org/abs/2203.04863))

Kaons and rare-decay reviews

Particle Data Group, 2024/2025 rare-kaon and conservation-law reviews. These remain the best compact review sources for ε'/ε , $K_{L,S} \rightarrow \pi^0 \ell^+ \ell^-$, $K_{L,S} \rightarrow \ell^+ \ell^-$, radiative modes, and LFV kaons. (pdg.lbl.gov)

X. Chang for NA62, *New measurement of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ branching ratio at the NA62 experiment*, 2026 preliminary result based on the 2016–2024 dataset, giving the current most useful status for the charged golden mode in kaon flavor. ([arxiv.org](https://arxiv.org/abs/2403.17867))

J. K. Ahn et al. for KOTO, *Search for the $K_L \rightarrow \pi^0 \nu \bar{\nu}$ decay at the J-PARC KOTO experiment*, PRL 134 (2025), the current benchmark result for the neutral golden mode. ([arxiv.org](https://arxiv.org/abs/2403.17867))

Charged-lepton flavor violation and EDMs

PDG conservation-laws review, 2024. Use this as the compact baseline source for the current best limits on $\mu \rightarrow 3e$, $\mu N \rightarrow e N$, K -LFV, and many older but still current LFV channels. (pdg.lbl.gov)

MEG II Collaboration, 2025 $\mu \rightarrow e \gamma$ result, the current flagship radiative muon-LFV constraint. ([arxiv.org](https://arxiv.org/abs/2403.17867))

Mu2e official status/projection material and COMET Phase-I documentation. These are the appropriate sources for future $\mu N \rightarrow e N$ conversion sensitivity in any lepton-bulk RS scenario. (fnal.gov)

JILA Cornell group, current HfF^+ electron-EDM programme page, describing the present best d_e sensitivity. (colorado.edu)

C. Abel et al., *Measurement of the permanent electric dipole moment of the neutron*, PRL 124 (2020), still the definitive neutron-EDM constraint used in generic CP-violating RS analyses. (cds.cern.ch)

W. B. Cairncross et al., *Precision measurement of the electron's electric dipole moment using trapped molecular ions*, and B. Graner et al., *Reduced limit on the permanent electric dipole moment of ^{199}Hg* , for the molecular and atomic EDM sectors. ([arxiv.org](https://arxiv.org/abs/2403.17867))

Top FCNC, Higgs-LFV, and collider limits

CMS public FCNC top-physics publication pages and preliminary top pages, together with recent ATLAS proceedings on $t \rightarrow qH$ and other top-FCNC channels, are the right sources for current limits on $t \rightarrow qZ$, $t \rightarrow q\gamma$, $t \rightarrow qH$, and the corresponding single-top FCNC production searches. (cms-results.web.cern.ch)

ATLAS and CMS direct Higgs-LFV searches — especially the recent ATLAS $h \rightarrow e\tau, \mu\tau$ Run-2 paper and the CMS Run-2 $h \rightarrow \mu\tau, e\tau$ paper — are the correct present sources for LFV Higgs constraints in bulk-lepton warped models. (arxiv.org)

B. Clerbaux and collaborators, *Searches for New High-Mass Resonances Decaying to Fermions* in *Annual Review of Nuclear and Particle Science* and the CMS review of vector-like quark searches. These are the most useful compact entry points for direct RS collider constraints, especially when you need to map benchmark-specific mass limits to KK gluons, KK gravitons, KK electroweak bosons, and custodial fermions. (annualreviews.org)

ATLAS and CMS, *Highlights of HL-LHC physics projections*, 2025. This is the right umbrella source for future impact statements on top FCNC, Higgs precision, EW precision, and some direct-search reach. (arxiv.org)

RS-specific theory anchors

K. Agashe, G. Perez, A. Soni, *Flavor from Minimal Flavor Violation and a Viable Randall-Sundrum Model*, arXiv: 0710.1869. This is one of the theory anchors for the MFV/RS-GIM direction you explicitly want to revisit.

C. Csáki, A. Falkowski, A. Weiler, *Flavor Alignment via Shining in RS*, arXiv:0907.0474. This is the theory anchor for the down-aligned/shining branch of the inventory.

A. J. Buras, M. Harlander, A. Hryczuk, J. Kumar, and collaborators' recent rare-decay/global-fit reviews and modern EFT mapping papers are the right bridge between today's experimental inputs and the Wilson-coefficient basis in which warped flavor models are most naturally constrained. (link.springer.com)
