

Three CP invariants, not one: what actually has to be aligned to fix ε_K in warped flavor

A survival-mechanism anatomy of anarchic Randall-Sundrum, with a numerically demonstrated negative result

5D-Neutrino-Mixing project, RS-FLAVOR-ALIGNMENT-2026-07

July 2026

Abstract

Anarchic Randall-Sundrum (RS) is pushed to $M_{KK} \gtrsim 10\text{--}20$ TeV almost entirely by the CP-violating kaon amplitude ε_K , generated by the chirality-enhanced KK-gluon operator C_4 . The standard responses align the Yukawas so as to suppress the *magnitude* of the dangerous coupling (right-handed down $U(3)$, 5D-MFV, flavor shining). We ask a sharper question: what is the minimal set of CP structures that a low- M_{KK} RS point must control, and does aligning the kaon operator suffice. Working from a forward Monte Carlo that retains the full complex down-sector couplings, we establish three things. (i) Anarchic ε_K survival at $M_{KK} \leq 3$ TeV is *magnitude-dominated*: about 57% of survivors have an accidentally small $|C_4|$ and only about 7% survive by a genuine CP-phase cancellation. This is a consequence of the cut itself, which weights survivors by a phase-volume factor $P_{\text{pass}}(R_K) \sim 1/R_K$. (ii) A “CP-only” alignment that makes the down Yukawa real kills the kaon phase exactly and passes ε_K at 100% *while keeping the coupling magnitude anarchic*, so Δm_K stays large. This is a distinct corner from the magnitude cures. (iii) That same point is *not* EDM-safe: the neutron-EDM CP invariant $I_d = \text{Im}[F_Q(Y_u Y_u^\dagger + Y_d Y_d^\dagger)Y_d F_d]_{11}$ stays $O(1)$, sourced by up-sector CP through $Y_u Y_u^\dagger$, independent of the kaon phase. The kaon operator and the neutron EDM are governed by *different* CP invariants; aligning one does not align the other. The practical consequence is that “sequester CP” is not one condition but (at least) two, and the $(\Delta m_K, d_n)$ plane separates every candidate mechanism. The genuinely open, buildable target is a calculable construction that suppresses the kaon operator *magnitude* (rank-one / $U(2)$ localization) *and* confines CP in the strong Nelson-Barr sense (real determinant), because neither alone is sufficient.

1 The obstruction, stated precisely

The RS-GIM mechanism suppresses most quark flavor violation once M_{KK} is electroweak-safe, with the single robust exception of ε_K [1, 2], and a subdominant D^0 concern. In this repo’s convention the binding operator is the color-octet KK-gluon left-right operator,

$$C_4 = -\frac{(G_L^d)_{12}(G_R^d)_{12}}{M_{KK}^2}, \quad (G_L^d)_{12} \sim g_{s*} f_{Q_1} f_{Q_2} e^{i\phi_L}, \quad (G_R^d)_{12} \sim g_{s*} f_{d_1} f_{d_2} e^{i\phi_R}, \quad (1)$$

with $\phi_{L,R}$ order-one anarchic phases. Writing $R_K \equiv 2|M_{12}^{LR}|/\Delta m_K$ and $\theta_K \equiv \arg[(G_L^d)_{12}(G_R^d)_{12}]$,

$$|\varepsilon_K^{\text{NP}}| \simeq \frac{\kappa_\epsilon}{2\sqrt{2}} R_K |\sin \theta_K| \simeq 0.332 R_K |\sin \theta_K|, \quad \Delta m_K^{\text{NP}} \propto R_K \cos \theta_K. \quad (2)$$

So ε_K measures $R_K \sin \theta_K$ and Δm_K measures $R_K \cos \theta_K$: the same magnitude R_K , projected on the imaginary and real axes. A low- M_{KK} survivor can therefore be built two ways, by small R_K (small magnitude) or by small $\sin \theta_K$ (aligned phase). These are physically different and, as we show, they predict different things.




Figure 1: Anarchic ε_K survival in the (magnitude, phase) plane at $M_{KK} = 3 \text{ TeV}$. Survivors (red) populate two disjoint regions: the left edge (small $|C_4|$, magnitude-suppressed) and the bottom edge (small $|\sin \Phi_{12}|$, phase-aligned). The bulk is the magnitude edge.

2 F1: anarchic survival is magnitude-dominated

From a 9.9×10^5 -draw anarchic ensemble (masses and CKM fit per draw), the ε_K survivors at $M_{KK} \leq 3 \text{ TeV}$ split (Fig. 1) into *magnitude-suppressed* ($\sim 57\%$, small $|C_4|$, which also makes Δm_K small) and *phase-aligned* ($\sim 7\%$, $|C_4|$ anarchic, $\sin \theta_K \rightarrow 0$, Δm_K saturating). The dominance of the magnitude mode is not an independent fact: the ε_K cut weights the pre-cut population by the phase volume

$$P_{\text{pass}}(R_K) = \frac{2}{\pi} \arcsin\left(\frac{\epsilon_{\text{max}}}{0.332 R_K}\right) \xrightarrow{R_K \gg} \frac{2 \epsilon_{\text{max}}}{\pi 0.332} \frac{1}{R_K}, \quad (3)$$

so large-magnitude points are penalized as $1/R_K$ and survivors tilt to small $|C_4|$. The measured correlation $\text{corr}(\log \varepsilon_K, \log \Delta m_K) = +0.82$ is the signature: a pure phase-cancellation picture would give an anticorrelation. The practical lesson is that “anarchy survives by fine-tuning the ε_K phase” is the *minority* path; most survivors simply draw a small coupling.

3 The three CP invariants

The point of this note is that a low- M_{KK} RS quark sector must control more than one CP structure, and they are independent:

- $\theta_K = \arg[(G_L^d)_{12}(G_R^d)_{12}]$ sets ε_K (kaon 1-2, left-right).
- $I_d = \text{Im}[F_Q(Y_u Y_u^\dagger + Y_d Y_d^\dagger)Y_d F_d]_{11}/(\text{mass})$ sets the neutron EDM. The tree KK-gluon EDM vanishes (the diagonal couplings are real); the leading RS contribution is the one-loop Higgs/KK-fermion dipole with the *misaligned* spurion $Y_u Y_u^\dagger + Y_d Y_d^\dagger$ [3].
- $\theta_D = \arg[(G_L^u)_{12}(G_R^u)_{12}]$ sets D^0 CP violation (up 1-2).

The crucial feature is that I_d feeds on $Y_u Y_u^\dagger$: up-sector CP enters the down-quark EDM directly, with no reference to the kaon phase θ_K .




Figure 2: The $(\Delta m_K, d_n)$ discriminator at $M_{KK} = 3$ TeV. Magnitude alignment moves left (small $|C_4|$, Δm_K quiet); CP-only alignment stays right (Δm_K loud) but high in $|I_d|$ (EDM loud). The green EDM-quiet, Δm_K -loud corner (genuine Nelson-Barr) is not reached by any single-axis anarchic draw.

4 F2/F3: CP-only alignment fixes ε_K but not the EDM

Consider the cleanest CP-only alignment: a real anarchic down Yukawa, with CP carried by the up sector (a Nelson-Barr-like routing that keeps a realistic Jarlskog J). By construction the down M_d is real, so $\theta_K = 0$ and ε_K vanishes at tree level, while $|C_4|$ is untouched. The forward Monte Carlo confirms this and adds the decisive point (Table 1, Fig. 2):

Table 1: Mechanism comparison at $M_{KK} = 3$ TeV (PDG+ J -passing points). $|C_4|$ is the Δm_K proxy; $|I_d|$ is the neutron-EDM CP invariant.

Lane	ε_K pass	$ \sin \theta_K $	$\log_{10} C_4 $	$ I_d $ (EDM)
flat anarchy	2%	0.54	-15.9	2.9
magnitude align ($U(3)_{d_R}$)	high	—	suppressed	suppressed
CP-only (real Y_d)	100%	10^{-16}	-15.9	4.1
sanity: fully real $Y_{u,d}$	—	10^{-16}	—	0.0

A real down sector passes ε_K at 100% with the coupling magnitude at its anarchic value (so Δm_K stays large), yet the EDM invariant is $|I_d| \simeq 4$, *larger* than flat anarchy. Turning up-sector CP off (fully real Yukawas) sends $I_d \rightarrow 0$ exactly, confirming both the estimator and the mechanism: the residual EDM is up-sector CP leaking through $Y_u Y_u^\dagger$. Aligning the kaon operator does nothing to it. This is the central result: **CP-only alignment of the kaon operator does not yield an EDM-safe theory**. An EDM cure requires controlling the up-sector CP as well, which is precisely what genuine Nelson-Barr (a real quark mass determinant, CP confined to a heavy sector) delivers and a merely real Y_d does not.

5 What is new, and what is not

The broad ideas here are not new and we do not claim them. CP-only / minimal-CP suppression of ε_K with reduced EDMs exists in composite/warped language (Redi-Weiler [8]); warped

Nelson-Barr with CP moved to a heavy sector exists (Girmohanta et al. [9]); the magnitude cures are FPR 5D-MFV [4], Santiago right-handed-down protection [5], and flavor shining [6]; CP sequestering in warped space is Cheung-Fitzpatrick-Randall [7]. The defensible, and to our knowledge unpublished, content is narrower:

1. the survival-mechanism taxonomy of Sec. 2, with the $P_{\text{pass}} \sim 1/R_K$ explanation of why anarchic survival is magnitude-dominated;
2. the explicit demonstration that the kaon phase and the neutron-EDM invariant are independent, so that a real down sector is ε_K -safe with anarchic Δm_K yet EDM-loud (Sec. 5), i.e. CP-only kaon alignment is provably insufficient;
3. the $(\Delta m_K, d_n)$ plane as a mechanism discriminator that separates magnitude alignment, CP-only alignment, and genuine Nelson-Barr into distinct, falsifiable corners.

6 The worked target

The synthesis the literature points to, and that our numerics motivate operationally, is a single calculable construction combining two independent ingredients, because we have shown neither suffices alone:

- **Magnitude:** a sequentially generated, low-rank Yukawa ensemble under an accidental light-family $U(2)$ symmetry (rather than i.i.d. anarchy), so that the KK-gluon 1-2 coupling is structurally, not accidentally, small. This is the “wrong ensemble” correction: natural Yukawas are rank-one towers, not flat random matrices, and the resulting ε_K floor is set by symmetry-breaking leakage rather than phase luck.
- **CP:** genuine Nelson-Barr confinement of CP (real mass determinant), so that both θ_K and I_d are spurion-suppressed, not just the kaon phase. A real Y_d alone is demonstrably not enough.

Each ingredient exists separately; no calculable 5D construction combines them with a controlled radiative-stability proof. The decisive metric for such a build is the scale M_{50} at which at least half of the points pass *all* constraints (masses, CKM, ε_K , Δm_K , $B_{d,s}$, D^0 , and the EDM invariant), compared across flat anarchy, magnitude-only, CP-only, and the combined construction. Our results predict that only the combined construction reaches the few-TeV regime with Δm_K retained and I_d suppressed.

7 Bottom line

The RS ε_K problem has been read as a problem of one delicate phase in the off-diagonal down couplings. It is really a problem of *three* independent CP structures, of which the kaon phase is only one. Aligning the kaon operator, by magnitude or by phase, leaves the neutron EDM untouched, because the EDM samples a flavor-diagonal invariant fed by the up sector. A low- M_{KK} warped flavor model is therefore not “anarchy plus one alignment relation”; it must suppress the kaon operator magnitude with a low-rank, $U(2)$ -symmetric ensemble and confine CP in the Nelson-Barr sense, together, in one calculable construction. That combined target, and the $(\Delta m_K, d_n)$ discriminator that certifies it, is the concrete next step.

References

- [1] C. Csáki, A. Falkowski, A. Weiler, arXiv:0804.1954.
- [2] M. Blanke et al., arXiv:0809.1073.

- [3] K. Agashe, G. Perez, A. Soni, hep-ph/0408134 (RS CP problem / neutron EDM).
- [4] A. L. Fitzpatrick, G. Perez, L. Randall, arXiv:0710.1869 (5D-MFV).
- [5] J. Santiago, arXiv:0806.1230 (minimal flavor protection).
- [6] C. Csáki, G. Perez, Z. Surujon, A. Weiler, arXiv:0907.0474 (flavor shining).
- [7] C. Cheung, A. L. Fitzpatrick, L. Randall, arXiv:0711.4421 (CP sequestering).
- [8] M. Redi, A. Weiler, arXiv:1106.6357 (minimal CP violation).
- [9] S. Girmohanta, S. J. Lee, Y. Nakai, M. Suzuki, arXiv:2203.09002 (warped Nelson-Barr).
- [10] A. Greljo, A. E. Thomsen (rank-one sequential Yukawas); Barbieri et al. $U(2)$ flavor.