
Reproducing RS Warped Flavor & Electroweak Constraints:

Fixed-Code Validation Against the Literature

Minimal Randall–Sundrum, quark sector.
Audit fixes + reproduction of published figures.

June 2026

1 Introduction

We work in a minimal Randall–Sundrum (RS) warped extra dimension; fermion mass hierarchies come from geometric (wavefunction) localization rather than hierarchical 5D Yukawas. An adversarial audit of our quark-sector constraint code found six implementation errors (three blocker-level), now fixed and dual-reviewed: **B1** the $Z \rightarrow b\bar{b}$ fermion-KK admixture (mistranslated from Casagrande et al. 0807.4937 — wrong sign and $\sim 190\times$ too large), **B2** the ε_K SVD→PDG rephasing (ε_K was convention-dependent), **B3** the $\Delta F = 2$ O_4/O_5 chiral coefficients + state normalization, **M1** the $Z \rightarrow b\bar{b}$ (T010) gate, **M2** the EW001 ΔT convention (oblique T was $\sim 6\times$ too weak), and **M5** a tagging bug. Net effect: the headline minimal-model floor changed qualitatively — the old “25–30 TeV (108 TeV at 1σ) $Z \rightarrow b\bar{b}$ floor” was an artifact of B1 and collapses to ~ 5 TeV once corrected.

We validate the fixed code two ways. (i) A **constrained-fit scan**: draw bulk masses + a seed, fit the Yukawas to reproduce the measured quark masses + CKM (a near-unique solution $\rightarrow$ a thin “locus”). (ii) An **anarchic-forward scan**: draw complex $O(1)$ anarchic Yukawas, compute observables forward, keep mass+CKM-reproducing points (the multi-decade “cloud” the literature uses). The papers scatter anarchic Yukawas; our production scan fits — so the two give very different spreads, which matters for how floors compare to the literature.

Headline findings

- The fixed code **reproduces the published RS literature** when run in anarchic-forward mode: ε_K 95%-quantile floor = **10 TeV** (= Bauer 0912.1625 headline), D^0 funnel containment 99.6% at 10 TeV (Gedalia), kaon $\text{Im}/\text{Re}(M_{12}) = \mathbf{106\times}$ ($\approx$ Blanke’s $\sim 100\times \varepsilon_K$ problem), $C_{B_d}/C_{B_s} \approx 1$.
- **Minimal-model floors (corrected):** *typical* anarchic floor with current data $\approx$ **30 TeV**, set by ε_K (flavor); *existence* (fine-tuned) floor $\approx$ **18–20 TeV**, set irreducibly by **S, T, U**. $Z \rightarrow b\bar{b}$ no longer leads (≈ 5 TeV).
- Flavor constraints are **tunable** (alignment drives the NP $\rightarrow 0$); **S, T, U is irreducible** (no Yukawa freedom). This is precisely why custodial RS exists.

2 Floor summary: typical vs. existence

The two scan modes give two distinct notions of a floor. The *typical* (median) floor is where the bulk of anarchic points become excluded; the *existence* (best/tuned) floor is where even the single best-aligned point crosses the bound. Flavor constraints separate the two by orders of magnitude (they are tunable); the oblique S, T, U constraint does not (it has no Yukawa freedom).

Constraint	Typical (median)	Existence (best/tuned)	Tunable?
ε_K	~ 30 TeV (anarchic)	$\lesssim 1$ TeV	yes (align $\text{Im } M_{12} \rightarrow 0$)
$\Delta m_d (B_d)$	few TeV	$\lesssim 1$ TeV	yes
$\Delta m_s (B_s)$	few TeV	$\lesssim 1$ TeV	yes
D^0	low	$\lesssim 1$ TeV	yes
Δm_K	low	$\lesssim 3$ TeV	yes
$Z \rightarrow b\bar{b}$ (T010)	~ 5 TeV	~ 4.6 TeV	no (gauge-dominated; c_{Q_3} pinned by m_t)
S, T, U (EW001)	~ 20 TeV	$\sim 18\text{--}20$ TeV	no (no Yukawa dependence)
COMBINED	~ 30 TeV (ε_K)	$\sim 18\text{--}20$ TeV (S, T, U)	—

3 Figures: reproduction and analysis

Constraint census — our 100k minimal scan (ours only)




Figure 1: Per-constraint veto fraction vs. M_{KK} from our 100k minimal-model scan with all fixes in.

Per-constraint veto fraction vs. M_{KK} from our 100k minimal-model scan with all fixes in. The corrected ranking: S, T, U (**EW001**) leads at ~ 20 TeV, ϵ_K (rigorous) at 7 TeV, $Z \rightarrow b\bar{b}$ (**T010**) collapsed to 5 TeV (it was the spurious ~ 25 – 30 TeV B1 artifact), Δm_s at 2 TeV; $D^0/\Delta m_d/Z \rightarrow b\bar{b}-A_b$ never bind. Dropping $Z \rightarrow b\bar{b}$ leaves the combined floor unchanged at 20 TeV — concrete proof it is no longer the driver.

S, T, U oblique — Casagrande et al. 0807.4937, Fig. 4 (paper left, ours right)




Figure 2: The S – T plane: paper (left) vs. ours (right). Minimal RS sweeps a wedge over M_{KK} ($1 \rightarrow 10$ TeV) and warp volume L against the CL ellipses.

The paper plots the S – T plane with 68/95/99% CL ellipses and the minimal-RS “wedge” swept over M_{KK} ($1 \rightarrow 10$ TeV) and warp volume L . Minimal RS climbs steeply in $\mathbf{T}$ — the volume-

ε_K — Bauer et al. 0912.1625, Fig. 4 (paper left, ours right)




Figure 4: Anarchy $|\varepsilon_K|$ vs. M_{KK} : paper (left) vs. ours (right), with 5/50/95% quantile curves. Paper scatters anarchic Yukawas; we reproduce the fat cloud in anarchy-forward mode.

The canonical RS ε_K problem: anarchic $|\varepsilon_K|$ vs. M_{KK} spanning ~ 6 decades, colored by $Z \rightarrow b\bar{b}$ / ε_K consistency, with 5/50/95% quantile curves. The key methodological point: the paper **scatters** anarchic $O(1)$ Yukawas (fat cloud) while our production scan **fits** to masses+CKM (thin locus). Running our fixed code (B2 rephasing + B3 coefficients) in anarchy-forward mode reproduces the fat cloud ($\sim 3+$ decades) and lands the **95%-quantile floor at $M_{KK} = 10$ TeV** — **exactly Bauer’s quoted “ $M_g^{(1)} \gtrsim 10$ TeV.”** With current (vs. paper-era) constraints the $\geq 50\%$ floor rises to ~ 30 TeV.

Consistency fraction — Bauer et al. 0912.1625, Fig. 5 (paper left, ours right)




Figure 5: Percentage of anarchic points consistent vs. M_{KK} : paper (left) vs. ours (right), for both paper-era and current inputs.

Percentage of anarchic points consistent vs. M_{KK} : a rising S-curve with $1/M_{KK}^2$ decoupling, where $Z \rightarrow b\bar{b}$ consistency turns on before the combined fraction. Ours reproduces the same ordering and decoupling, shown for both paper-era and current inputs.

B_d mixing — Bauer et al. 0912.1625, Fig. 6 (paper left, ours right)




Figure 6: The B_d -mixing NP plane (C_{B_d} vs. ϕ_{B_d}): paper (left) vs. ours (right), centered near the SM point (1, 0).

The B_d -mixing new-physics plane: amplitude C_{B_d} vs. phase ϕ_{B_d} . The paper’s anarchic cloud sits near the SM point $(1, 0)$ with the experimental CL ellipse offset to $\approx -6^\circ$. Our anarchic cloud centers on the SM cross with few-degree phases — $C_{B_d}[p5, 50, 95] = [0.97, 1.00, 1.06]$, $\phi_{B_d} \approx \pm 1^\circ$ — matching Bauer’s structure (their $C_{B_d} = 0.89 \pm 0.17$); our spread is narrower because of our $O(1)$ Yukawa window.

B_s mixing — Bauer et al. 0912.1625, Fig. 7 (paper left, ours right)




Figure 7: The analogous B_s plane (C_{B_s} , ϕ_{B_s} , $S_{\psi\phi}$): paper (left) vs. ours (right).

The analogous B_s plane (C_{B_s} , ϕ_{B_s} , and $S_{\psi\phi}$). Ours: $C_{B_s} \approx [0.97, 1.01, 1.20]$, $\phi_{B_s} \approx \pm 2^\circ$, $S_{\psi\phi}$ median 0.04 (= SM) — matching Bauer ($C_{B_s} = 0.93 \pm 0.19$).

D^0 mixing — Gedalia, Grossman, Nir, Perez 0906.1879, Fig. 1 (paper left, ours right)




Figure 8: The allowed “funnel” in $(x_{12}^{\text{NP}}/x, \sin 2\sigma_D)$: paper (left) vs. ours (right). Our cloud sinks into the allowed funnel as M_{KK} rises.

The allowed “funnel” in $(x_{12}^{\text{NP}}/x, \sin 2\sigma_D)$. Our anarchic D -meson cloud (now carrying the complex M_{12} phase) sinks into the grey allowed funnel as M_{KK} rises — inside-funnel fraction 57.5% at 1 TeV $\rightarrow$ 92.7% at 3 TeV $\rightarrow$ 99.6% at 10 TeV — a faithful reproduction.

$\Delta F = 2$ amplitudes — Blanke et al. 0809.1073, Fig. 2 (paper left, ours right)




Figure 9: Re/Im of the KK contribution to M_{12} for the K and B_s systems: paper (left) vs. ours (right).

Re/Im of the KK contribution to M_{12} for the K and B_s systems. For kaons, Blanke note $\text{Im}(M_{12}) \gg \text{Re}$ by $\sim 100\times$ (the ε_K problem); **ours** = $106\times$ (ratio of medians at 3 TeV). For B_s the contribution is $\mathcal{O}(1)$. Both reproduce the paper.

Existence vs. typical floors (our analysis, ours only)




Figure 10: Per constraint: **minimum** ratio-to-bound (the best fine-tuned point $\rightarrow$ existence floor) and **median** ratio (typical anarchic point) vs. M_{KK} .

For each constraint we plot the **minimum** ratio-to-bound (the best, fine-tuned point $\rightarrow$ the *existence* floor) and the **median** ratio (the *typical* anarchic point) vs. M_{KK} . The flavor constraints (ε_K , Δm_d , Δm_s , D^0 , Δm_K) are fully **tunable**: even at 1 TeV the best point beats the bound by $\sim 1000\times$ (alignment sends $\varepsilon_K \propto \text{Im } M_{12} \rightarrow 0$), so their existence floor is $\lesssim 1$ TeV. S, T, U is **irreducible** — min $\approx$ median (spread only $\sim 1.5\times$) because it has no Yukawa dependence — crossing the bound only at $\sim 18\text{--}20$ TeV. Hence even with maximally fine-tuned Yukawas, minimal RS cannot exist below $\sim 18\text{--}20$ TeV, set by oblique S, T, U ; $Z \rightarrow b\bar{b}$ is the next irreducible one at ~ 4.6 TeV.

4 Conclusion

The fixed, dual-reviewed code reproduces the published RS flavor/EWPT literature across seven figures from four papers, matching their quoted numbers (ε_K 10 TeV floor, D^0 containment, kaon $100\times \varepsilon_K$ problem, $C_{B_d}/C_{B_s} \approx 1$). The corrected minimal-model picture: the *typical* anarchic floor with current data is ~ 30 TeV, dominated by the (tunable) ε_K constraint; the *existence* floor for a fine-tuned point is $\sim 18\text{--}20$ TeV, dominated by the (irreducible) S, T, U oblique constraint. The earlier 25–30 TeV $Z \rightarrow b\bar{b}$ “floor” was a sign-bug artifact (now 5 TeV). Pushing below ~ 20 TeV requires custodial protection of T (custodial RS), which is the next branch of this program.