

Series 2a

1. Monte Carlo Approximation for Numerical Integration

Let $A, B \subseteq \mathbb{R}^2$ be the sets defined by

$$A = \{(x, y) \in \mathbb{R}^2 : (x - 2)^2 + y^2 \leq 4\} \quad \text{and} \quad B = \{(x, y) \in \mathbb{R}^2 : x^2 + (y - 2)^2 \leq 4\},$$

and consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) = \mathbb{1}_{A \cap B}(x, y) \cdot |x|^{2/3}.$$

Let (Ω, \mathcal{F}, P) be a probability space, and let $Y_n, Z_n : \Omega \rightarrow \mathbb{R}, n \in \mathbb{N}$, be independent $\mathcal{U}_{(0,1)}$ -distributed random variables. For each $N \in \mathbb{N}$, define the random variable $I_N : \Omega \rightarrow \mathbb{R}$ by

$$I_N = \frac{4}{N} \sum_{n=1}^N f(2Y_n, 2Z_n).$$

The random variables $I_N, N \in \mathbb{N}$, are thus Monte Carlo approximations of $\mathbb{E}_P[4f(2Y_1, 2Z_1)]$.

a) Prove or disprove the following statement: The random variables $I_N, N \in \mathbb{N}$, are P -unbiased estimators of the integral

$$\int_0^2 \int_0^2 f(x, y) \, dx \, dy.$$

b) Write a MATLAB function `MonteCarlo(N)` with input $N \in \mathbb{N}$ and output a realization of I_N .

c) Write a MATLAB function `MonteCarloPlot()` which, for each $k \in \{2, 3, 4, 5, 6\}$, plots five realizations of I_{10^k} , each represented by a blue star in a coordinate plane. Plot the values of k on the x -axis and the corresponding realizations of I_{10^k} on the y -axis. The plot should contain a total of 25 blue stars.

2. Approximative Realizations of a One-Dimensional Standard Brownian Motion

In this exercise, we do not distinguish between pseudo-random numbers and actual random numbers. Let A be the set defined by

$$A = \bigcup_{n=1}^{\infty} \{\mathbf{t} = (t_1, \dots, t_n) \in [0, \infty)^n : \#\mathbb{R}(\{t_1, \dots, t_n\}) = n\},$$

and define the function $\text{length} : A \rightarrow \mathbb{N}$ such that for all $n \in \mathbb{N}$ and all $\mathbf{t} = (t_1, \dots, t_n) \in [0, \infty)^n \cap A$,

$$\text{length}(\mathbf{t}) = n.$$

Let further $Q : A \rightarrow \bigcup_{n=1}^{\infty} \mathbb{R}^{n \times n}$ be the function defined such that for all $n \in \mathbb{N}$ and for all $\mathbf{t} = (t_1, \dots, t_n) \in [0, \infty)^n \cap A$, we have

$$Q(\mathbf{t}) = (\min\{t_i, t_j\})_{(i,j) \in \{1, \dots, n\}^2}.$$

Write a MATLAB function `StandardBrownianMotion(t)` with input $\mathbf{t} \in A$ and output a realization of an $\mathcal{N}_{0, Q(\mathbf{t})}$ -distributed random variable. The MATLAB function `StandardBrownianMotion(t)` may use at most $\text{length}(\mathbf{t})$ realizations of an $\mathcal{N}_{0, I_{\mathbb{R}}}$ -distributed random variable.

Test your implementation by calling the following MATLAB commands:

```

1 % Set random number generator to default
2 rng('default');
3
4 % Define parameters
5 N = 10^3;
6 preimage = linspace(0, 1, N+1);
7
8 % Generate and plot realizations of the standard Brownian motion
9 X = StandardBrownianMotion(preimage);
10 plot(preimage, X, 'b');
11 hold on;
12
13 X = StandardBrownianMotion(preimage);
14 plot(preimage, X, 'r');
15
16 X = StandardBrownianMotion(preimage);
17 plot(preimage, X, 'g');
18
19 % Add labels and a title
20 xlabel('t');
21 ylabel('X(t)');
22 title('Realizations of Standard Brownian Motion');
23 legend('BM (blue)', 'BM (red)', 'BM (green)');
24
25 hold off;

```

3. Brownian Motion and Monte Carlo Simulation

Let (Ω, \mathcal{F}, P) be a probability space and $W: [0, T] \times \Omega \rightarrow \mathbb{R}$ be a standard Brownian motion ($T > 0$).

a) For $N \in \mathbb{N}$, define a temporal discretization (or mesh) on the time interval $[0, T]$ by:

$$0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T, \quad t_n := \frac{n}{N}T. \quad (1)$$

Write a MATLAB function **BrownianMotion**(T, N) with inputs $T \in (0, \infty)$ and $N \in \mathbb{N}$, and output a realization of a $(W_0, W_{t_1}, \dots, W_{t_{N-1}}, W_T)$ $(P_{\mathcal{B}(\mathbb{R}^{N+1})})$ -distributed random variable. (You may use your results from Problem 2.) Now, assume $T = 1$. Write a MATLAB function **BrownianMotionPlot**() which uses the function **BrownianMotion**(T, N) to generate and plot 5 sample paths of the process \widetilde{W}^{1000} over the time interval $[0, 1]$, where, for $N \in \mathbb{N}$, the process $\widetilde{W}^N: [0, T] \times \Omega \rightarrow \mathbb{R}$ is defined by the linear interpolation of W at the mesh points $\{t_0, t_1, \dots, t_N\}$, that is:

$$\widetilde{W}_t^N = \left(n + 1 - \frac{tN}{T}\right) W_{t_n} + \left(\frac{tN}{T} - n\right) W_{t_{n+1}},$$

for all $n \in \{0, 1, \dots, N-1\}$ and each $t \in [t_n, t_{n+1}]$.

b) Let $X = (X_t)_{t \in [0, 1]}$ be the stochastic process defined by $X_t = 1 + t + 3W_t$. First, prove that

$$\mathbb{E}[e^{\sigma W_t}] = \exp\left(\frac{1}{2}\sigma^2 t\right), \quad \text{for all } \sigma \in \mathbb{R} \text{ and } t \in [0, 1].$$

Then, perform the following tasks:

- (i) Plot 10 sample paths of X , using the discretization (1) for $N = 10^3$.
- (ii) Compute $\mu := \mathbb{E}[\exp(X_1)]$ exactly and then approximate it using a Monte Carlo simulation with $M = 10^5$ samples. Provide a 95% confidence interval for your estimate based on the Central Limit Theorem (CLT), and compare the exact value of μ with your simulation result.

Submission Deadline: Wednesday, 23 October 2024, by 2:00 PM.