

STA 4102/5106: Homework Assignment #7

(Wednesday, October 29)

Due: Wednesday, November 5

1. Write a matlab program to implement a uniform random number generator using a multiplicative congruential method with $m = 2^{13} - 1$ and $a = 17$. Generate 500 numbers for the starting point $x_0 = 100$. You can use the **mod** command in matlab for evaluating the modulus function. For the sequence $u_i = x_i/m$:
 - (a) Plot a histogram to display the results.
 - (b) Calculate the coefficient correlation of the pairs of successive number u_i and u_{i+1} . (You can use **corrcoef** in matlab).
 - (c) Plot the pairs (u_i, u_{i+1}) on a 2D plot.
2. Repeat Problem 1 with $a = 85$.
3. Derive and implement a method to generate samples of a Weibull random variable whose probability distribution function is given by:

$$F(x) = 1 - \exp(-\alpha x^\beta), \quad 0 < x < \infty$$

Run your program to simulate 1000 values of Weibull random variable with $\alpha = 1$ and $\beta = 0.5$. Draw the histogram of the simulated values with 100 bins.

4. **(STA 5106 Students Only)** Let $X_i, i = 1, 2, \dots, n$ be a sequence of independent and identically distributed exponential random variables with mean 1. Define a random variable Z_i , for $i > 1$

$$Z_i = \begin{cases} 1 & \text{if } X_i \geq \max(X_1, X_2, \dots, X_{i-1}) \\ 0 & \text{otherwise} \end{cases}$$

We will assume that $Z_1 = 1$. In the sequence $Z_i, i = 1, 2, \dots, n$, we are interested in how many times $Z_i = 1$ are separated by k time points, where $k = 0, 1, \dots, 8$.

- (a) Write a matlab program to simulate the sequence Z_i for $n = 200$.
- (b) Compute the number of times $Z_i = 1$ are separated by k time points. Call this random value Y_k .
- (c) Repeat the above simulation $m = 3000$ times. Draw the histogram the values of Y_k and compute the mean of those simulated values. Compare the histogram of Y_k with the probability mass function of a Poisson random variable with mean $1/(k+1)$.