

PROJECT III

SIMULATION-BASED INFERENCE USING BOOTSTRAP AND MONTE CARLO TECHNIQUE

III

SIMULATION-BASED INFERENCE USING BOOTSTRAP AND MONTE CARLO TECHNIQUE

Bootstrap · Monte Carlo · Statistical Simulation

Submitted: March 23, 2022

Oscar Bismark Tetteh

Advanced Econometrics · 2022

Description of Data

This study used the same data sets as Project II. The deflated total commodity price and the nominal Canada-US exchange rate will be used in this study. Because there was no recent data for the Canada-US nominal exchange rate, the study used data spanning from January 1972 to April 2017.

The objective of the study

This project aims to use simulation-based inference to revise the inferences made in project II using conventional tests. The main reason for using the simulation-based inference approach is to see if it improves on the conventional inference I performed in project II. Based on simulation-based inference, the conventional test will be used to test for cointegration, residual normality (Jarque - Bera Test), Granger - causality, residual autocorrelation, and skewness.

The theory of Simulation-Based Inference

Statistical inference consists of the use of statistics to draw conclusions about some unknown aspect of a population based on a random sample from that population. However, when making such inferences, the test statistic plays a significant role. This is because the test statistic contains information about the data that is relevant for deciding whether to reject the null hypothesis. Sometimes, we are unable to conduct such inferences due to the nature of the distribution of the test statistics (especially when the test statistic has no analytical distribution). It has been demonstrated that simulation-based techniques are more appropriate or provide better estimates for test statistics in such cases.

In general, simulation-based techniques are used when no analytical distribution for a test statistic is available, as well as when asymptotic distributions are available, but simulation is expected to improve finite-sample properties. Simulation-based techniques are used to estimate the distribution of an estimator or test statistic when the analytical distribution is unknown. Monte Carlo simulation techniques, bootstrap techniques, and resampling techniques are a few examples of simulation techniques.

The central idea behind simulation-based techniques is to create a large number of simulated samples and then use them to calculate the test statistic or estimator of interest. The calculated test statistics are used to build the empirical distribution, which

is then used to answer the question of how likely the actual test statistics are to behave under the null hypothesis.

The Kendal Density Graph in Simulation-Based – Inference

A kernel density curve may follow the shape of the distribution more closely. In short, the technique allows one to create a smooth curve given a set of random data. However, the estimation can also be used to generate points that only appear to have come from a specific sample set. This feature is particularly useful in simulation-based inference. One of the principal advantages of using kernel density graphs to represent the test statistic from a simulation is that they are better at determining the distribution shape because the number of bins used has no effect on them.

I used the kernel density graph in this project to compare the simulated test statistics to the normal density. This gives an idea of how well the simulated test statistic performs in comparison to the test statistic computed using the conventional approach to inference.

Normality Tests (Jarque-Bera test) using Simulation-Based Inference

The assumption of normality is that the underlying residuals are normally distributed, or approximately so. The normality of the residuals is an assumption made when running a linear model. So, if your residuals are normal, your assumption is correct, and your model inference (confidence intervals, model predictions) should be correct as well. If your residuals are not normal, there could be a problem with the model's fit, stability, or reliability. The Jarque - Beta test was used in the previous project to determine the residuals' normality. The test results indicated that the residuals are not normally distributed at a 5% significance level, indicating possible misspecification of the VAR model.

The conventional test will be employed here to test for normality in residuals. The test's null hypothesis states that residuals are normally distributed, while the alternate hypothesis states that residuals are not normally distributed.

The conventional Jarque-Bera test for normality shows that the null hypothesis is rejected at a 5% level of significance (see table 1.1). As a result of this result, the disturbance term is not normally distributed. Since the null hypothesis has been rejected, it is expected that bootstrap and Monte Carlo simulation-based techniques will improve the test statistic's ability to reject the null hypothesis when it is actually false.

The Jarque – Bera test for normality test is pivotal.

$$JB = \frac{n}{6} \left(S^2 + \frac{1}{4} (K - 3)^2 \right)$$

It does not depend on any parameters of the model. The analytical distribution in a finite sample is unknown, but the asymptotic distribution is known which is Chi-square. In this situation, the simulation-based techniques are more appropriate or provide better estimates for test statistics. The simulation results confirm that the null of normality should be rejected (Table 1.2). The conventional and simulation tests revealed that the residuals are not normally distributed, implying that the vector autoregressive model was misspecified. This means that the model does not account for all of the trends in the dataset. Both the normal distribution line and the simulated normal distribution are depicted on the kernel density graph (Table 1.3).

Conventional Approach

Selection of lag order based on AIC (Table 1.0)

. varsoc d.logDTCP d.logEX, exo(mon)								
Selection-order criteria								
Sample: 1972m6 - 2017m4					Number of obs		=	539
lag	LL	LR	df	p	FPE	AIC	HQIC	SBIC
0	2581.11				2.4e-07	-9.56254	-9.55009	-9.53071
1	2619.38	76.555*	4	0.000	2.1e-07*	-9.68973*	-9.66483*	-9.62606*
2	2623.06	7.3529	4	0.118	2.1e-07	-9.68853	-9.65118	-9.59303
3	2624.68	3.2504	4	0.517	2.1e-07	-9.67972	-9.62991	-9.55238
4	2629.08	8.7959	4	0.066	2.1e-07	-9.6812	-9.61894	-9.52202
Endogenous: D.logDTCP D.logEX								
Exogenous: mon _cons								

Vector autoregression result using lag order of 1

```
. var d.logDTCP d.logEX, lags(1/1)
```

Vector autoregression

Sample: 1972m3 - 2017m4
 Log likelihood = 2636.389
 FPE = 2.09e-07
 Det(Sigma_ml) = 2.04e-07

Number of obs = 542
 AIC = -9.706231
 HQIC = -9.687638
 SBIC = -9.658682

Equation	Parms	RMSE	R-sq	chi2	P>chi2
D_logDTCP	3	.035836	0.0815	48.09719	0.0000
D_logEX	3	.013904	0.0889	52.91689	0.0000

	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
D_logDTCP					
logDTCP	.2552216	.0460356	5.54	0.000	.1649935 .3454497
LD.					
logEX	-.1489792	.1180214	-1.26	0.207	-.3802969 .0823385
LD.					
_cons	-.0004033	.0015361	-0.26	0.793	-.0034139 .0026073
D_logEX					
logDTCP	-.0380915	.017861	-2.13	0.033	-.0730984 -.0030845
LD.					
logEX	.2416676	.0457903	5.28	0.000	.1519203 .331415
LD.					
_cons	.0003813	.000596	0.64	0.522	-.0007868 .0015493

Results from the conventional test (Table 1.1)

```
. varnorm, jbera
```

Jarque-Bera test

Equation	chi2	df	Prob > chi2
D_logDTCP	284.507	2	0.00000
D_logEX	166.073	2	0.00000
ALL	450.580	4	0.00000


Simulation Results (Table 1.2)

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
jqb	1,500	1395.695	75.38054	1033.403	1725.32
rej	1,500	1	0	1	1

```
. histogram jqb,normal kdensity  
(bin=31, start=1033.403, width=22.319903)
```

The kernel density graph for the simulation results (Table 1.3)



Lagrange Multiplier test for residual autocorrelation using Simulation-Based Inference

The LM test for residual autocorrelation determines whether autocorrelation exists in the residuals for a given lag order. In general, the residuals from a VAR model should be white noise with no autocorrelation. If there is autocorrelation among the residuals, it indicates that some information was not accounted for by the model, such as insufficient lags. LM test for residual autocorrelation demonstrates that the VAR in the levels model is correctly specified. In the previous project (project II), I investigated whether there is autocorrelation in the residuals for a given lag order. The results showed that no autocorrelation exists in the residuals for any lag order. This shows that the VAR in the levels model was properly specified. For the purposes of this study, I will employ the conventional method of testing for autocorrelation in residuals for any lag orders estimated after the vector autoregressive model. The null hypothesis states that there is no autocorrelation in the residuals for any lag order, whereas the alternative hypothesis states that there is autocorrelation in the residuals for any lag orders estimated after the vector autoregressive model. The LM test for autoregression test is pivotal. It does not depend on any parameters of the model. The analytical distribution in a finite sample is unknown, but the asymptotic distribution is known which is Chi-square. The conventional test revealed no autocorrelation in the residuals for any lag order. Because the significance probability is greater than the significance level. (See Table 2.1). However, since the LM test statistic is asymptotically pivotal and does not depend on any parameter, we anticipate that parametric bootstrap techniques will improve the test for autocorrelation in residuals. After the simulation test, the outcome of the test revealed that there is no autocorrelation in the residuals for any lag order. Because the significance probability is greater than the significance level. (See Table 2.2). In this case, the simulation-based technique produced results, which is consistent which the conventional approach.

Conventional Approach (Table 2.0)

```
. var d.logDTCP d.logEX, lags(1/1) exog(mon)
```

Vector autoregression

Sample: 1972m3 - 2017m4 Number of obs = 542
 Log likelihood = 2636.677 AIC = -9.699917
 FPE = 2.10e-07 HQIC = -9.675126
 Det(Sigma_ml) = 2.04e-07 SBIC = -9.636518

Equation	Parms	RMSE	R-sq	chi2	P>chi2
D_logDTCP	4	.035863	0.0818	48.2969	0.0000
D_logEX	4	.013915	0.0892	53.08976	0.0000

	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
D_logDTCP						
logDTCP						
LD.	.2544378	.0460642	5.52	0.000	.1641537	.3447219
logEX						
LD.	-.1507089	.1180705	-1.28	0.202	-.3821229	.080705
mon	-4.21e-06	9.82e-06	-0.43	0.668	-.0000234	.000015
_cons	.0013485	.004369	0.31	0.758	-.0072146	.0099117
D_logEX						
logDTCP						
LD.	-.0383732	.0178725	-2.15	0.032	-.0734028	-.0033437
logEX						
LD.	.2410458	.0458105	5.26	0.000	.1512589	.3308326
mon	-1.51e-06	3.81e-06	-0.40	0.691	-8.98e-06	5.95e-06
_cons	.0010111	.0016952	0.60	0.551	-.0023114	.0043335

LM test results (Table 2.1)

```
. varlmar, mlag (3)
```

Lagrange-multiplier test

lag	chi2	df	Prob > chi2
1	7.1779	4	0.12678
2	8.5441	4	0.07356
3	4.0325	4	0.40163

H0: no autocorrelation at lag order

Simulation results (Table 2.2)

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
LM	500	67.29871	14.2565	33.08839	107.2492
rej	500	1	0	1	1

```
. histogram LM,kdensity  
(bin=22, start=33.088394, width=3.3709474)
```

Cointegration test using simulation-based inference

The Augmented Dickey-Fuller test for a unit root in the residual was used to perform the conventional test for cointegration. The Augmented Dickey-Fuller test was used because there is the possibility of autocorrelation in the residuals. The null hypothesis of the test states that there is a unit root in the residuals, whereas the alternative hypothesis states that there is no unit root in the residuals. I proceed with implementing the Augmented Dickey-Fuller (ADF) Test. The test statistic in ADF showed a value of -3.643 (Table 3.2) and the asymptotic critical value of 2.89 at a 5% level of significance. In reference to Table 3.2, the output shows that the test statistic value -3.643 of the test statistic is smaller than the 5% asymptotic critical value of 2.89 . Therefore, at a 5% level of significance, there is enough evidence to fail to reject the null hypothesis, which means we can confidently conclude that the Canada-US Nominal Spot Exchange rate and deflated commodity prices has a unit root in the residuals and hence, not cointegrated.

The simulation-based inference techniques such as bootstrap and Monte Carlo simulation were used to improve test inference. The main assumption of these techniques is that there is no analytical distribution available or that it cannot be derived. Bootstrap and Monte Carlo are expected to improve the test statistic's finite sample properties can be enhanced its power. The bootstrap and Monte Carlo simulation-based techniques are expected to improve the test statistic's ability to reject the null hypothesis when it is actually false.

Also, if the test is asymptotically pivotal, then the parametric bootstrap will be best, and it can improve inference in finite samples. Residuals will be estimated based on simulated data.

After simulation, the null hypothesis that there is a unit root in the residual was rejected, this is because it was revealed that the test statistic for the simulated distribution is greater than the test statistic for the asymptotic distribution and the significant probability from the simulated distribution is statistically. The results of the simulation-based inference show that commodity prices and the Canada-US exchange rate are cointegrated. Because the null is false, we can conclude that the simulated-based inference improved the test statistic's ability to reject the null of a unit root in the residuals. Kernel density estimation is a non-parametric method for estimating the probability density function of a random variable from a finite data sample. The kernel density, however, illustrates that the distribution is normally distributed (Table 3.4).

Conventional Approach (Table 3.0)

. reg logDTCP logEX mon						
Source	SS	df	MS	Number of obs	=	544
Model	26.7820389	2	13.3910194	F(2, 541)	=	563.84
Residual	12.8485174	541	.02374957	Prob > F	=	0.0000
				R-squared	=	0.6758
				Adj R-squared	=	0.6746
Total	39.6305562	543	.07298445	Root MSE	=	.15411
logDTCP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
logEX	-1.383363	.0493556	-28.03	0.000	-1.480315	-1.286411
mon	-.0005653	.0000428	-13.22	0.000	-.0006493	-.0004814
_cons	1.338811	.0197106	67.92	0.000	1.300092	1.377529


Simulation results (Table 3.3)

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
df_test	987	-13.57424	.7176953	-15.78872	-11.45785
rej	987	0	0	0	0

```
. histogram df_test,kdensity  
(bin=29, start=-15.788724, width=.14934063)
```

The kernel density graph for the simulation results (Table 3.4)



Granger Causality Test using simulation-based Inference

Granger causality is a causality concept derived from the idea that causes may not always occur aftereffects and that if one variable is the cause of another, knowing the status of the cause at an earlier point in time can help predict the effect at a later point in time (Granger, 1969; Lütkepohl 2005, p.41). The granger causality test was used in the previous project (Project II) to determine whether the Canada-US Spot Exchange Rate does not cause or has no explaining power in predicting Deflated Commodity Prices. At the 5% significance level, we fail to reject the null hypothesis, implying that the exchange rate does not granger-cause commodity prices. However, a granger causality test was performed on the normal variables again, and the results were compared to the variables in the first difference, revealing that the exchange rate granger causes commodity price. Bootstrap and Monte Carlo simulation techniques will be applied to improve inference. I then used the conventional approach to test whether Canada-US exchange rates granger-cause commodity prices, and then added one extra lag number to overfit the model for generalization and inference. The null hypothesis of the test is that the Canada-US exchange rate does not granger cause commodity prices, whereas the alternative indicates that exchange rates granger cause commodity prices. The conventional test results for the granger causality test (Table 4.1) show that the F-statistic value (3.18) is greater than the F-critical value (2.9957) at a 5% significance level, and the significant probability of 0.0422 is also statistically significant at a 5% significance level, providing enough evidence to reject the null and conclude that Canada-US exchange rates do granger-cause commodity prices. Because the null is false, we can conclude that the simulated-based inference improved the test statistic's ability to reject the null hypothesis that Canada-US exchange rates do not granger cause commodity prices. The test statistics (F - Statistic) is pivotal. This implies that the test statistic does not depend on the nuisance parameter. This is due to the fact that the F-statistic used in this analysis does not depend on other parameters and the probability of type I error will be exactly significant level.

After several simulations, the simulated test statistic and its p-value (0.037333) (Table 4.2) confirm the conventional test result. At a 5% significance level, the null hypothesis is rejected. Both tests indicate that the granger exchange rate causes deflated commodity prices. The kernel density, on the other hand, shows that the distribution is positively skewed (Table 4.3).

Conventional Approach

Selection of lag order (Table 4.0)

```
. varsoc logDTCP logEX, exo(mon)
```

Selection-order criteria
Sample: 1972m5 - 2017m4

Number of obs = 540

lag	LL	LR	df	p	FPE	AIC	HQIC	SBIC
0	575.247				.000413	-2.11573	-2.1033	-2.08394
1	2593.14	4035.8	4	0.000	2.4e-07	-9.57459	-9.54973	-9.51101
2	2633.87	81.459*	4	0.000	2.1e-07*	-9.71063*	-9.67333*	-9.61526*
3	2637.5	7.2585	4	0.123	2.1e-07	-9.70925	-9.65952	-9.5821
4	2638.89	2.7891	4	0.594	2.1e-07	-9.6996	-9.63744	-9.54066

Endogenous: logDTCP logEX
Exogenous: mon _cons

```
. regress logDTCP mon L(1/2).logDTCP L(1/2).logEX
```

Source	SS	df	MS	Number of obs	=	542
Model	38.9479113	5	7.78958226	F(5, 536)	=	6155.19
Residual	.67832419	536	.00126553	Prob > F	=	0.0000
Total	39.6262355	541	.073246276	R-squared	=	0.9829
				Adj R-squared	=	0.9827
				Root MSE	=	.03557

logDTCP	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
mon	-.0000235	.0000114	-2.06	0.040	-.000046 -1.05e-06
logDTCP					
L1.	1.235257	.0462774	26.69	0.000	1.14435 1.326165
L2.	-.267745	.0460439	-5.81	0.000	-.3581936 -.1772964
logEX					
L1.	-.1722598	.1180758	-1.46	0.145	-.4042078 .0596882
L2.	.1329381	.1177631	1.13	0.259	-.0983956 .3642719
_cons	.0442082	.0142007	3.11	0.002	.0163123 .072104

The conventional test statistic (Table 4.1)

```
. test(L.logEX) (L2.logEX)

( 1)  L.logEX = 0
( 2)  L2.logEX = 0

          F( 2, 536) = 3.18
          Prob > F = 0.0422
```


Simulation from the test statistic (Table 4.2)

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
F_test	1,500	.9865299	.9935846	7.65e-06	7.716124
rej	1,500	.0373333	.1896405	0	1

```
. histogram F_test, kdensity
(bin=31, start=7.649e-06, width=.24890697)
```

The kernel density graph for the simulation results (Table 4.3).



Simulation-based inference for granger causality in first difference (Table 4.4)

```
. var d.logDTCP d.logEX, lags(1/1)
```

Vector autoregression

Sample: 1972m3 - 2017m4
 Log likelihood = 2636.389
 FPE = 2.09e-07
 Det(Sigma_ml) = 2.04e-07

Number of obs = 542
 AIC = -9.706231
 HQIC = -9.687638
 SBIC = -9.658682

Equation	Parms	RMSE	R-sq	chi2	P>chi2
D_logDTCP	3	.035836	0.0815	48.09719	0.0000
D_logEX	3	.013904	0.0889	52.91689	0.0000

	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
D_logDTCP						
logDTCP						
LD.	.2552216	.0460356	5.54	0.000	.1649935	.3454497
logEX						
LD.	-.1489792	.1180214	-1.26	0.207	-.3802969	.0823385
_cons	-.0004033	.0015361	-0.26	0.793	-.0034139	.0026073
D_logEX						
logDTCP						
LD.	-.0380915	.017861	-2.13	0.033	-.0730984	-.0030845
logEX						
LD.	.2416676	.0457903	5.28	0.000	.1519203	.331415
_cons	.0003813	.000596	0.64	0.522	-.0007868	.0015493

```
. vargranger
```

Granger causality Wald tests

Equation	Excluded	chi2	df	Prob > chi2
D_logDTCP	D.logEX	1.5934	1	0.207
D_logDTCP	ALL	1.5934	1	0.207
D_logEX	D.logDTCP	4.5482	1	0.033
D_logEX	ALL	4.5482	1	0.033

Simulated results from the test statistic – the first difference (Table 4.5)

. summarize					
Variable	Obs	Mean	Std. Dev.	Min	Max
F_test	1,500	.9865299	.9935846	7.65e-06	7.716124
rej	1,500	.0373333	.1896405	0	1

Skewness and Kurtosis test using Simulation-Based Inference

Skewness allows you to determine how much a distribution's overall shape deviates from the shape of the normal distribution. Kurtosis is a measure of how different the tails of a distribution are from the tails of a normal distribution. Kurtosis focuses on the tail shape, whereas skewness focuses on the overall shape. If skewness is equal to zero, the data is perfectly symmetrical. Real-world data is rarely if ever perfectly symmetric, instead, we get data that is approximately symmetric. The null hypothesis of the test is that the residual term has a skewness of zero. The conventional test was performed, and the results indicate that the null should be rejected at a 5% level of significance (see Table 5.2). As a result, the residual term appears to be skewed. The simulated test statistic, on the other hand, failed to reject the null of zero skewness. This means that the simulated test result indicates that the normal distribution is skewed. I used the kernel density graph, which shows that the distribution is not skewed and deviates significantly from the standard normal distribution. As a result, the conclusions from the two tests differ depending on the power of each test's test statistic.

Conventional Approach

Vector autoregression (Table 5.1)

Vector autoregression

Sample: 1972m4 - 2017m4
Log likelihood = 2634.24
FPE = 2.10e-07
Det(Sigma_ml) = 2.02e-07

Number of obs = 541
AIC = -9.701441
HQIC = -9.670406
SBIC = -9.62208

Equation	Parms	RMSE	R-sq	chi2	P>chi2
D_logDTCP	5	.035847	0.0860	50.88424	0.0000
D_logEX	5	.013913	0.0925	55.11625	0.0000

	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
D_logDTCP						
logDTCP						
LD.	.2464391	.0471523	5.23	0.000	.1540222	.338856
L2D.	.0221817	.0472906	0.47	0.639	-.0705061	.1148696
logEX						
LD.	-.1049544	.1212342	-0.87	0.387	-.3425692	.1326603
L2D.	-.1486392	.1210338	-1.23	0.219	-.3858612	.0885828
_cons	-.0003591	.0015355	-0.23	0.815	-.0033687	.0026505
D_logEX						
logDTCP						
LD.	-.0362533	.0183006	-1.98	0.048	-.0721219	-.0003847
L2D.	-.01433	.0183543	-0.78	0.435	-.0503038	.0216438
logEX						
LD.	.2553083	.0470531	5.43	0.000	.1630859	.3475307
L2D.	-.0673661	.0469753	-1.43	0.152	-.159436	.0247039
_cons	.0004103	.000596	0.69	0.491	-.0007578	.0015784

Results from Conventional test (Table 5.2)

. varnorm, skewness					
Skewness test					
Equation	Skewness	chi2	df	Prob > chi2	
D_logDTCP	-.5033	22.841	1	0.00000	
D_logEX	.19629	3.474	1	0.06234	
ALL		26.315	2	0.00000	


Simulation results for Skewness (Table 5.3)

```
. summarize
```

Variable	Obs	Mean	Std. Dev.	Min	Max
sk	1,500	56.19085	4.300095	34.69073	70.78248
rej	1,500	1	0	1	1

```
. histogram sk,normal kdensity
(bin=31, start=34.690735, width=1.1642498)
```

The kernel density graph for the simulation results (Table 5.4)



STATA CODES

Stata Codes for Normality test using simulation

```
//%%% Normality test using simulation-based inference%%%%%%%%//
```

```
clear all
```

```
import excel "C:\Users\bisma\OneDrive\Desktop\Winter\Econometrics\Module  
3\Data_Project 2.xlsx", sheet("Data for Project 1 - Copy") cellrange(A1:I545) firstrow  
gen logEX = ln( CanadaUSSpotExchangerate)  
gen logDTCP =ln( TotalAllCommodities/ US_CPI_Index)
```

```
generate mon = tm(1972m1)+_n-1
```

```
format mon %tm
```

```
tsset mon, monthly
```

```
//#####conventional testing#####//
```

```
varsoc d.logDTCP d.logEX, exo(mon)
```

```
var d.logDTCP d.logEX, lags(1/1)
```

```
varnorm, jbera
```

```
//#####bootstrapping for normality #####//
```

```
predict rhs, xb
```

```
set seed 42457786
```

```

gen y=.
tempname jarcque
postfile `jarcque' jqb rej using resultsjqb, replace
quietly {
  forvalues i=1/1500 {
    replace y=rhs+rnormal(0,1)
    var d.y d.logEX, lags(1/1)
    varnorm, jbera
    matrix v=r(jb)
    scalar jqb=v[3,1]
    scalar rej=(jqb> 450.580 )
    post `jarcque' (jqb) (rej)
  }
}
postclose `jarcque'

use resultsjqb, clear
summarize
histogram jqb,normal kdensity

```



```

replace y=rnormal(0,1)
var d.y d.logEX, lags(1/1)
varlmar, mlag (3)
matrix v=r(lm)
scalar LM= v[1,2]
scalar rej=(LM> 7.1779)
post `Lagrangm' (LM) (rej)
}
}
postclose `Lagrangm'

```

```
use results, clear
```

```
summarize
```

```
histogram LM,kdensity
```

Stata Code for Cointegration Test

```

////////////////Cointegration      test      using      Simulation
test////////////////

```

```
clear all
```

```

import excel "C:\Users\bisma\OneDrive\Desktop\Winter\Econometrics\Module
3\Data_Project 2.xlsx", sheet ("Data for Project 1 - Copy") cellrange(A1:I545) firstrow

```

```
gen logEX = ln( CanadaUSSpotExchangerate)
```

```
gen logDTCP =ln( TotalAllCommodities/ US_CPI_Index)
```

```
generate mon = tm(1972m1) +_n-1
```

```
format mon %tm
```

```
tsset mon, monthly
```

```
//////////#####conventional test#####//////////
```

```
reg logDTCP logEX mon
```

```
predict e, residuals
```

```
varsoc e //lag selection- order criteria is based on AIC//
```

```
dfuller e, regress lag (2)
```

```
//////////##### Bootstrapping - Simulation #####//////////
```

```
set seed 48889900
```

```
scalar constant=_b[_cons]
```

```
gen y=.
```

```
gen u=.
```

```
gen resd=.
```

```
tempname dfulco
```

```
postfile `dfulco' df_test rej using results, replace
```

```
quietly{
```

```
forvalues i=1/987 {
```

```
replace u=rnormal(0,1)
```

```
replace y=constant+u
```

```
reg y logEX mon
```

```
predict e1, residuals
```

```
replace resd=e1
```

```
drop e1
```

```

dfuller resd, regress lag(2)
scalar df_test=r(Zt)
scalar rej=(df_test> -3.643)
post `dfulco' (df_test) (rej)
}
}
postclose `dfulco'

```

```

use results, clear
summarize
histogram df_test,kdensity

```

Stata Codes for Granger Causality Test

```

/////////%%%%%%%%%Simulation Based Inference for Granger Causality
Test%%%%%%%%////////

```

```

clear all

```

```

import excel "C:\Users\bisma\OneDrive\Desktop\Winter\Econometrics\Module
3\Data_Project 2.xlsx", sheet("Data for Project 1 - Copy") cellrange(A1:I545) firstrow
gen logEX = ln( CanadaUSSpotExchangerate)
gen logDTCP =ln( TotalAllCommodities/ US_CPI_Index)

```

```

generate mon = tm(1972m1)+_n-1
format mon %tm

```

```
tsset mon, monthly
```

```
//#####conventional testing#####//
```

```
varsoc logDTCP logEX, exo(mon)
```

```
regress logDTCP mon L(1/2).logDTCP L(1/2).logEX
```

```
test(L.logEX) (L2.logEX)
```

```
//#####bootstrapping#####//
```

```
set seed 6743987
```

```
scalar constant=_b[_cons]
```

```
scalar b1=_b[L1.]
```

```
scalar b2=_b[L2.]
```

```
gen y=.
```

```
gen u=.
```

```
tempname Ftest
```

```
postfile `Ftest' F_test rej using results, replace
```

```
quietly {
```

```
forvalues i=1/1500 {
```

```
replace u=rnormal()
```

```
replace y=constant+b1+b2+u
```

```
regress y mon L(1/2).y L(1/2).logEX
```

```
test (L.logEX) (L2.logEX)
```

```
scalar F_test=r(F)
```

```
scalar rej=(F_test>3.18)
```

```
post `Ftest' (F_test) (rej)
```

```
}
```

```
}
```

```
postclose `Ftest'
```

```
use results, clear
```

```
summarize
```

```
histogram F_test, kdensity
```

```
//// %%%%%%%%%%Granger Causality - Simulation for first difference  
Variable %%%%%%%%%%///////
```

```
clear all
```

```
import excel "C:\Users\bisma\OneDrive\Desktop\Winter\Econometrics\Module  
3\Data_Project 2.xlsx", sheet("Data for Project 1 - Copy") cellrange(A1:I545) firstrow
```

```
gen logEX = ln( CanadaUSSpotExchangerate)
```

```
gen logDTCP =ln( TotalAllCommodities/ US_CPI_Index)
```

```
generate mon = tm(1972m1)+_n-1
```

```
format mon %tm
```

```
tsset mon, monthly
```

```
//conventional testing//
```

```
var d.logDTCP d.logEX, lags(1/1)
```

```
vargranger
```

```

//bootstrapping//
predict rhs, xb
set seed 678544

gen y=.

tempname granger
postfile `granger' grg rej using resultsgrg, replace
quietly {
  forvalues i=1/500 {
    replace y=rhs+rnormal(0,1)
    var d.y d.logEX, lags(1/1)
    vargranger
    matrix v=r(gstats)
    scalar grg= v[2,1]
    scalar rej=(grg> 1.5934)
    post `granger' (grg) (rej)
  }
}

postclose `granger'
use results, clear
summarize
histogram grg, kdensity

```

Skewness test using Simulations

```
//%%%%%%%%Skewness test using simulation %%%%%%%%%//
```

```
clear all
```

```
import excel "C:\Users\bisma\OneDrive\Desktop\Winter\Econometrics\Module  
3\Data_Project 2.xlsx", sheet("Data for Project 1 - Copy") cellrange(A1:I545) firstrow  
gen logEX = ln( CanadaUSSpotExchangerate)  
gen logDTCP =ln( TotalAllCommodities/ US_CPI_Index)
```

```
generate mon = tm(1972m1)+_n-1
```

```
format mon %tm
```

```
tsset mon, monthly
```

```
///##### conventional testing #####///
```

```
var d.logDTCP d.logEX, lags(1/2)
```

```
varnorm, skewness
```

```
/////#####bootstrapping#####/////
```

```
predict rhs, xb
```

```
set seed 866567435
```

```
gen y=.
```

```
tempname skew
```

```

postfile `skew' sk rej using resultssk, replace
quietly {
  forvalues i=1/1500 {
    replace y=rhs+rnormal(0,1)
    var d.y d.logEX, lags(1/2)
    varnorm, skewness
    matrix v=r(skewness)
    scalar sk=v[3,2]
    scalar rej=(sk>26.315)
    post `skew' (sk) (rej)
  }
}
postclose `skew'

use resultssk, clear
summarize
histogram sk,normal kdensity

```