

# A Tutorial for the package CAXA (conspecific attraction experiment analysis)

Sean E. MacDonald

December 13, 2017

## 1 Introduction

The package contains three functions that help clean and analyze data. The `Remove.WRSH.ZEDO` function removes experiments containing trials of the track species White-rumped shama (WRSH) and Zebra dove (ZEDO) due to low sampling effort. The `Keep.15.Tracks` function keeps experiments containing trials equal to 15 minutes in order to standardize methodology and low sampling efforts with track lengths not equal to 15 minutes. Lastly, the `GLMM.Bin` function applies a generalized linear mixed model with a binary distribution to the data in order to see what bird species significantly respond to track species.

The dataset used in this package is a series of 80 experiments with control and treatment periods. Control periods did not include a speaker. During treatment periods bird species' tracks were broadcasted in various orders and lengths (min). Whether or not the experiment was conducted during the bird's breeding season (0 = no, 1 = yes), on a native plant species (0 = no, 1 = yes), and the behavioral response of fruit-eating birds (0 = no response, 1 = attracted) are included. Japanese white-eye (jawe), Red-billed leiothrix (rble), Red-vented bulbul (rvbu), Red-whiskered bulbul (rwbu), and all non-fruit eating species (other). The data was collected on the Island of Oahu, Hawaii, USA from 06/2016 - 07/2017. Sampling took place across 26 native and exotic plant species. Bird breeding season ranges from approximately March - August.

## 2 Example

```
> ## Load the required packages and dataset
> library(CAXA)
> data("cax_data")
> ##
> View(cax_data)
> ## Remove all experiments with track species containing wrsh
> ## or zedo. The newly cleaned dataframe (cax_data_1) should
```

```

> ## contain 573 observations now.
> cax_data_1 <- Remove.WRSH.ZEDO(cax_data)

# A tibble: 573 x 12
# Groups:   experiment [73]
  experiment track.order track.spp track.type track.length breeding.season
    <fctr>      <fctr>      <chr>      <chr>      <int>      <int>
1         1         1         1 control      con         15         1
2         1         2         1 control      con         15         1
3         1         3         1 control      con         15         1
4         1         4         1 control      con         15         1
5         1         5         1 rvbu        treat        60         1
6         2         1         1 control      con         15         1
7         2         2         1 control      con         15         1
8         2         3         1 control      con         15         1
9         2         4         1 control      con         15         1
10        2         5         1 jawe        treat        15         1
# ... with 563 more rows, and 6 more variables: native.plant <int>, jawe <int>,
#   rble <int>, rvbu <int>, rwbu <int>, other <int>

> ## Utilizing the cleaned dataframe (cax_data_1) from
> ## Remove.WRSH.ZEDO, this function will now keep only experiments
> ## with all trials of track lengths equal to 15 minutes. The newly
> ## cleaned dataframe (cax_data_2) should contain 448 observations now.
> cax_data_2 <- Keep.15.Tracks(cax_data_1)

# A tibble: 448 x 12
# Groups:   experiment [56]
  experiment track.order track.spp track.type track.length breeding.season
    <fctr>      <fctr>      <chr>      <chr>      <int>      <int>
1         13         1         1 control      con         15         0
2         13         2         1 control      con         15         0
3         13         3         1 control      con         15         0
4         13         4         1 control      con         15         0
5         13         5         1 rvbu        treat        15         0
6         13         6 jawe.rble treat        15         0
7         13         7         1 rwbu        treat        15         0
8         13         8         1 rble        treat        15         0
9         15         1         1 control      con         15         0
10        15         2         1 control      con         15         0
# ... with 438 more rows, and 6 more variables: native.plant <int>, jawe <int>,
#   rble <int>, rvbu <int>, rwbu <int>, other <int>

> ## Utilizing the latest cleaned dataframe (cax_data_2), this
> ## function applys a generalized linear mixed model with a binomial
> ## distribution. Bird species is the response variable; track species,
> ## breeding season, and plant origin as fixed effects; order as the

```

```
> ## one random effect. The output shows that the track species
> ## call's jawe, jawe.rble, rble, and rvbu significantly attract
> ## Japanese white-eyes in comparison to the intercept (i.e. control).
> ## The breeding season and plant origin did not significantly
> ## influence strength of response.
> GLMM.Bin(cax_data_2)
```

Generalized linear mixed model fit by maximum likelihood (Laplace  
Approximation) [glmerMod]

Family: binomial ( logit )

Formula:

species ~ track.spp + breeding.season + native.plant + (1 | track.order)

Data: cax\_data\_2

AIC	BIC	logLik	deviance	df.resid
498	535	-240	480	439

Scaled residuals:

Min	1Q	Median	3Q	Max
-2.3400	-0.5263	-0.4832	0.4631	2.1722

Random effects:

Groups	Name	Variance	Std.Dev.
track.order	(Intercept)	0.01191	0.1091

Number of obs: 448, groups: track.order, 8

Fixed effects:

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	-1.2871	0.2805	-4.589	4.46e-06 ***
track.sppjawe	2.9869	0.4133	7.226	4.96e-13 ***
track.sppjawe.rble	3.0937	0.6680	4.632	3.63e-06 ***
track.spprble	1.2887	0.3783	3.407	0.000658 ***
track.spprvbu	1.1971	0.3240	3.695	0.000220 ***
track.spprwbu	0.2429	0.3705	0.656	0.511997
breeding.season	-0.1335	0.2302	-0.580	0.562028
native.plant	-0.0773	0.2610	-0.296	0.767146

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Correlation of Fixed Effects:

	(Intr)	trck.sppj	trck..	trck.spprb	trck.spprv	trck.spprw	brdng.
track.sppjw	-0.212						
trck.sppjw.	-0.275	0.106					
trck.spprb	-0.257	0.217	0.143				
trck.spprv	-0.334	0.261	0.147	0.271			
trck.spprw	-0.316	0.217	0.155	0.260	0.278		

breedng.ssn	-0.353	-0.063	0.137	-0.074	-0.016	0.012	
native.plnt	-0.655	-0.044	0.062	-0.027	-0.001	0.005	-0.084