

# FYS3500 Spring 2024 - Problem set 2

Topic: Binding energy, Properties of the nuclear force

## Concepts of the week

Explain these concepts: Semi-Empirical Mass Formula (SEMF), Liquid Drop Model (LDM), the Nuclear Force

### Problem 1: Binding energy

- a) The mass of  $^{27}\text{Al}$  and  $^{235}\text{U}$  has been measured to be 26.9815 u and 235.0439 u respectively. Use this to estimate the binding energy of the two nuclei from equation (1)
- b) Find the binding energies of  $^{27}\text{Al}$  and  $^{235}\text{U}$  using the semi empirical binding energy formula in equation (2). How well does it agree with the result in a) and what are possible reasons for the deviations?  
(Use  $a_v = 15.5$  MeV,  $a_s = 16.8$  MeV,  $a_c = 0.72$  MeV and  $a_{\text{sym}} = 23$  MeV).

### Problem 2: Semi-empirical mass formula

The relation between the mass and the binding energy is

$$M(A, Z) = Z(m_p + m_e) + (A - Z)m_n - \frac{1}{c^2}B(A, Z) \quad (1)$$

The semi-empirical binding energy formula is

$$B(A, Z) = a_v A - a_s A^{2/3} - a_c \frac{Z(Z-1)}{A^{1/3}} - a_{\text{sym}} \frac{(A-2Z)^2}{A} (+\delta) \quad (2)$$

where  $a_v$ ,  $a_s$ ,  $a_c$  and  $a_{\text{sym}}$  are constants.

- a) Explain the origin of the different terms in  $B(A, Z)$ . Why is the formula called semi-empirical?
- b) Explain which terms in  $B(A, Z)$  that dominates for light and heavy nuclei respectively. What does this mean for the observed  $\frac{N}{A}$  ratio among light and heavy nuclei?
- c) Use (1) and 2 to show that for a given  $A$ , the  $Z$  value that corresponds to the lowest-mass isobar is

$$Z_{\min} = \frac{m_n c^2 - (m_p + m_e) c^2 + a_c A^{-1/3} + 4a_{\text{sym}}}{2a_c A^{-1/3} + 8a_{\text{sym}} A^{-1}}$$

### Problem 3: Nuclear force

- a) We say the nuclear force is charge independent. Still, a proton and a neutron can form a bound state but two protons or two neutrons cannot. Why?
- b) List and briefly explain the most important contributions to the nuclear potential.
- c) We know from experiment that the parity  $P$  of the ground state of the deuteron is  $(+1)$ . What can we then learn about the ground state spin and orbital angular momentum configurations? (Hint: you also need to know about the spin-dependence of the nuclear force)

## Problem 4 Deuteron and the nuclear force

A simple quantum mechanical model of the deuteron is based on a spherically symmetric potential well given by the following equation:

$$\begin{aligned} V(r) &= -V_0 & \text{for } r < R \\ &= 0 & \text{for } r > R \end{aligned} \quad (3)$$

- a) \*Extra\* Use the continuity and normalization conditions to evaluate the coefficients  $A$  and  $C$  in equation (4.4) and (4.5) in the deuteron wave function, pg 82 in Krane. Show that

$$A = [2\pi(R + 1/k_2)]^{-\frac{1}{2}} \quad (4)$$

and

$$C = [2\pi(R + 1/k_2)]^{-\frac{1}{2}} \cdot \frac{\sin(k_1 R)}{e^{-k_2 R}} \quad (5)$$

- b) What is the origin of the magnetic moment of a nucleon? Why does the neutron have a non-zero magnetic moment?
- c) Search online to find the experimentally determined value for the magnetic moment of the deuteron. Using the values  $\mu_{d,L=0} = 0.8798 \mu_N$  and  $\mu_{d,L=2} = 0.3101 \mu_N$ , find the percentage admixture of the ( $L = 2$ ) state in the ground state of the deuteron. Remember, the wave function must be normalized!