

# Mandatory Assignment 1

Oskar Idland

## Repo

The code for this assignment can be found at:

<https://github.com/Oskar-Idland/MAT-MEK4270-mandatory1>

## Dirichlet Problem

The Dirichlet problem is given by:

$$u(x, y, t) = \sin(k_x x) \sin(k_y y) \cos(\omega t).$$

Using that

$$\frac{\partial^2 u}{\partial t^2} = c^2 \nabla^2 u$$

we get:

$$\begin{aligned} \frac{\partial^2 u}{\partial t^2} &= -\omega^2 \sin(k_x x) \sin(k_y y) \cos(\omega t) \\ \frac{\partial^2 u}{\partial t^2} &= c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) = -c^2 \left( -k_x^2 \sin(k_x x) \sin(k_y y) \cos(\omega t) - k_y^2 \sin(k_x x) \sin(k_y y) \cos(\omega t) \right) \\ \omega^2 &= c^2 (k_x^2 + k_y^2) \implies \omega = c \sqrt{k_x^2 + k_y^2}. \end{aligned}$$

## Exact Solutions

To find the relation between  $\omega$ ,  $k_x$  and  $k_y$  we use the exact solution:

$$u(x, y, t) = e^{i(k_x x + k_y y - \omega t)}$$

$$\frac{\partial^2 u}{\partial t^2} = c^2 \nabla^2 u$$

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

$$\frac{\partial^2 u}{\partial x^2} = -k_x^2 e^{i(k_x x + k_y y - \omega t)}$$

$$\frac{\partial^2 u}{\partial y^2} = -k_y^2 e^{i(k_x x + k_y y - \omega t)}.$$

This shows their relationship to be:

$$\omega^2 = c^2 (k_x^2 + k_y^2) \implies \omega = c \sqrt{k_x^2 + k_y^2}.$$

## Dispersion Coefficient $\omega$

If we assume that  $k_x = k_y = k$ , and  $m_x = m_y$ , we get the following discrete solution:

$$u_{ij}^n = e^{i(kh(i+j) - \tilde{\omega}n\Delta t)}.$$

Using equation 1.3 from the assignment, we get:

$$\frac{u_{ij}^{n+1} - 2u_{ij}^n + u_{ij}^{n-1}}{\Delta t^2} = c^2 \left( \frac{u_{i+1,j}^n - 2u_{ij}^n + u_{i-1,j}^n}{h^2} + \frac{u_{i,j+1}^n - 2u_{ij}^n + u_{i,j-1}^n}{h^2} \right).$$

Adding this to the left-hand side of the equation for  $u_{ij}^n$ , we get:

$$\begin{aligned} \frac{u_{ij}^{n+1} - 2u_{ij}^n + u_{ij}^{n-1}}{\Delta t^2} &= \frac{e^{ikh(i+j)}}{\Delta^2} (e^{i\tilde{\omega}(n+1)\Delta t} - 2e^{-i\tilde{\omega}} + e^{-i\tilde{\omega}(n-1)\Delta t}) \\ \frac{u_{ij}^{n+1} - 2u_{ij}^n + u_{ij}^{n-1}}{\Delta t^2} &= \frac{e^{ikh(i+j)}}{\Delta^2} (e^{i\tilde{\omega}\Delta t} - 2 + e^{-i\tilde{\omega}\Delta t}). \end{aligned}$$

Doing the same for the right-hand side of the equation for  $u_{ij}^n$ , we get:

$$\begin{aligned} \frac{u_{i+1,j}^n - 2u_{ij}^n + u_{i-1,j}^n}{h^2} &= \frac{e^{i\tilde{\omega}n\Delta t}}{h^2} (e^{ikh(i+1+j)} - 2e^{ikh(i+j)} + e^{ikh(i-1+j)}) \\ \frac{u_{i+1,j}^n - 2u_{ij}^n + u_{i-1,j}^n}{h^2} &= \frac{e^{i(kh(i+j) - \tilde{\omega}n\Delta t)}}{h^2} (e^{ikh} - 2 + e^{-ikh}). \end{aligned}$$

Combining these results, we get:

$$\frac{e^{i(kh(i+j) - \tilde{\omega}n\Delta t)}}{\Delta^2} (e^{i\tilde{\omega}\Delta t} - 2 + e^{-i\tilde{\omega}\Delta t}) = c^2 \frac{e^{-i(\tilde{\omega}n\Delta t - kh(i+j))}}{h^2} (e^{ikh} - 2 + e^{-ikh} + e^{ikh} - 2 + e^{-ikh}).$$

Simplifying this, we get:

$$\frac{e^{-i\tilde{\omega}\Delta t} - 2 + e^{i\tilde{\omega}\Delta t}}{\Delta t^2} = 2c^2 \frac{e^{ikh} - 2 + e^{-ikh}}{h^2}.$$

Using the CLF number  $C = c\Delta t/h = 1/\sqrt{2}$ :

$$\begin{aligned} e^{i\tilde{\omega}\Delta t} - 2 + e^{-i\tilde{\omega}\Delta t} &= 2C (e^{ikh} - 2 + e^{-ikh}) \\ 2\cos(\tilde{\omega}\Delta t) - 2 &= 2C^2 (2\cos(kh) - 2) \\ \cos(\tilde{\omega}\Delta t) &= 1 + 2C^2 \cos(kh) - 2C^2 \\ \cos(\tilde{\omega}\Delta t) &= \cos(kh) \end{aligned}$$