

# Beginnings in Algebraic Geometry Notes for CH 9 - 12

SFSU FALL 2026

## Contents

|          |                                                  |          |
|----------|--------------------------------------------------|----------|
| <b>1</b> | <b>CH 9 Projective Varieties</b>                 | <b>2</b> |
| 1.1      | 9.1 Projective Space . . . . .                   | 2        |
| 1.2      | 9.2 Projective Vanishing Operator . . . . .      | 3        |
| 1.3      | 9.3 Projective $\mathcal{I}$ -operator . . . . . | 4        |
| 1.4      | 9.4 Affine Restrictions . . . . .                | 5        |
| 1.5      | 9.5 Projective Closure . . . . .                 | 6        |
| 1.6      | 9.6 Project Nullstellensatz . . . . .            | 8        |

# 1 CH 9 Projective Varieties

## 1.1 9.1 Projective Space

**Definition 1.1** (Projective Space). Let  $n \in \mathbb{N}$  a projective space over  $K$  denoted  $\mathbb{P}_K^n$  is defined as

$$\frac{K^{n+1}/\{(0, \dots, 0)\}}{\sim}$$

where  $(a_0, \dots, a_n) \sim (b_1, \dots, b_n)$  iff there exists a  $\lambda \in K/\{0\}$  such that

$$(a_0, \dots, a_n) = (\lambda b_0, \dots, \lambda b_n).$$

An equivalence class is denoted  $[a_0 : \dots : a_n]$ .

*Remark 1.1.* Note here that we aren't really putting any structural requirements on  $K$ . My previous encounters with projective spaces almost always used a field and the notation of  $K$  is consistent with the textbooks use of fields. With affine spaces we noted that the Nullstellensatz was only consistent when  $K$  was an algebraically closed field. I imagine we will soon face a similar restriction. *Does  $K$  need to be a field?*

Projective spaces are particularly interesting because there are many related ways of presenting them. The first thing we'll do as a responsible mathematicians is go through a couple of examples.

**Ex** Let  $n = 0$ , that is  $\mathbb{P}_K^0$  for some  $K$ . This means we are looking at equivalence classes of  $\frac{K/\{0\}}{\sim}$ . Note that for any  $a \in K/\{0\}$  we can always take  $\lambda = \frac{1}{a}$  therefore all of  $\mathbb{P}^0 = \{[1]\}$ .

**Ex** Take  $n = 1$ , for any  $[a_0 : a_1] \in \mathbb{P}^1$  observe that by taking  $\lambda = \frac{1}{a_0}$  then every equivalence class where  $a_0 \neq 0$  is simply  $[1 : \frac{a_1}{a_0}]$ . On the other hand when the first term is zero,  $[0 : a_1]$  we may always choose  $\lambda = \frac{1}{a_1}$  so that it may be scaled to  $[0 : 1]$ . This means that  $\mathbb{P}^1$  can instead be expressed as a disjoint union

$$\mathbb{P}^1 = \{[1 : a] \mid a \in K\} \sqcup \{[0 : 1]\}.$$

*Observe that the left term is in bijection with  $\mathbb{A}^1$*

**Ex** Let  $n = 2$ . Similar to  $\mathbb{P}^1$ , equivalence classes  $[a_0 : a_1 : a_2] \in \mathbb{P}^2$  can always be rescaled to  $[1 : b_1 : b_2]$  when  $a_0 \neq 0$ . If the first term is zero then again every term can be scaled to  $[0 : b_1 : b_2]$ . Then,

$$\mathbb{P}^2 = \{[1 : b_1 : b_2] \mid b_1, b_2 \in K\} \sqcup \{[0 : b_1 : b_2] \mid b_1, b_2 \in K\}$$

Interestingly the by dropping the first term; the left portion is in bijection with  $\mathbb{A}^2$  while the right term is in bijection with  $\mathbb{P}^1$  which gives the decomposition

$$\mathbb{P}^2 = \mathbb{A}^2 \sqcup \mathbb{P}^1.$$

**Theorem 1.2.** For  $n \geq 1$  that there exists a natural bijection  $\mathbb{P}^n = \mathbb{A}^n \sqcup \mathbb{P}^{n-1}$

*Proof.* The proof follows the structure provided in problem 9.1.3 and consists of three portions. First define

$$U = \{[a_0 : \dots : a_n] \in \mathbb{P}^n \mid a_0, \dots, a_n \in K, a_0 \neq 0\} \quad \text{and} \quad V = \{[a_0 : \dots : a_n] \in \mathbb{P}^n \mid a_0, \dots, a_n \in K, a_0 = 0\}$$

We begin by showing that  $\mathbb{P}^n = U \sqcup V$ . The containment of  $\mathbb{P}^n$  in  $U \cup V$  and vice verse is automatic given the definition of  $U$  and  $V$ . It remains to show that  $U \cup V$  can be upgraded to  $U \sqcup V$ . Suppose towards a contradiction that  $U \cap V \neq \emptyset$  then there exists  $[b_0 : \dots : b_n] \in U \cap V$ . Note that for any  $[a_1 : \dots : a_n] \in U$ , because  $a_0$  is non-zero, we may always take  $\lambda = \frac{1}{a_0}$  so that every equivalence class contained in  $U$  is of the form  $[1 : c_1 : \dots : c_n]$  for some  $c_1, \dots, c_n \in K$ . The assumption that  $U \cap V$  is non-empty implies that  $b_0$  is both zero and one. Hence  $U \cap V = \emptyset$  and  $\mathbb{P}^n = U \sqcup V$ . The isomorphism  $V \simeq \mathbb{P}^{n-1}$  and  $U \simeq \mathbb{A}^n$  follow immediately by simply dropping the leading zero or one.  $\square$

There are a few more ways that projective spaces are represented. The most well known is that  $\mathbb{P}^n$  is isomorphic to the set of points through the origin of  $\mathbb{A}^{n+1}$

## 1.2 9.2 Projective Vanishing Operator

We say that a polynomial  $f \in K[x_0, \dots, x_n]$  only vanishes on a point  $a$  if it vanishes on every representative of  $a$  in the desired projective space.

**Non-Example:** Consider the polynomial  $f(x, y) = x^2 - y$  which vanishes on  $[2, 4]$  but it doesn't vanish on  $[4, 8]$

**Example:** The polynomial  $f(x, y) = x^2 - y^2$  on the other hand vanishes on  $[2, 2]$  and every possible representative of the solution

We end up with a problem: How do we check that every representative vanishes? Instead of checking we simply restrict ourselves to the case of homogenous polynomials. A polynomial is considered homogenous of degree  $d$  if every non-zero term of  $f$  has degree  $d$ . For example  $f(x, y, z) = 3x^2 + 2yz + 3zx$  is a polynomial homogenous of degree 2 because every non-zero term has degree 2.

**Proposition 1.1.** Let  $f \in K[x_0, \dots, x_n]$  be a homogenous polynomial of degree  $d$  and  $[a_0 : \dots : a_n] = [b_0 : \dots : b_n]$  in  $\mathbb{P}^n$  then  $f(a_0, \dots, a_n) = 0 \iff f(b_0, \dots, b_n) = 0$ .

*Proof.* If  $f$  is a homogenous polynomial of degree  $d$ , then each non-zero term of  $f$  takes the form  $cx_0^{\alpha_0} \dots x_n^{\alpha_n}$  for some non-zero  $c \in K$ ,  $\alpha_0, \dots, \alpha_n \in \mathbb{N}$  such that  $\sum \alpha_i = d$ . Evaluating  $f$  at a representative of  $[a_0 : \dots : a_n]$  means each non-zero term takes the form

$$c\lambda^{\sum \alpha_i} a_0 \dots a_n = c\lambda^d a_0 \dots a_n.$$

Then  $\lambda^d$  can be factored out of every non-zero term of  $f$  evaluating at so that  $f(\lambda a_0, \dots, \lambda a_n) = \lambda^d f(a_0, \dots, a_n)$ . If  $f(a_0, \dots, a_n) = 0$  and, by assumption, there exists a  $\lambda \in K$  such that

$$(\lambda a_0, \dots, \lambda a_n) = (b_1, \dots, b_n)$$

with the desired implication following directly. Following identical logic the reverse implication completes the desired if and only if. N

Of course in true algebraist sense, the fact that we can determine that a homogenous polynomial vanishes on all representatives as long as it vanishes on one, we restrict ourselves to looking only at homogenous polynomials. Yet we would like to still work with non-homogenous as a strict homogeneity condition is too restrictive. Lucky for us we can consider homogenous components of a polynomial. For example the polynomial  $f(x, y) = x^2 + xy + x$  has a homogenous component of degree 2 and degree 1 denoted

$$f_2(x, y) = x^2 + xy \quad \text{and} \quad f_1(x, y) = x$$

. The next proposition follows quite naturally

**Proposition 1.2.** Let  $f \in K[x_0, \dots, x_n]$  be a polynomial and  $a \in \mathbb{P}^n$ . Then  $f$  vanishes at  $a$  if and only if each homogenous component of  $f$  vanishes at  $a$ .

*Proof.* Suppose that each homogenous component  $f_k$  of  $f$  vanishes on  $a$  for all  $k \leq d$  where  $d$  is the degree of polynomial  $f$ . As

$$f = \sum_{k=1}^d f_k$$

and  $f_k(a) = 0$  for every representative of  $a$ , then  $f$  vanishes at every representative of  $a$ . Now suppose that  $f$  vanishes on every representative of  $a$ . Then for any  $\lambda \in K \setminus \{0\}$ , by assumption it follows

$$0 = f(\lambda a_0, \dots, \lambda a_n) \tag{1}$$

$$= \sum_{k=1}^d \lambda^d f_k(a_0, \dots, a_n). \tag{2}$$

Since (2) vanishes of infinitely many  $\lambda$  we may view (2) as a univariate polynomial

$$f(x) = \sum_{k=1}^d x^d f_k(a_0, \dots, a_n) = 0$$

vanishes on infinitely many values of  $K$ . This implies that  $f(x)$  must be the zero polynomial then,  $f_k(a_0, \dots, a_n) = 0$  for all  $k$ . Since  $f_k$  is homogenous then each homogenous component of  $f$  must vanish on every representative of  $a$ .  $\aleph$

**Definition 1.2** (Projective  $\mathcal{V}$  Operator). *Let  $\mathcal{S} \subseteq K[x_0, \dots, x_n]$  be a set of polynomials. The projective vanishes set of  $\mathcal{S}$  is*

$$\mathcal{V}_{\mathbb{P}}(\mathcal{S}) = \{a \in \mathbb{P}^n \mid f(a) = 0 \forall f \in \mathcal{S}\}$$

*We say that  $X \subseteq \mathbb{P}^n$  is projective variety if  $X = \mathcal{V}_{\mathbb{P}}(\mathcal{S})$  for some set  $\mathcal{S} \subseteq K[x_0, \dots, x_n]$ .*

Analogously to the affine case, a projective variety is defined by a ideals, i.e.

**Proposition 1.3.** *If  $\mathcal{S} \subseteq K[x_0, \dots, x_n]$  is a set of polynomials then*

$$\mathcal{V}_{\mathbb{P}}(\mathcal{S}) = \mathcal{V}_{\mathbb{P}}(\langle \mathcal{S} \rangle)$$

*Proof.* Let  $a \in \mathcal{V}_{\mathbb{P}}(\mathcal{S})$ , implies that for any polynomial  $f \in \mathcal{S}$  that  $f$  vanishes on  $a$ . Let  $g \in K[x_0, \dots, x_n]$  be a polynomial. The product  $fg$  evaluated at  $a$  must also vanish. Therefore  $fg \in \mathcal{S}$  hence  $\mathcal{S}$  must be an ideal. The reverse inclusion comes immediately.  $\aleph$

Further specifying, again like the affine case, we see that projective varieties are finitely generated by homogenous polynomials.

**Proposition 1.4.** *Any projective variety  $X \subseteq \mathbb{P}^n$  is of the form  $X = \mathcal{V}_{\mathbb{P}}(f_1, \dots, f_k)$  where  $f_1, \dots, f_k \in K[x_0, \dots, x_n]$  are homogenous polynomials.*

*Proof.* Let  $X \subseteq \mathbb{P}^n$  be a projective variety then  $X = \mathcal{V}_{\mathbb{P}}(\mathcal{S})$  where  $\mathcal{S} \subseteq K[x_0, \dots, x_n]$  is an ideal. Hilbert's Basis theorem ensures that  $K[x_0, \dots, x_n]$  is Noetherian implying there exist polynomials  $f_1, \dots, f_k \in K[x_0, \dots, x_n]$  such that  $\mathcal{S} = \langle f_1, \dots, f_k \rangle$ . Using lemma 9.16 note that if each polynomial  $f_i$  of degree  $d$  vanishes on a point  $a \in \mathbb{P}^n$  then every homogenous component  $f_{i,k}$  of degree  $k$  for  $k \leq d$  must also vanishes on  $a$ . Since  $f_i = \sum_p = 1^d f_{i,p}$  then we may further refine our list of generating polynomials. Hence any projective variety is the vanishing of a finite set of homogenous polynomials.  $\aleph$

### 1.3 9.3 Projective $\mathcal{I}$ -operator

It is defined essentially identically to the affine vanishing operator

**Definition 1.3** (Projective  $\mathcal{I}$ -operator). *Let  $X \subseteq \mathbb{P}^n$  be a subset. The vanishing ideal of  $X$  is the set*

$$\mathcal{I}_{\mathbb{P}}(X) = \{f \in K[x_0, \dots, x_n] \mid f(a) = 0 \forall a \in X\}$$

**Ex:** Let  $X = \{[1 : 0]\} \subseteq \mathbb{P}^n$ . Observe that for the polynomial  $f(x_0, x_1) = x_1$  any representative of  $[1 : 0]$  will always vanish. In fact we will show that  $\langle x_1 \rangle = \mathcal{I}_{\mathbb{P}}(X)$ . The inclusion  $\langle x_1 \rangle \subseteq \mathcal{I}_{\mathbb{P}}(X)$  follows immediately as  $x_1$  is fixed to zero. Now take  $f \in \mathcal{I}_{\mathbb{P}}(X)$  then  $f(a, 0) = 0$  for all  $a \in K$ , then we can view the polynomial as univariate in the first coordinate  $f(x_0, 0)$  with infinitely many zeroes implying it must be the zero polynomial. Take a polynomial  $f \in K[x_0, x_1]$  and view it as a univariate polynomial  $f \in (K[x_0])[x_1]$  which has a general form

$$f = \sum_{d \geq 0} f_d(x_0)x_1^d$$

If  $f$  vanishes on  $X$ , then  $f(x_0, 0) = 0$  implying that the degree zero homogenous component is also zero leaving us with an extra  $x_1$  term

$$f = x_1 \sum_{d \geq 1} f_d(x_0)x_1^{d-1} \in \langle x_1 \rangle.$$

**Ex:** Let  $X = \mathcal{V}(x_0 + x_1 - x_2) \subseteq \mathbb{P}^2$  then  $\mathcal{I}(X) = \langle x_0 + x_1 - x_2 \rangle$ . The ideal into vanishing ideal inclusion is immediate by definition. Now

**Definition 1.4.** An ideal  $I \subseteq K[x_0, \dots, x_n]$  is a homogenous ideal if for all  $f \in I$  every homogenous component of  $f$  is also in  $I$ .

In fact we can immediately generate a more useful and straightforward characterization of a homogenous ideal.

**Proposition 1.5.** An ideal  $I \subseteq K[x_0, \dots, x_n]$  is homogenous if and only if it admits a set of homogenous generators.

*Proof.* Suppose that  $I = \langle S \rangle$  where  $S$  is composed of all homogenous polynomials contained in  $I$ . Let  $f \in I$ , then  $f = \sum_{k \geq 0} f_k$ . Since  $f_k \in \langle S \rangle$  by construction then the sum  $f$  must also be contained in  $\langle S \rangle$ . The other inclusion follows immediately from the fact that  $S$  is a subset of the ideal  $I$ . Now suppose that  $I$  admits a set of homogenous generators, more precisely that  $I = \langle S \rangle$  where  $S$  is generated by a collection of homogenous polynomials. Let  $f \in I$ , then by definition

$$f = \sum_{i=1}^m g_i h_i$$

where  $g_i \in K[x_0, \dots, x_n]$  and  $h_i \in \langle S \rangle$  of homogenous degree  $d_i$  for all  $i$ . Observe that homogenous components of  $f$  of degree  $k$  take the form

$$f_k = \sum_{i=1}^m (g_i h_i)_k$$

. Note that every monomial in the product  $g_i h_i$  is a monomial of  $g_i$  multiplied by all of  $h_i$ . This means that every monomial of the product is of homogenous degree at least  $d_i$ . This ensures that for  $k < d_i$ , the homogenous component  $(g_i h_i)_k = 0$ . For  $k \geq d_i$ , monomials of homogenous degree  $k$  must be the product  $(g_i)_{k-d_i} h_i$  as every monomial of the product  $g_i h_i$  is a product of a monomial of  $g_i$  with  $h_i$ . This implies that  $(g_i h_i)_k \in I$  then  $f_k$  is a sum of elements of  $I$  implying that it must be contained in  $I$ . Therefore  $I$  must be a homogenous ideal.  $\aleph$

With this we can give a characterization of not just ideals, but vanishing ideals.

**Proposition 1.6.** If  $X \subseteq \mathbb{P}^n$  is any subset, then  $\mathcal{I}(X)$  is a homogenous radical ideal.

*Proof.* We begin by proving that  $\mathcal{I}(X)$  is a radical ideal. Suppose there exists an integer  $m$  such that  $f^m \in \mathcal{I}(X)$ , then for a point  $a \in X$

$$0 = f^m(a) = (f(a))^m.$$

Since  $K$  is an infinite field it has no zero divisors implying that  $f(a) = 0$  and that  $f \in \mathcal{I}(X)$ . Hence  $\mathcal{I}(X)$  is a radical ideal. Since  $f$  vanishes on  $a$  then every homogenous component of  $f$  must also vanish on  $a$  and is therefore contained in  $\mathcal{I}(X)$  implying that it is homogenous.  $\aleph$

With this we go on to establish identical properties to those of affine varieties related to the existence of unique irreducible decompositions of projective varieties. The proofs themselves follow exactly as in the affine case.

## 1.4 9.4 Affine Restrictions

We saw in the very first section of this chapter that a projective space  $\mathbb{P}^n$  could be viewed as  $\mathbb{A}^n \sqcup \mathbb{P}^{n-1}$ . Quite reasonably we ask whether a projective variety could be stratified in this way. Consider the projective variety  $X = \mathcal{V}_{\mathbb{P}}(x_0^2 + x_1^2 - x_2^2) \subseteq \mathbb{P}_{\mathbb{C}}^2$ . we saw that the affine component of a projective space is the collection of points where the first coordinate is non-zero or more succinctly when the first coordinate was fixed at 1. Similarly, fix  $x_0 = 1$ , then we have a subvariety

$$X_0 = \mathcal{V}_{\mathbb{A}}(1 + x_1^2 - x_2^2) \subseteq X$$

whose solutions are entirely contained in  $\mathbb{A}_{\mathbb{C}}^2$ . We give this phenomenon a more precise definition:

**Definition 1.5** (Affine Patch). For each  $i \in \{0, \dots, n\}$  the  $i$ -th affine patch of  $\mathbb{P}^n$  is the set

$$\mathbb{A}_i^n = \{[a_0 : \dots : a_n] \in \mathbb{P}^n \mid a_i \neq 0\}$$

and for any set  $X \subseteq \mathbb{P}^n$ , the intersection  $X \cap \mathbb{A}_i^n$  is called the  $i$ -th affine restriction of  $X$ .

Of course by simply dropping the  $i$ -th component, which is fixed at 1, we are in bijection with  $\mathbb{A}^n$ . Then the  $i$ -th restriction of a projective variety  $X = \mathcal{V}_{\mathbb{P}}(\mathcal{S})$  is  $\mathcal{V}_{\mathbb{A}}(\mathcal{S}_i)$  where

$$\mathcal{S}_i = \{f(x_0, \dots, x_{i-1}, 1, x_{i+1}, \dots, x_n) \mid f \in \mathcal{S}\}$$

## 1.5 9.5 Projective Closure

In the previous section we developed a way to take the global projective variety and investigate restriction to local affine patches. Now we aim to develop a method to take some affine patches and join them together into a projective variety. Notice though that in the process of restricting to affine patches we lose the points at infinity. So if we were to start with affine patches and attempt to move back into a projective variety, we would need to devise a way to add point at infinity and ensure that the resulting set is actually a projective variety. To do this we define the projective closure.

**Definition 1.6** (Projective Closure). Let  $X \subseteq \mathbb{A}^n$  be an affine variety, and let  $j_0(X)$  be the image of  $X$  in the first affine patch  $\mathbb{A}_0^n \subseteq \mathbb{P}^n$ . The projective closure of  $X$ , denoted  $\overline{X}$  is the intersection of all projective varieties that contain  $j_0(X)$ .

*Remark 1.3.* Observe that, under the zariski topology, this is exactly the definition of a closed set.

- Consider the affine variety  $\mathcal{V}_{\mathbb{A}}(1 + x_1 - x_2) \subseteq \mathbb{A}^2$  which has the vanishing set  $\{(a, 1 + a) : a \in K\}$ . Under the josure,  $j_0$  we can embed this variety as a set in  $\mathbb{P}^2$  defined as

$$j_0(X) = \{[1 : a : a + 1] : a \in K\} \subseteq \mathbb{P}^2.$$

Observe that  $[1 : a : a + 1] = [\frac{1}{a} : 1 : 1 + \frac{1}{a}]$  means that  $a$  must be further restricted to  $a \in K \setminus \{0\}$ . Take  $b = \frac{1}{a}$  then we can rewrite the elements of  $j_0(X)$  to be  $[b : 1 : 1 + b]$ . Then in terms of our polynomial, for every point in  $j_0(X)$ ,  $f(b, 1, 1 + b) = 0$  can be viewed as a univariate polynomial  $f(x, 1, 1 + x) = 0$  which means that  $f$  is the univariate polynomial. If we were to treat  $j_0(X)$  as a projective variety then we have a univariate polynomial vanishing on  $\mathbb{P}^2 = \mathbb{A}^2 \sqcup \mathbb{P}^1$ . In particular this polynomial is vanishing on  $\mathbb{P}^1$  which in this case is only the point  $[0 : 1]$  then  $f(0, 1, 1)$  must be a solution so we have the point  $[0 : 1 : 1]$  as a solution. So we determine that we must augment  $j_0(X)$  as  $j_0(X) \cup \{[0 : 1 : 1]\}$ . Note that we STILL don't yet know whether this is a projective variety because restricting to the first affine patch agrees with  $j_0(X)$  but we don't know how this variety is changed when restricting to different affine patches. This means that we must determine the polynomial that vanishes and luck for us there is one that I am too lazy to compute so I will take from the book:

$$\overline{X} = \mathcal{V}_{\mathbb{P}}(x_0 + x_1 - x_2)$$

- Let  $\mathcal{V}_{\mathbb{A}}(x_2 + x_1^2) \subseteq \mathbb{A}^2$  has josure

$$j_0(X) = \{[1 : a : a^2] \mid a \in K\}$$

which under  $b = \frac{1}{a}$  can be viewed as the set of points  $\{[b^2 : b : 1] : b \in K \setminus \{0\}\}$ . Similarly we must add a single point and then determine whether this point with the josure is actually the vanishing of the polynomial. It is hence  $\overline{X} = \mathcal{V}_{\mathbb{P}}(x_0x_1 - x_2^2)$ .

Observe that the resulting polynomial of the variety is now of some homogenous degree. In fact, as we will state soon, the projective closure of an affine variety is always the "homogenization" of the ideal generating the affine variety. This gives us a much simpler characterization of the projective closure than the rather abstract intersection of all varieties containing the josure.

**Definition 1.7.** Let  $f \in K[x_1, \dots, x_n]$  be a polynomial of degree  $d$ . The homogenization of the polynomial is define as

$$\overline{f} = x_0^d f\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in K[x_0, \dots, x_n].$$

Observe that if we were to look at the josure of a homogenized polynomial, that is we evaluate  $\bar{f}(1, x_1, \dots, x_n)$  we will get back our unhomogenized polynomial  $f$ . Reframing our previous examples we would like to determine whether if  $X = \mathcal{V}_{\mathbb{A}}(f_1, \dots, f_k)$  that  $\bar{X} = \mathcal{V}_{\mathbb{P}}(\bar{f}_1, \dots, \bar{f}_k)$  yet the accursed twisted cubic will laugh in our face. I don't feel like walking through this example so I will make some general remarks about it instead. The affine twisted cubic  $\mathcal{V}_{\mathbb{A}}(x_2 - x_1^2, x_3 - x_1^3)$  will have a josure  $j_0(X) = \{[1 : a : a^2 : a^3] \mid a \in K\} \subseteq \mathbb{P}^3$  and to make it a projective variety we only need to add the point  $[0 : 0 : 0 : 1]$ . If we were to look at a variety defined by the homogenization of the polynomials then we would arrive at  $j_0(X) \sqcup \{0 : 0 : b : c\}$ , i.e. we would add too many points. Instead we actually another defining polynomial rather than simply homogenizing the defining polynomials of the affine variety. So instead of homogenizing some polynomials we will actually just homogenize the entire ideal.

**Definition 1.8** (Homogenization of a Set). *Let  $\mathcal{S} \subset K[x_1, \dots, x_n]$  be a set of polynomials. The homogenization of  $\mathcal{S}$  is the set*

$$\bar{\mathcal{S}} = \{\bar{f} : f \in \mathcal{S}\} \subseteq K[x_0, \dots, x_n]$$

One thing to note is that generally the homogenization of an ideal is not also an ideal. If we were to take a product  $g\bar{f}$  where  $g \in K[x_0, \dots, x_n]$  and inhomogenous and  $\bar{f} \in \bar{\mathcal{S}}$  then our product is not homogenous and so can not be in  $\bar{\mathcal{S}}$ . With the homogenization of a set we get the right information to recharacterize the affine closure in a way that plays better with the vanishing and ideal operators.

**Proposition 1.7.** *If  $X \subseteq \mathbb{A}^n$  and affine variety then  $\bar{X} = \mathcal{V}_{\mathbb{P}}(\overline{\mathcal{I}_{\mathbb{A}}(X)}) \subseteq \mathbb{P}^n$ .*

*Proof.* ( $\subseteq$ ) Let  $[1 : a_1 : \dots : a_n] \in j_0(X)$  and  $g \in \overline{\mathcal{I}_{\mathbb{A}}(X)}$  then  $g = \bar{f}$  for some  $f \in \mathcal{I}_{\mathbb{A}}(X)$  and  $g(1, a_1, \dots, a_n) = f(a_1, \dots, a_n) = 0$  then  $j_0(X)$  is contained in  $\mathcal{V}_{\mathbb{P}}(\overline{\mathcal{I}_{\mathbb{A}}(X)})$ . Since  $\bar{X}$  is the smallest projective variety to contain  $j_0(X)$  then  $\bar{X}$  must be contained in  $\mathcal{V}_{\mathbb{P}}(\overline{\mathcal{I}_{\mathbb{A}}(X)})$ . ( $\supseteq$ ) It suffices to show that

$$\mathcal{I}_{\mathbb{P}}(\bar{X}) \subseteq \mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(\overline{\mathcal{I}_{\mathbb{A}}(X)}))$$

because application of the projective vanishing operator will give us the desired containment. Suppose that  $g \in \mathcal{I}_{\mathbb{P}}(\bar{X})$ , as  $g$  is contained in a homogenous ideal it must also be homogenous. The homogenous polynomial  $g$  is not necessarily the homogenization of a polynomial so to force this condition let  $k \in \mathbb{N}$  be the maximal choice such that  $g = x_0^k h$  for some  $h \in K[x_0, \dots, x_n]$ . Observe that forcing  $k$  to be maximal implies that from at least one monomial of  $g$  we must be able to factor the entirety of an  $x_0$  component out, if this weren't the case then there would be some  $x_0$  terms in every monomial and  $k$  would not be maximal. Since  $g$  is homogenous of some degree  $d$ , then this monomial has degree  $d - k$ . Note that this monomial must have minimal degree after factoring as any term with some  $x_0$  has less remaining degree distributed over terms  $x_i$  where  $i \neq 0$ . Take  $h_0 = h(1, x_1, \dots, x_n)$  and observe that this removes the  $x_0$  term from every monomial and utilizing the conclusion of the previous two sentences this means that the monomial of degree  $d - k$  is now of maximal degree in  $h_0$ . The  $\bar{h}_0 = x_0^{d-k} h_0$  which returns us back to the polynomial  $h$ . Since  $g$  vanishes on all of  $\bar{X}$  then it must vanish on all of  $j_0(X)$ , in particular this means that  $g$  must vanish on all of  $X$ . Since

$$h_0(a_1, \dots, a_n) = h(1, a_1, \dots, a_n) = g(1, a_1, \dots, a_n) = 0$$

we see that  $h_0 \in \mathcal{I}_{\mathbb{A}}(X)$  and by definition  $h \in \overline{\mathcal{I}_{\mathbb{A}}(X)}$  and that  $g \in \langle \overline{\mathcal{I}_{\mathbb{A}}(X)} \rangle$ . Then

$$g \in \langle \overline{\mathcal{I}_{\mathbb{A}}(X)} \rangle \subseteq \mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(\overline{\mathcal{I}_{\mathbb{A}}(X)}))$$

and we are done. \(\square\)

While we can't in general conclude that  $\overline{\mathcal{V}_{\mathbb{A}}(f_1, \dots, f_k)} = \mathcal{V}_{\mathbb{P}}(\bar{f}_1, \dots, \bar{f}_k)$  we get a special case when  $i = 1$ .

**Proposition 1.8.** *If  $f \in K[x_0, \dots, x_n]$  then  $\overline{\mathcal{V}_{\mathbb{A}}(f)} = \mathcal{V}_{\mathbb{P}}(\bar{f})$ .*

*Proof.* Let  $X = \mathcal{V}_{\mathbb{A}}(f)$ , the Nullstellensatz says that  $\mathcal{I}_{\mathbb{A}}(\mathcal{V}_{\mathbb{A}}(X)) = \langle q_1 \cdots q_{\ell} \rangle$  where  $f = q_1^{\alpha_1} \cdots q_{\ell}^{\alpha_{\ell}}$  is an irreducible decomposition of  $f$ . Using the previous proposition  $\overline{\mathcal{V}_{\mathbb{A}}(X)} = \mathcal{V}_{\mathbb{P}}(\overline{\mathcal{I}_{\mathbb{A}}(X)}) = \mathcal{V}_{\mathbb{P}}(\langle \overline{q_1 \cdots q_{\ell}} \rangle)$ . It remains to show that this is equivalent to  $\mathcal{V}_{\mathbb{P}}(\bar{f})$ . Let  $f \in \langle q_1 \cdots q_{\ell} \rangle$  then  $\bar{f} \in \langle \overline{q_1 \cdots q_{\ell}} \rangle$  which implies that

$$\mathcal{V}_{\mathbb{P}}(\bar{f}) \supseteq \mathcal{V}_{\mathbb{P}}(\overline{\langle q_1 \cdots q_{\ell} \rangle}).$$

Now suppose that  $a \in \mathcal{V}_{\mathbb{P}}(\overline{\langle q_1 \cdots q_\ell \rangle})$  then for some  $\bar{f} \in \overline{\langle q_1 \cdots q_\ell \rangle}$ ,  $\bar{f}(a) = 0$ . Note that  $\bar{f} = \bar{q}_1^{\alpha_1} \cdots \bar{q}_\ell^{\alpha_\ell}$ . The fact that  $\bar{f}$  vanishes on  $a$  implies that  $\bar{q}_i^{\alpha_i}$  vanishes on  $a$  for some  $i$ . Then for any  $g \in \overline{\langle q_1 \cdots q_\ell \rangle}$  it follows that  $g$  must also vanish on  $a$  because it will have the vanishing homogenized  $q_i$  component.  $\aleph$

One thing that we may have noticed in the examples working through projective closures was the fact that we never seemed to be adding affine points, only points at infinity. In fact these are examples of the following proposition.

**Proposition 1.9.** *The affine restriction of the projective closure of an affine variety  $X$  is always  $X$ .  $(\bar{X})_0 = X$ .*

*Proof.* Let  $X \subseteq \mathbb{A}^n$  be an affine variety. Define  $\mathcal{S} = \overline{\mathcal{I}_{\mathbb{A}}(X)}$  and note that  $\bar{X} = \mathcal{V}_{\mathbb{P}}(\mathcal{S})$ . Then the affine restriction  $(\bar{X})_0$  vanishes on a set of polynomials

$$\mathcal{S}_0 = \{g(1, a_1, \dots, a_n) \mid g \in \mathcal{S}\} = \{f(a_1, \dots, a_n) \mid f \in \mathcal{I}_{\mathbb{A}}(X)\}$$

where the second equality comes from the fact that  $g$  must be the homogenization of some polynomial  $f$  and evaluating  $g$  at  $x_0 = 1$  simply returns us to evaluation  $f$  at  $a_1, \dots, a_n$ . The final set is just  $\mathcal{I}_{\mathbb{A}}(X)$  then  $\mathcal{V}_{\mathbb{A}}(\mathcal{I}_{\mathbb{A}}(X)) = X$  and we are done.  $\aleph$

## 1.6 9.6 Project Nullstellensatz

If we are to take inspiration from the affine NS then we aim to prove that

$$\mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(I)) = \sqrt{I}$$

for some ideal  $I \in K[x_0, \dots, x_n]$ . But we have a problem here if  $I = \langle x_0, x_1, \dots, x_n \rangle$ . The only point that vanishes on a polynomial in here would be  $[a_0 : \cdots : a_n]$  where each  $a_i = 0$ . Since there is no such point in  $\mathbb{P}^n$  then  $\mathcal{V}_{\mathbb{P}}(I) = \emptyset$ . Yet every polynomial vanishes on the empty set hence  $\mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(I)) = K[x_0, \dots, x_n] \neq I$  where we note that  $I$  is already radical. So with this we aim to disregard this annoying ideal.

**Definition 1.9.** *An ideal  $I \subseteq K[x_0, \dots, x_n]$  is irrelevant if  $I \supseteq \langle x_0, \dots, x_n \rangle$ . Otherwise  $I$  is called relevant.*

With this the projective NS is the affine NS reframed with these classification.

**Proposition 1.10.** *Let  $I \subseteq K[x_0, \dots, x_n]$  be a homogenous ideal.*

1.  $\mathcal{V}_{\mathbb{P}}(I) = \emptyset$  if and only if  $\sqrt{I} \supseteq \langle x_0, \dots, x_n \rangle$ .
2. If  $\sqrt{I}$  is relevant then  $\mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(I)) = \sqrt{I}$ .

But before proving this we will utilize a different way to move between affine and projective varieties through cones.

**Definition 1.10.** *Let  $X \subseteq \mathbb{P}^n$  be a subset. The affine cone over  $X$  is the set*

$$C(X) = \{(0, \dots, 0)\} \cup \{(a_0, \dots, a_n) \mid [a_0 : \cdots : a_n] \in X\} \subseteq \mathbb{A}^{n+1}$$

With the affine cone in mind we'll prove the following proposition that allow interchanges between the affine and projective vanishing and ideal operators.

**Proposition 1.11.** *Let  $I \subseteq K[x_0, \dots, x_n]$  be a homogenous ideal then*

$$\mathcal{V}_{\mathbb{A}}(I) = C(\mathcal{V}_{\mathbb{P}}(I))$$

*Proof.* Since we are looking to show that  $\mathcal{V}_{\mathbb{A}}(I)$  is a cone then we must first show that for any homogenous ideal that this variety admits  $(0, \dots, 0)$ . Suppose that it did not admit the origin, this means that there exists a homogenous polynomial  $f \in I$  such that  $f(0, \dots, 0) \neq 0$ . This is only possible if the polynomial is non-zero and constant which forces  $I$  to admit such a generator. But if it were to admit such a generator then  $I = K[x_0, \dots, x_n]$  contradicting our choice of  $I$ . Since each polynomial in  $I$  is homogenous, if it were to vanish on a point  $(a_0, \dots, a_n)$  then it must vanish on any representative  $[a_0 : \cdots : a_n]$  and vice versa hence

$$(a_0, \dots, a_n) \in \mathcal{V}_{\mathbb{A}}(I) \iff [a_0 : \cdots : a_n] \in \mathcal{V}_{\mathbb{P}}(I) \iff (a_0, \dots, a_n) \in C(\mathcal{V}_{\mathbb{P}}(I))$$

therefore we are done.  $\aleph$



**Proposition 1.12.** *Let  $X \subseteq \mathbb{P}^n$  be a non-empty subset. Then*

$$\mathcal{I}_{\mathbb{P}}(X) = \mathcal{I}_{\mathbb{A}}(C(X)).$$

*Proof.* Note that  $\mathcal{I}_{\mathbb{P}}(X)$  is a non-empty homogenous ideal which means that none of the generating polynomials can be constant. Then on any point  $(0, \dots, 0) \in \mathbb{A}^{n+1}$  a polynomial in this projective vanishing ideal might vanish. Similarly the fact that a polynomial vanishes on the projective variety  $X$  means that it must vanish on its affine restriction. Then we can redefine the projective vanishing ideal as

$$\mathcal{I}_{\mathbb{P}}(X) = \{f \in K[x_0, \dots, x_n] \mid f(a_0, \dots, a_n) = 0 \text{ if } [a_0 : \dots : a_n] \in X, f(a_0, \dots, a_n) = 0 \text{ if } (a_0, \dots, a_n) = (0, \dots, 0)\} \quad (3)$$

$$= \mathcal{I}_{\mathbb{A}}(\{0, \dots, 0\} \cup \{(a_0, \dots, a_n) \mid [a_0 : \dots : a_n] \in X\}) \quad (4)$$

$$= \mathcal{I}_{\mathbb{A}}(C(X)) \quad (5)$$

⋈

Now that we have the ability to move between projective and affine varieties and vanishing ideals through cones we can prove the projective NS.

*Proof.* (a) Suppose that  $\sqrt{I} \supseteq \langle x_0, \dots, x_n \rangle$ . Note that if  $a \in \mathcal{V}_{\mathbb{P}}(I)$  then there exists some polynomial  $f \in I$  such that  $f(a) = 0$  then  $f \in \sqrt{I}$  hence  $a \in \mathcal{V}_{\mathbb{P}}(\sqrt{I})$ . Suppose that  $a \in \mathcal{V}_{\mathbb{P}}(\sqrt{I})$  then for some  $m > 0$  there exists an  $f^m \in \sqrt{I}$  such that  $f^m(a) = 0$ . By definition it follows that  $f \in I$  and as  $f^m(a) = f^{m-1}(a)f(a) = 0$  it follows that  $f(a) = 0$ . Therefore  $\mathcal{V}_{\mathbb{P}}(I) = \mathcal{V}_{\mathbb{P}}(\sqrt{I})$ . Then

$$\mathcal{V}_{\mathbb{P}}(I) = \mathcal{V}_{\mathbb{P}}(\sqrt{I}) \subseteq \mathcal{V}_{\mathbb{P}}(\langle x_0, \dots, x_n \rangle) = \emptyset.$$

Now suppose that  $\mathcal{V}(I) = \emptyset$ , then either  $I = K[x_0, \dots, x_n]$  in which case  $\sqrt{I} = I \supseteq \langle x_0, \dots, x_n \rangle$ . Suppose that  $I \subsetneq K[x_0, \dots, x_n]$ . Utilizing prop 1.11

$$\mathcal{V}_{\mathbb{A}}(I) = C(\mathcal{V}_{\mathbb{P}}(I)) = C(\emptyset) = \{(0, \dots, 0)\}.$$

Under the affine NS

$$\sqrt{I} = \mathcal{I}_{\mathbb{A}}(\mathcal{V}_{\mathbb{A}}(I)) = \mathcal{I}_{\mathbb{A}}(\{(0, \dots, 0)\}) = \langle x_0, \dots, x_n \rangle$$

(b) Now assume that  $\sqrt{I}$  is relevant. Suppose that  $\mathcal{V}_{\mathbb{P}}(I) = \emptyset$ . The first case tells us that  $\sqrt{I} \supseteq \langle x_0, \dots, x_n \rangle$  but the only relevant ideal that can contain the maximal irrelevant ideal is the full ring i.e.  $\sqrt{I} = K[x_0, \dots, x_n]$ . Then  $\mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(I)) = \mathcal{I}_{\mathbb{P}}(\emptyset) = K[x_0, \dots, x_n]$  and we are done. Now suppose that  $\mathcal{V}_{\mathbb{P}}(I) \neq \emptyset$  which means that  $I \subsetneq K[x_0, \dots, x_n]$ . Then,

$$\mathcal{I}_{\mathbb{P}}(\mathcal{V}_{\mathbb{P}}(I)) = \mathcal{I}_{\mathbb{A}}(C(\mathcal{V}_{\mathbb{P}}(I))) = \mathcal{I}_{\mathbb{A}}(\mathcal{V}_{\mathbb{A}}(I)) = \sqrt{I}$$

hence we are done.

⋈