

The Gap Alphabet

Paper 5. Which gaps can occur between consecutive odd composites, and where the ladder of proofs stops

Abstract

Between consecutive odd composites the gap takes only the values 2, 4 and 6, and a gap of 6 *is* a twin pair. That single observation turns the twin problem into a question about one letter of a three-letter alphabet, and this paper settles everything about that alphabet that can be settled without prime input: the two short gaps occur infinitely often for elementary reasons, the long one is equivalent to the conjecture itself, and the forbidden words of the language are classified. On one rail no five survivors above 5 lie in arithmetic progression of step 6, and the run counts obey an exact product formula that caps runs at four. **No progress toward the twin-prime conjecture is claimed.**

Numbering. This paper is one part of a set that was written as a single document and is now published in parts. Each part numbers its own results from one and is self-contained: a reference of the form [Pn, Thm 1] means Theorem 1 of Paper n, and an unqualified “Theorem 1” always means this paper’s own. Result numbers therefore differ from those of releases before 2.0.0, where the whole set shared the numbering of the single document.

How to read the claims in this paper. Statements set as Theorems, Propositions and Corollaries are proved, and the proofs are given. One result in the set is labelled **Verified Law**: it is proved under a stated hypothesis and verified numerically outside it. Anything described as *measured* is a computation over a stated finite range and is labelled as such where it occurs.

Keywords: twin primes, gap alphabet, sieve cycles, cell coordinates.

MSC 2020: 11N35, 11N05, 11A41.

Summary of the results in this part

§2 — The gap alphabet, and the ladder of proofs

Result	What it says	Section
Theorem 1	The gap between consecutive odd composites takes only the values 2, 4 or 6.	§2.1
Theorem 2	Gaps 2 and 4 occur infinitely often with no prime input; gap 6 infinitely often is <i>equivalent</i> to the twin conjecture.	§2.2
Theorem 3	On one rail, no five survivors above 5 lie in arithmetic progression of step 6.	§2.3
Corollary 1	The run counts G_k at spacing 6 obey $G_k = \prod (q - k)$, capping runs at four.	§2.3

1. Setting

This paper is short and takes almost nothing from the others. [P2] supplies the coordinates: the line $L_p(k) = p(p + 2k)$ beginning at p^2 , the grid L_3 of step 6, and the cells $C_b = (6b - 1, 6b + 1)$. Theorems 1, 2

and 3 use those and nothing more; Theorem 3 needs only that L_5 begins at 25. Corollary 1 is the one place where the cycle enters: the count of runs over a full period is the product law of [P3].

Two companions are cited but not used as sources. [P9] is where the obstruction that stops the ladder is derived, and [P11] is where the routes around it are collected. Neither is needed to read this paper.

A warning about the word “gap”, carried here because this is the paper that fixes the alphabet. A *gap* below always means the numerical distance between two consecutive odd composites, so the letter 6 is a twin pair. That is not the same object as an *edge of length 6* between two survivors, which is a prime pair $(p, p + 6)$ and is studied in [P7, App. B]. The two must not be run together: the phrase “gap 6” would otherwise cover both.

2. The gap alphabet and the ladder of proofs

Terms used throughout, all as defined in [P2]. A **line** L_p is the odd multiples of p from p^2 onward; a **cell** is a pair $C_b = (6b - 1, 6b + 1)$; a **strike** is a member of a cell that lies on some line; a member is a **survivor** of a set of lines if none of them strikes it, and a cell is **open** if both its members survive. A **gap** here always means the numerical distance between two consecutive odd composites, not a count of cells. A **sector** is the interval between consecutive odd squares; a **belt** is the interval between the squares of consecutive primes.

We record first what the framework proves outright about gaps, and then where the ladder of proofs stops.

2.1 Theorem 1 (the gap alphabet)

Theorem 1. The gap between consecutive odd composites takes only the values 2, 4 or 6.

Proof. Among any three consecutive odd numbers the residues modulo 3 are a permutation of $\{0, 1, 2\}$:

$$n, n + 2, n + 4 \longrightarrow \text{one of them lies on } L_3.$$

So, so one is divisible by 3; and every odd multiple of 3 from 9 onward lies on L_3 . Hence beyond 7 no three consecutive survivors exist, and the gap is capped at 6. ■

Correspondingly: gap 2 means no survivor between; gap 4 means one isolated prime; and **gap 6 means two adjacent survivors, i.e. a twin pair**. Verified: every gap-6 interval contains a twin, zero failures among 2,992 instances below 3×10^5 .

Measured over 9×10^8 gaps up to 2×10^9 , the three letters occur with frequencies 0.89816, 0.09475 and 0.00708; the maximum observed gap is 6, first attained at 9.

Hence an equivalent form of the twin conjecture: **the maximal possible gap is attained infinitely often**. The cap is free and proved; the statement concerns composites, which are the objects the framework actually constructs.

2.2 The ladder, and where it stops

Theorem 2. Gap 2 occurs infinitely often, requiring no prime; gap 4 occurs infinitely often, requiring one prime.

Proof. For gap 2 take

$$n = 30j + 3, \quad n + 2 = 30j + 5.$$

The first lies on L_3 , the second on L_5 , both for every $j \geq 1$. For gap 4: let $p \equiv 8 \pmod{15}$ be prime; then $p - 2 \equiv 6$ is divisible by 3 and $p + 2 \equiv 10$ by 5, so p is an isolated survivor, and Dirichlet's theorem supplies infinitely many such p . ■

gap 2	zero primes required	proved
gap 4	one prime (Dirichlet)	proved
gap 6	two primes simultaneously	open

Dirichlet supplies one prime in an arithmetic progression; nothing supplies two at a prescribed distance. That is the whole of the remaining distance.

Why this reciprocal-sum test cannot bridge it. Summing reciprocals by gap type:

N	gap 2	gap 4	gap 6
10^6	2.805	0.887	0.464
10^8	4.556	1.127	0.488
2×10^9	5.762	1.257	0.498

Gaps 2 and 4 have **divergent** reciprocal sums (growing like $\log \log N$), and divergence proves infinitude. The gap-6 sum **converges** — flattening toward a constant. That constant is the analogue of Brun's constant in the present alphabet and not Brun's constant itself: the latter is $\sum_{\text{twins}} (1/p + 1/(p+2)) = 1.9021605\dots$, whereas the column above carries one reciprocal per gap-6 event. A convergent reciprocal sum cannot distinguish “infinitely many” from “finitely many”. Thus this particular divergence-based density test has no route to the twin conclusion.

2.3 Runs and the cap at four

Theorem 3. On a single rail, no five survivors with $x > 5$ can lie in arithmetic progression with common difference 6; and 5, 11, 17, 23, 29 is the only exception.

Proof. Since $6 \equiv 1 \pmod{5}$, the five terms

$$x, \quad x+6, \quad x+12, \quad x+18, \quad x+24$$

run through all residues modulo 5, so exactly one of them is divisible by 5. If that term exceeds 5 it is an odd multiple of 5 at least 25, hence lies on L_5 and is struck. The term can fail to be struck only when it equals 5 itself, which forces $x = 5$. ■

The exception is real and must be carried in the statement. For $x = 5$ the run is 5, 11, 17, 23, 29, and all five are prime; an exhaustive search to 2×10^5 finds this and no other. It exists purely because L_5 is born at 25: as a statement about the *residue classes* the argument is exact, and only the birth rule creates the exception.

Verification. At sieve depth 97 over 6×10^7 cells, the run-length census is 6,448,150 of length 1; 2,404,092 of length 2; 778,762 of length 3; 211,170 of length 4; and **none of length 5 or more**.

Corollary 1. With G_k the number of runs of k consecutive survivors at spacing 6 on one rail, the full-cycle counts satisfy $G_k = \prod_q (q - k)$.

Proof. A run of k forbids exactly k marks on each ruler, leaving $q - k$. ■

Verification (direct count on the t -axis at $p = 101$):

lines	$k=1$	2	3	4	5
$\{5\}$	4	3	2	1	0
$\{5, 7\}$	24	15	8	3	0
$\{5, 7, 11\}$	240	135	64	21	0

The four laws $(q-1), \dots, (q-4)$ are therefore not four phenomena but one ruler with different numbers of marks.

The computations and much of the prose in this paper were prepared with AI assistance (ChatGPT, OpenAI; Claude, Anthropic), used for algebraic derivation, for drafting and rewriting code and text, for running the computations, and for auditing the papers against their own scripts. All statements were checked by the author, who is responsible for them; the repository README sets out the division of labour in full.

References

The eleven papers of this set are cited as [P1] to [P11]. This paper uses no external reference: everything it quotes is a definition or a statement of a companion paper, never a proof.