

DON'T SAY CAT: NEW ITEM RESPONSE THEORY APPROACHES FOR DEVELOPING SHORT TEST FORMS

Ottavia M. Epifania^{1,2}, Pasquale Anselmi¹, Egidio Robusto¹

ottavia.epifania@unipd.it

¹University of Padova

²Catholic University of the Sacred Heart

September 30th 2022, Padova

XXX Annual Conference of the Italian Psychology Association (AIP)



1 Introduction

2 Item Response Theory and information functions

3 IRT procedures for shortening tests

Benchmark procedure

Procedures based on θ targets

4 Simulation study

5 Some final remarks

CAT



CAT



Computerized Adaptive Testing

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., for each respondent) → (**CAT**: Computerized Adaptive Testing)

STATIC SHORT FORMS: Static tests equal for all respondents → The information is maximized across θ levels (i.e., across all respondents)

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., for each respondent) → (**CAT**: Computerized Adaptive Testing)

Issue

Different short test forms for each respondent → Potential fairness issues in assessments, e.g. for recruitment

STATIC SHORT FORMS: Static tests equal for all respondents → The information is maximized across θ levels (i.e., across all respondents)

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., for each respondent) → (**CAT**: Computerized Adaptive Testing)

Issue

Different short test forms for each respondent → Potential fairness issues in assessments, e.g. for recruitment

STATIC SHORT FORMS: Static tests equal for all respondents → The information is maximized across θ levels (i.e., across all respondents)

Issue

Not being tailored to any θ level of interest → Potentially more items are needed to cover a wide range of θ s

Aim

New IRT-based procedures for shortening tests

Aim


New IRT-based procedures for shortening tests



Equal for all respondents

Aim

New IRT-based procedures for shortening tests



Equal for all respondents

Tailored to specific levels of
the latent trait

1 Introduction

2 Item Response Theory and information functions

3 IRT procedures for shortening tests

Benchmark procedure

Procedures based on θ targets

4 Simulation study

5 Some final remarks

Item Response Theory

2-PL Model

$$P(x_{pj} = 1 | \theta_p, b_j, a_j) = \frac{\exp[a_j(\theta_p - b_j)]}{1 + \exp[a_j(\theta_p - b_j)]}$$


where:


$P(x_{pj} = 1)$: Probability of a correct response to item j by respondent p


θ_p : Ability of respondent p


b_j : Difficulty of item j

a_j : Discrimination of item j









Information functions

Item Information Function

$$IIF_j = a_j^2 [P(\theta)(1 - P(\theta))]$$

Information functions

Item Information Function

$$IIF_j = a_j^2 [P(\theta)(1 - P(\theta))]$$




Figure 1: $a = 0.20$, $a = 0.70$, $a = 1.90$,
 $b = 0$

Information functions

Item Information Function

$$IIF_j = a_j^2 [P(\theta)(1 - P(\theta))]$$

Test Information Function

$$TIF = \sum_{j=1}^J IIF_j$$




Figure 1: $a = 0.20$, $a = 0.70$, $a = 1.90$,
 $b = 0$

Information functions

Item Information Function

$$IIF_j = a_j^2 [P(\theta)(1 - P(\theta))]$$




Figure 1: $a = 0.20$, $a = 0.70$, $a = 1.90$,
 $b = 0$

Test Information Function

$$TIF = \sum_{j=1}^J IIF_j$$




Figure 2: $TIF = IIF_1 + IIF_2 + IIF_3$

1 Introduction

2 Item Response Theory and information functions

3 IRT procedures for shortening tests

Benchmark procedure

Procedures based on θ targets

4 Simulation study

5 Some final remarks

- └ Short form procedures
- └ Benchmark procedure

1 Introduction

2 Item Response Theory and information functions

3 IRT procedures for shortening tests

Benchmark procedure

Procedures based on θ targets

4 Simulation study

5 Some final remarks

Benchmark procedure

Selected items \rightarrow items with the highest *IIFs*

e.g.: 3-item short form from 10-item full-length test

| item | <i>b</i> | <i>a</i> | <i>IIF</i> |
|------|----------|----------|------------|
| 1 | -0.67 | 0.71 | 0.08 |
| 2 | 0.50 | 1.19 | 0.15 |
| 3 | -2.43 | 0.25 | 0.01 |
| 4 | 2.12 | 1.98 | 0.24 |
| 5 | 1.72 | 0.39 | 0.03 |
| 6 | -2.28 | 1.62 | 0.19 |
| 7 | 0.64 | 0.50 | 0.05 |
| 8 | -2.51 | 1.68 | 0.19 |
| 9 | -0.66 | 0.44 | 0.04 |
| 10 | 0.72 | 0.33 | 0.02 |

Benchmark procedure

Selected items → items with the highest *IIFs*

e.g.: 3-item short form from 10-item full-length test

| item | <i>b</i> | <i>a</i> | <i>IIF</i> |
|------|----------|----------|------------|
| 4 | 2.12 | 1.98 | 0.24 |
| 8 | −2.51 | 1.68 | 0.19 |
| 6 | −2.28 | 1.62 | 0.19 |
| 2 | 0.50 | 1.19 | 0.15 |
| 1 | −0.67 | 0.71 | 0.08 |
| 7 | 0.64 | 0.50 | 0.05 |
| 9 | −0.66 | 0.44 | 0.04 |
| 5 | 1.72 | 0.39 | 0.03 |
| 10 | 0.72 | 0.33 | 0.02 |
| 3 | −2.43 | 0.25 | 0.01 |

Don't say CAT

└ Short form procedures

└ Procedures based on θ targets

1 Introduction

2 Item Response Theory and information functions

3 IRT procedures for shortening tests

Benchmark procedure

Procedures based on θ targets

4 Simulation study

5 Some final remarks

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| | θ'_1 | θ'_2 | θ'_3 |
|------|-------------|-------------|-------------|
| item | −2.67 | 0.01 | 2.67 |
| 1 | | | |
| 2 | | | |
| 3 | | | |
| 4 | | | |
| 5 | | | |
| 6 | | | |
| 7 | | | |
| 8 | | | |
| 9 | | | |
| 10 | | | |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| | θ'_1 | θ'_2 | θ'_3 |
|------|-------------|-------------|-------------|
| item | -2.67 | 0.01 | 2.67 |
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| item | θ'_1 -2.67 | θ'_2 0.01 | θ'_3 2.67 |
|------|----------------------|---------------------|---------------------|
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| item | θ'_1 -2.67 | θ'_2 0.01 | θ'_3 2.67 |
|------|----------------------|---------------------|---------------------|
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| item | θ'_1 -2.67 | θ'_2 0.01 | θ'_3 2.67 |
|------|----------------------|---------------------|---------------------|
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| item | θ'_1 -2.67 | θ'_2 0.01 | θ'_3 2.67 |
|------|----------------------|---------------------|---------------------|
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| item | θ'_1 -2.67 | θ'_2 0.01 | θ'_3 2.67 |
|------|----------------------|---------------------|---------------------|
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

θ -target procedures

Selected items \rightarrow items with highest $IIFs$ in respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

| item | θ'_1 -2.67 | θ'_2 0.01 | θ'_3 2.67 |
|------|----------------------|---------------------|---------------------|
| 1 | 0.04 | 0.12 | 0.08 |
| 2 | 0.09 | 0.33 | 0.03 |
| 3 | 0.01 | 0.01 | 0.02 |
| 4 | 0.73 | 0.06 | 0.01 |
| 5 | 0.04 | 0.03 | 0.02 |
| 6 | 0.01 | 0.06 | 0.59 |
| 7 | 0.05 | 0.06 | 0.03 |
| 8 | 0.01 | 0.04 | 0.69 |
| 9 | 0.03 | 0.05 | 0.04 |
| 10 | 0.02 | 0.03 | 0.02 |

Don't say CAT

- └ Short form procedures

- └ Procedures based on θ targets


Segmenting the latent trait

Don't say CAT

└ Short form procedures

└ Procedures based on θ targets

Segmenting the latent trait




Equal Intervals Procedure
Equal segmentation

Don't say CAT

└ Short form procedures

└ Procedures based on θ targets

Segmenting the latent trait



Equal Intervals Procedure
Equal segmentation

Unequal Intervals Procedure
Clustering

1 Introduction

2 Item Response Theory and information functions

3 IRT procedures for shortening tests

Benchmark procedure

Procedures based on θ targets

4 Simulation study

5 Some final remarks

Comparison between the item selection procedures:

- **Benchmark procedure (BP)**: The N items with the highest $IIFs$ are selected from the full-length test
- **Equal Intervals Procedure (EIP)**: The N items that maximize the information for each θ' obtained by dividing the latent trait into equal intervals are selected
- **Unequal Intervals Procedure (UIP)**: The N items that maximize the information for each θ' obtained by clustering the latent trait are selected
- **Random Procedure (RP)**: N items are randomly selected from the full-length tests

10, 30, 50, 70, 90-item short test forms from a 100-item full-length test

1000 respondents p

- ① Normal distribution
 $p \sim \mathcal{N}(0, 1)$
- ② Positive skewed distribution
 $p \sim \text{Beta}(1, 100)$ (linearly transformed
to obtain negative values)
- ③ Uniform distribution
 $p \sim \mathcal{U}(-3, 3)$

100 items j :

- $b \sim \mathcal{U}(-3, 3)$
- $a \sim \mathcal{U}(0.40, 2)$

An overall look




Figure 3: Overall Information of the short test forms

A closer look




Figure 4: TIF of the 10-item short test form

An even closer look




Figure 5: $\text{bias} = \theta - \hat{\theta}$ of the 10-item short test form

- ① Introduction
- ② Item Response Theory and information functions
- ③ IRT procedures for shortening tests
 - Benchmark procedure
 - Procedures based on θ targets
- ④ Simulation study
- ⑤ Some final remarks

Good!

There's no “one-fits-all” solution

The θ distribution is a key element

Good!

There's no “one-fits-all” solution

The θ distribution is a key element

..but work is still needed

Real life applications are missing

The CAT is missing

Thank you!

`ottavia.epifania@unipd.it`