

DON'T SAY CAT: NEW ITEM RESPONSE THEORY APPROACHES FOR DEVELOPING SHORT TEST FORMS

Ottavia M. Epifania^{1,2}, Pasquale Anselmi¹, Egidio Robusto¹

ottavia.epifania@unipd.it

¹University of Padova

²Catholic University of the Sacred Heart

September 30th 2022, Padova

XXX Annual Conference of the Italian Psychology Association (AIP)



① Introduction

② Item Response Theory and information functions

③ IRT procedures for shortening tests

④ Simulation study

⑤ Some final remarks

CAT



CAT



Computerized **A**daptive **T**esting

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., each respondent) → (**CAT**: Computerized Adaptive Testing)

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., each respondent) → (**CAT**: Computerized Adaptive Testing)

STATIC SHORT FORMS: Static tests equal for all respondents → the information is maximized across θ levels (i.e., across all respondents)

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., each respondent) → (**CAT**: Computerized Adaptive Testing)

Issue

Different short test forms for each respondent → Potential fairness issues in assessments for recruitment

STATIC SHORT FORMS: Static tests equal for all respondents → the information is maximized across θ levels (i.e., across all respondents)

Item Response Theory and short test forms

ADAPTIVE SHORT FORMS: *Ad-hoc* tests for each person → The information is maximized for each level of θ (i.e., each respondent) → (**CAT**: Computerized Adaptive Testing)

Issue

Different short test forms for each respondent → Potential fairness issues in assessments for recruitment

STATIC SHORT FORMS: Static tests equal for all respondents → the information is maximized across θ levels (i.e., across all respondents)

Issue


Not being tailored to any θ level of interest → Potentially more items are needed to cover a wide range of θ s

Aim

New IRT-based procedures for shortening tests

Aim

New IRT-based procedures for shortening tests



Equal for all respondents

Tailored to specific levels of
the latent trait

① Introduction

② Item Response Theory and information functions

③ IRT procedures for shortening tests

④ Simulation study

⑤ Some final remarks

Item Response Theory

2-PL

$$P(x_{ps} = 1 | \theta_p, b_s, a_s) = \frac{\exp[a_s(\theta_p - b_s)]}{1 + \exp[a_s(\theta_p - b_s)]} \quad (1)$$

where:

$P(x_{ps} = 1)$: Probability of a correct response to item s by respondent p

θ_p : Ability of respondent's p

b_s : Difficulty of item s

a_s : Discrimination of item s

Information functions

Item Information Function

$$IIF_s = a_s^2 [P(\theta)(1 - P(\theta))] \quad (2)$$

Information functions

Item Information Function

$$IIF_s = a_s^2 [P(\theta)(1 - P(\theta))] \quad (2)$$




Figure 1: $a = 0.20$, $a = 0.70$, $a = 1.90$,
 $b = 0$

Information functions

Item Information Function

$$IIF_s = a_s^2 [P(\theta)(1 - P(\theta))] \quad (2)$$

Test Information Function

$$TIF = \sum_{s=1}^S IIF_s \quad (3)$$




Figure 1: $a = 0.20$, $a = 0.70$, $a = 1.90$,
 $b = 0$

Information functions

Item Information Function

$$IIF_s = a_s^2 [P(\theta)(1 - P(\theta))] \quad (2)$$




Figure 1: $a = 0.20$, $a = 0.70$, $a = 1.90$,
 $b = 0$

Test Information Function

$$TIF = \sum_{s=1}^S IIF_s \quad (3)$$




Figure 2: $TIF = IIF_1 + IIF_2 + IIF_3$

① Introduction

② Item Response Theory and information functions

③ IRT procedures for shortening tests

④ Simulation study

⑤ Some final remarks

Benchmark procedure

Selected items → items with the highest *IIFs*

e.g.: 3-item short form from 10-item full-length test

Benchmark procedure

Selected items \rightarrow items with the highest $IIFs$

e.g.: 3-item short form from 10-item full-length test

item	b	a	IIF
1	-0.67	0.71	0.08
2	0.50	1.19	0.15
3	-2.43	0.25	0.01
4	2.12	1.98	0.24
5	1.72	0.39	0.03
6	-2.28	1.62	0.19
7	0.64	0.50	0.05
8	-2.51	1.68	0.19
9	-0.66	0.44	0.04
10	0.72	0.33	0.02

Benchmark procedure

Selected items → items with the highest *IIFs*

e.g.: 3-item short form from 10-item full-length test

item	<i>b</i>	<i>a</i>	<i>IIF</i>
4	2.12	1.98	0.24
8	−2.51	1.68	0.19
6	−2.28	1.62	0.19
2	0.50	1.19	0.15
1	−0.67	0.71	0.08
7	0.64	0.50	0.05
9	−0.66	0.44	0.04
5	1.72	0.39	0.03
10	0.72	0.33	0.02
3	−2.43	0.25	0.01

Benchmark procedure

Selected items → items with the highest *IIFs*

e.g.: 3-item short form from 10-item full-length test

item	<i>b</i>	<i>a</i>	<i>IIF</i>
4	2.12	1.98	0.24
8	−2.51	1.68	0.19
6	−2.28	1.62	0.19
2	0.50	1.19	0.15
1	−0.67	0.71	0.08
7	0.64	0.50	0.05
9	−0.66	0.44	0.04
5	1.72	0.39	0.03
10	0.72	0.33	0.02
3	−2.43	0.25	0.01

Benchmark procedure

Selected items → items with the highest *IIFs*

e.g.: 3-item short form from 10-item full-length test

item	<i>b</i>	<i>a</i>	<i>IIF</i>
4	2.12	1.98	0.24
8	−2.51	1.68	0.19
6	−2.28	1.62	0.19
2	0.50	1.19	0.15
1	−0.67	0.71	0.08
7	0.64	0.50	0.05
9	−0.66	0.44	0.04
5	1.72	0.39	0.03
10	0.72	0.33	0.02
3	−2.43	0.25	0.01

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test

	θ'_1	θ'_2	θ'_3
item	-2.67	0.01	2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test


item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Procedures based on θ targets

Selected items \rightarrow items with highest *IIFs* in respect to θ targets (θ')
e.g.: 3-item short form from 10-item full-length test


item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

Segmenting the latent trait



Equal Intervals Procedure
Equal segmentation

Segmenting the latent trait



Equal Intervals Procedure
Equal segmentation

Unequal Intervals Procedure
Clustering

① Introduction

② Item Response Theory and information functions

③ IRT procedures for shortening tests

④ Simulation study

⑤ Some final remarks

Comparison between the item selection procedures:

- **Benchmark procedure (BP)**: The N items with the highest $IIFs$ are selected from the full-length test
- **Unequal Intervals Procedure (UIP)**: The N items that maximize the information for each θ' obtained by clustering the latent trait are selected
- **Equal Intervals Procedure (EIP)**: The N items that maximize the information for each θ' obtained by dividing the latent trait into equal intervals are selected
- **Random Procedure (RP)**: N items are randomly selected from the full-length tests

10, 30, 50, 70-item short test forms from a 100-item full-length test

1000 respondents p

- ① Normal distribution
 $p \sim \mathcal{N}(0, 1)$
- ② Positive skewed distribution
 $p \sim \text{Beta}(1, 100)$ (linearly transformed to obtain negative values)
- ③ Uniform distribution
 $p \sim \mathcal{U}(-3, 3)$

100 items s :

- $b \sim \mathcal{U}(-3, 3)$
- $a \sim \mathcal{U}(0.40, 2)$

An overall look




Figure 3: Overall Information of the short test forms

A closer look




Figure 4: TIF of the 10-item short test form

An even closer look




Figure 5: $\text{bias} = \theta - \hat{\theta}$ of the 10-item short test form

① Introduction

② Item Response Theory and information functions

③ IRT procedures for shortening tests

④ Simulation study

⑤ Some final remarks

Good!

There's no "one-fits-all" solution

The θ distribution is a key element

Good!

There's no "one-fits-all" solution

The θ distribution is a key element

..but work is still needed

Real life applications are missing

Thank you!

`ottavia.epifania@unipd.it`