

Research activity, interests and beyond

Ottavia M. Epifania, Ph.D.



Università di Trento
Rovereto, Italia

May, 22nd 2024

Item Response Theory and Rasch modeling

```
( TITLE-ABS-KEY ( "Item response theory" ) OR TITLE-ABS-KEY ( "Rasch" ) )  
AND PUBYEAR > 1959 AND PUBYEAR < 2025 AND ( LIMIT-TO ( DOCTYPE , "ar" ) )
```

$n = 18,661$ documents (May, 17th 2024) on Scopus:

Some fields of application:



Some reasons for their application:

Measure validation

Measure refinement

Item selection

Computerized adaptive testing


...

Rasch revised & beyond
oooooooo

IRT and short test forms
ooooo

Measurement theory
oooo

Teaching Activity
oo



Rasch modeling of complex data structures



A_{Bart}



A_{Lisa}



A_{Bart}

Q1

$$4 + 5 = ?$$

d_{q1}

Q2

$$\frac{3}{2}x^2 + \frac{5}{4}x = ?$$

d_{q2}



A_{Lisa}



A_{Bart}

Q1

$$4 + 5 = ?$$

d_{q1}

Q2

$$\frac{3}{2}x^2 + \frac{5}{4}x = ?$$

d_{q2}




A_{Lisa}


$$P(X_{pi} = 1) = \frac{\frac{A_p}{d_i}}{1 + \frac{A_p}{d_i}}$$

$$P(X_{pi} = 1 | \theta_p, b_i) = \frac{\exp(\theta_p - b_i)}{1 + \exp(\theta_p - b_i)}$$


$$P(X_{pi} = 1|\theta_p, b_i) = \frac{\exp(\theta_p - b_i)}{1 + \exp(\theta_p - b_i)}$$




$$P(X_{pi} = 1|\theta_p, b_i) = \frac{\exp(\theta_p - b_i)}{1 + \exp(\theta_p - b_i)}$$




$$P(X_{pi} = 1|\theta_p, b_i) = \frac{\exp(\theta_p - b_i)}{1 + \exp(\theta_p - b_i)}$$






Generalized linear model (GLM) for dichotomous responses



Generalized linear model (GLM) for dichotomous responses



Random effects and random factors

Linear combination of predictors in a Linear Model:

$$\eta = X\beta,$$

where β indicates the coefficients of the fixed intercept and slope(s), and X is the model-matrix.

Linear combination of predictors in a Linear Mixed-Effects Model (LMM):

$$\eta = X\beta + Zd,$$

where Z is the matrix and d is the vector of the random effects (not parameters!)

Best Linear Unbiased Predictors

Future lines and developments

Application to data from experiments typically used in different fields of study, such:


- Go/No-Go in addiction studies
- SNARC effect in cognitive psychology
- ...

Integration between the information obtained from the accuracy responses and the response times

Item Response Theory and short test forms development


2-Parameter Logistic Model

Item Response Function: $P(x_{pi} = 1|\theta_p, b_i, a_i) = \frac{\exp[a_i(\theta_p - b_i)]}{1 + \exp[a_i(\theta_p - b_i)]}$




2-Parameter Logistic Model

Item Information Function: $I_i(\theta) = a_i^2 P_i(\theta, b_i, a_i)[1 - P_i(\theta, b_i, a_i)]$



2-Parameter Logistic Model

Test Information Function: $I(\theta) = \sum_{i=1}^I I_i(\theta)$



θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

	θ'_1	θ'_2	θ'_3
item	-2.67	0.01	2.67
1			
2			
3			
4			
5			
6			
7			
8			
9			
10			

θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

	θ'_1	θ'_2	θ'_3
item	-2.67	0.01	2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02

θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')

e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02


θ -target procedure

Selected items \rightarrow items with highest *IIFs* with respect to θ targets (θ')


e.g.: 3-item short form from 10-item full-length test

item	θ'_1 -2.67	θ'_2 0.01	θ'_3 2.67
1	0.04	0.12	0.08
2	0.09	0.33	0.03
3	0.01	0.01	0.02
4	0.73	0.06	0.01
5	0.04	0.03	0.02
6	0.01	0.06	0.59
7	0.05	0.06	0.03
8	0.01	0.04	0.69
9	0.03	0.05	0.04
10	0.02	0.03	0.02


The other way around




The other way around



The other way around



The other way around



Future lines and directions


- Development of short test forms for the diagnosis of clinically relevant constructs (e.g., addiction) → Focused on specific regions of the latent trait
- Extension to multidimensional/bifactor latent structures (e.g., content balancing) and to polytomous data

Measurement theory


Measuring fluid intelligence

Raven-like measure for the assessment of fluid intelligence in children and adults:
From stimulus generation to measure validation under three different approaches

Classical Test Theory
Factorial solution



Rasch modeling
Wright map



The meaningfulness of Psychological Measures



The ratio between the measures of a and b is constant and independent of the measurement unit:



$$\frac{\varphi(a)}{\varphi(b)} = \frac{\varphi'(a)}{\varphi'(b)},$$

where φ and φ' are two different scales of measurement of the same variable.



Meaningful comparisons

The comparison between a and b is meaningful if it is invariant under all the unit transformations.







Item	t_A	t_B	φ_A	φ_B	φ'_A	φ'_B
1	14	17	4	3	4	4
2	26	19	2	3	3	4
3	31	18	1	3	2	4
4	47	33	0	1	1	2
5	17	47	3	0	4	1
Sum scores			10	10	14	15



Item	t_A	t_B	φ_A	φ_B	φ'_A	φ'_B
1	14	17	4	3	4	4
2	26	19	2	3	3	4
3	31	18	1	3	2	4
4	47	33	0	1	1	2
5	17	47	3	0	4	1
Sum scores			10	10	14	15



Item	t_A	t_B	φ_A	φ_B	φ'_A	φ'_B
1	14	17	4	3	4	4
2	26	19	2	3	3	4
3	31	18	1	3	2	4
4	47	33	0	1	1	2
5	17	47	3	0	4	1
Sum scores			10	10	14	15

Teaching Activity

From theory...

Psychometrics

Statistics, data analysis, and experiment design in Social Sciences

Measurement Theory and Measurement models in Psychology

Rasch modeling and Item Response Theory

...to practice

R for beginneRs

RMarkdown in an Open Science perspective

Developing web applications with shiny

Implementing experiments in Inquisit

Thank you!

