



TECHNISCHE  
UNIVERSITÄT  
WIEN

## **165.144 Simulation of condensed matter**

2025   Ass.Prof. Esther Heid   [esther.heid@tuwien.ac.at](mailto:esther.heid@tuwien.ac.at)

# Topics covered

## Properties

- Semester hours: 2.0
- Credits: 3.0
- Type: VU Lecture and Exercise
- Format: Presence

## Learning outcomes

After successful completion of the course, students are able to...

- Understand the methodology, approximations, advantages and disadvantages of molecular dynamics
- Use and operate molecular dynamics simulations for soft matter
- Design simple classical and machine-learning based force fields
- Analyze trajectories obtained from molecular dynamics simulations to extract physical/chemical properties

## Subject of course

Molecular dynamics methods (force fields, thermostats, and involved approximations). Molecular dynamics simulations of liquids and analysis of their properties (d force fields obtained from machine learning.

## Teaching methods

- Lectures about molecular dynamics
- Practical exercises on a PC

# About the course


**Mode:** In-person (but no attendance enforced), recordings

**Grading:**

- Oral exam at the end of the semester
- Exercise assignments (JupyterHub)
- Voluntary TUWEL quizzes for bonus points

**Prerequisites:** Theoretical Chemistry (recommended, but not mandatory)

**Language:** English



## **Chapter 1: Classical mechanics in the Newtonian formulation**

**Goal:** Study the movements of atoms in a condensed system, here liquids, and relate them to a macroscopic property.

## Theory (Statistical Mechanics)

---

Averages via phase-space integrals

$$\langle G \rangle = \frac{\int G(r_1 \dots p_N) e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}{\int e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}$$

$G$  ... any quantity

$$\beta = \frac{1}{k_B T}$$

$k_B$  ... Boltzmann constant

→ only expectation value

**Goal:** Study the movements of atoms in a condensed system, here liquids, and relate them to a macroscopic property.

### Theory (Statistical Mechanics)

### Simulation (Monte Carlo)

---

Averages via phase-space integrals

$$\langle G \rangle = \frac{\int G(r_1 \dots p_N) e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}{\int e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}$$

$G$  ... any quantity

$$\beta = \frac{1}{k_B T}$$

$k_B$  ... Boltzmann constant

→ only expectation value

Sample over states

$$\langle G \rangle = \frac{1}{S} \sum_{s=1}^S G(s)$$

where states are chosen

with probability  $e^{-\beta E_{\text{pot}}(s)}$

→ only configuration space,  
no dynamics

**Goal:** Study the movements of atoms in a condensed system, here liquids, and relate them to a macroscopic property.

### Theory (Statistical Mechanics)

Averages via phase-space integrals

$$\langle G \rangle = \frac{\int G(r_1 \dots p_N) e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}{\int e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}$$

$G$  ... any quantity

$$\beta = \frac{1}{k_B T}$$

$k_B$  ... Boltzmann constant

→ only expectation value

### Simulation (Monte Carlo)

Sample over states

$$\langle G \rangle = \frac{1}{S} \sum_{s=1}^S G(s)$$

where states are chosen

with probability  $e^{-\beta E_{\text{pot}}(s)}$

→ only configuration space,  
no dynamics

### Simulation (Molecular dynamics)

Follow the dynamics of a system

$$\langle G \rangle = \frac{1}{M} \sum_{\mu=1}^M G_{\mu}$$

$\mu$  ... measurement

→ average over measurements  $\mu$   
during dynamics

**Goal:** Study the movements of atoms in a condensed system, here liquids, and relate them to a macroscopic property.

### Theory (Statistical Mechanics)

Averages via phase-space integrals

$$\langle G \rangle = \frac{\int G(r_1 \dots p_N) e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}{\int e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}$$

$G$  ... any quantity

$$\beta = \frac{1}{k_B T}$$

$k_B$  ... Boltzmann constant

→ only expectation value

### Simulation (Monte Carlo)

Sample over states

$$\langle G \rangle = \frac{1}{S} \sum_{s=1}^S G(s)$$

where states are chosen

with probability  $e^{-\beta E_{\text{pot}}(s)}$

→ only configuration space,  
no dynamics

### Simulation (Molecular dynamics)

Follow the dynamics of a system

$$\langle G \rangle = \frac{1}{M} \sum_{\mu=1}^M G_{\mu}$$

$\mu$  ... measurement

→ average over measurements  $\mu$   
during dynamics

**Example: Oscillator**  $\frac{k}{2}(x - x_0)^2$

$$\langle x \rangle = \frac{\int \dots}{\int \dots} = x_0$$



**Goal:** Study the movements of atoms in a condensed system, here liquids, and relate them to a macroscopic property.

### Theory (Statistical Mechanics)

Averages via phase-space integrals

$$\langle G \rangle = \frac{\int G(r_1 \dots p_N) e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}{\int e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}$$

$G$  ... any quantity

$$\beta = \frac{1}{k_B T}$$

$k_B$  ... Boltzmann constant

→ only expectation value

### Simulation (Monte Carlo)

Sample over states

$$\langle G \rangle = \frac{1}{S} \sum_{s=1}^S G(s)$$

where states are chosen

with probability  $e^{-\beta E_{\text{pot}}(s)}$

→ only configuration space,  
no dynamics

### Simulation (Molecular dynamics)

Follow the dynamics of a system


$$\langle G \rangle = \frac{1}{M} \sum_{\mu=1}^M G_{\mu}$$

$\mu$  ... measurement

→ average over measurements  $\mu$   
during dynamics

**Example: Oscillator**  $\frac{k}{2}(x - x_0)^2$

$$\langle x \rangle = \frac{\int \dots}{\int \dots} = x_0$$



**Goal:** Study the movements of atoms in a condensed system, here liquids, and relate them to a macroscopic property.

## Theory (Statistical Mechanics)

Averages via phase-space integrals

$$\langle G \rangle = \frac{\int G(r_1 \dots p_N) e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}{\int e^{-\beta E_{\text{pot}}(r_1 \dots p_N)} dr_1 \dots dp_N}$$

$G$  ... any quantity

$$\beta = \frac{1}{k_B T}$$

$k_B$  ... Boltzmann constant

→ only expectation value

## Simulation (Monte Carlo)

Sample over states

$$\langle G \rangle = \frac{1}{S} \sum_{s=1}^S G(s)$$

where states are chosen

with probability  $e^{-\beta E_{\text{pot}}(s)}$

→ only configuration space,  
no dynamics

## Simulation (Molecular dynamics)

Follow the dynamics of a system



$$\langle G \rangle = \frac{1}{M} \sum_{\mu=1}^M G_{\mu}$$

$\mu$  ... measurement

→ average over measurements  $\mu$   
during dynamics

**Example: Oscillator**  $\frac{k}{2}(x - x_0)^2$

$$\langle x \rangle = \frac{\int \dots}{\int \dots} = x_0$$




## For an MD simulation, we need to know:

- ① How atoms interact, i.e. given coordinates  $\vec{r}$ , obtain forces  $\vec{f}(\vec{r})$
- ② How atoms move, i.e. given forces  $\vec{f}$ , obtain coordinates at next time step  $\vec{r}(t + \Delta t)$

## For an MD simulation, we need to know:

- ① How atoms interact, i.e. given coordinates  $\vec{r}$ , obtain forces  $\vec{f}(\vec{r})$
- ② How atoms move, i.e. given forces  $\vec{f}$ , obtain coordinates at next time step  $\vec{r}(t + \Delta t)$


Problem 2 is old: N-body problem to study celestial objects, e.g. sun, earth and jupyter. No analytic solution for  $N \geq 3$



## For an MD simulation, we need to know:


- ① How atoms interact, i.e. given coordinates  $\vec{r}$ , obtain forces  $\vec{f}(\vec{r})$
- ② How atoms move, i.e. given forces  $\vec{f}$ , obtain coordinates at next time step  $\vec{r}(t + \Delta t)$

Problem 2 is old: N-body problem to study celestial objects, e.g. sun, earth and jupyter. No analytic solution for  $N \geq 3$



For studying a liquid, we need to follow the motions of thousands of atoms, so it is only possible to solve this problem numerically. For this, we only need Newton's classical mechanics

**Classical mechanics:** Physical theory describing the motion of macroscopic objects (disregarding quantum mechanics + relativity)




! The laws of classical mechanics can be derived from the asymptote of quantum mechanics (QM) with increasingly large systems

**Newtonian mechanics:** Earliest and simplest formulation from the 17th century which is directly applicable to the computer simulation of matter

## 1.1 Single point particle


**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...



# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...

Position  $\mathbf{x}(t)$  ... vector of cartesian coordinates






# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...

Position  $\mathbf{x}(t)$  ... vector of cartesian coordinates

Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$




# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...

Position  $\mathbf{x}(t)$  ... vector of cartesian coordinates

Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$

Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$  (=0 only for free particles)



# 1.1 Single point particle


**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...

Position  $\mathbf{x}(t)$  ... vector of cartesian coordinates

Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$

Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$  (=0 only for free particles)

Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$



# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


Position  $\mathbf{x}(t)$  ... vector of cartesian coordinates

Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$

Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$  (=0 only for free particles)

Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$

Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$  = Newton's second law



# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x}$$

# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt$$

# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


Position  $\mathbf{x}(t)$  ... vector of cartesian coordinates

Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$

Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$  (=0 only for free particles)

Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$

Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt = \int_{t_A}^{t_B} \mathbf{f} \cdot \mathbf{v} dt$$

# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt = \int_{t_A}^{t_B} \mathbf{f} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \frac{d\mathbf{v}}{dt} \cdot \mathbf{v} dt$$



# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt = \int_{t_A}^{t_B} \mathbf{f} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \frac{d\mathbf{v}}{dt} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \mathbf{v} d\mathbf{v}$$

# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$\begin{aligned} W_{A \rightarrow B}[C] &= \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt = \int_{t_A}^{t_B} \mathbf{f} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \frac{d\mathbf{v}}{dt} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \mathbf{v} d\mathbf{v} \\ &= \frac{m|\mathbf{v}|^2}{2} \Big|_{t_A}^{t_B} \end{aligned}$$

# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$\begin{aligned} W_{A \rightarrow B}[C] &= \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt = \int_{t_A}^{t_B} \mathbf{f} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \frac{d\mathbf{v}}{dt} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \mathbf{v} d\mathbf{v} \\ &= \frac{m|\mathbf{v}|^2}{2} \Big|_{t_A}^{t_B} = \frac{m}{2} [|\mathbf{v}(t_B)|^2 - |\mathbf{v}(t_A)|^2] \end{aligned}$$

# 1.1 Single point particle

**Point particle:** Particle is represented using position + mass only, disregarding its structure.  
Can be a ball, apple, chicken, planet, ...


**Position  $\mathbf{x}(t)$**  ... vector of cartesian coordinates

**Velocity  $\mathbf{v}(t) = \frac{d\mathbf{x}}{dt}$**

**Acceleration  $\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$**  (=0 only for free particles)

**Jerk  $\mathbf{j}(t) = \frac{d\mathbf{a}}{dt} = \frac{d^2\mathbf{v}}{dt^2} = \frac{d^3\mathbf{x}}{dt^3}$**

**Force  $\mathbf{f}(t) = m \frac{d^2\mathbf{x}}{dt^2} = m\mathbf{a}$**  = Newton's second law



**Work  $W$**  ... if we want to move a particle from a point  $A$  to a point  $B$  along a trajectory, we need to apply a force:

$$\begin{aligned} W_{A \rightarrow B}[C] &= \int_C \mathbf{f} d\mathbf{x} = \int_{t_A}^{t_B} \mathbf{f} \cdot \frac{d\mathbf{x}}{dt} dt = \int_{t_A}^{t_B} \mathbf{f} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \frac{d\mathbf{v}}{dt} \cdot \mathbf{v} dt = m \int_{t_A}^{t_B} \mathbf{v} d\mathbf{v} \\ &= \frac{m|\mathbf{v}|^2}{2} \Big|_{t_A}^{t_B} = \frac{m}{2} [|\mathbf{v}(t_B)|^2 - |\mathbf{v}(t_A)|^2] = E_k(t_B) - E_k(t_A) \text{ with kinetic energy } E_k = \frac{1}{2} m |\mathbf{v}|^2 \end{aligned}$$

→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0 \dots$  conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0$  ... conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

Going from  $A$  to  $B$  via curve  $C_1$  and back via  $C_2$ :

$$0 = \int_{C_1} \mathbf{f} d\mathbf{x} + \int_{C_2} \mathbf{f} d\mathbf{x} =$$

→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0$  ... conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

Going from  $A$  to  $B$  via curve  $C_1$  and back via  $C_2$ :

$$0 = \int_{C_1} \mathbf{f} d\mathbf{x} + \int_{C_2} \mathbf{f} d\mathbf{x} = W_{A \rightarrow B}[C_1] + W_{B \rightarrow A}[C_2] =$$

→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0 \dots$  conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

Going from  $A$  to  $B$  via curve  $C_1$  and back via  $C_2$ :

$$0 = \int_{C_1} \mathbf{f} d\mathbf{x} + \int_{C_2} \mathbf{f} d\mathbf{x} = W_{A \rightarrow B}[C_1] + W_{B \rightarrow A}[C_2] = W_{A \rightarrow B}[C_1] - W_{A \rightarrow B}[C_2] = 0$$



→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0$  ... conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

Going from  $A$  to  $B$  via curve  $C_1$  and back via  $C_2$ :

$$0 = \int_{C_1} \mathbf{f} d\mathbf{x} + \int_{C_2} \mathbf{f} d\mathbf{x} = W_{A \rightarrow B}[C_1] + W_{B \rightarrow A}[C_2] = W_{A \rightarrow B}[C_1] - W_{A \rightarrow B}[C_2] = 0$$

→ Work is trajectory-independent, i.e.  $W_{A \rightarrow B}[C_1] = W_{A \rightarrow B}[C_2]$

→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0$  ... conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

Going from  $A$  to  $B$  via curve  $C_1$  and back via  $C_2$ :

$$0 = \int_{C_1} \mathbf{f} d\mathbf{x} + \int_{C_2} \mathbf{f} d\mathbf{x} = W_{A \rightarrow B}[C_1] + W_{B \rightarrow A}[C_2] = W_{A \rightarrow B}[C_1] - W_{A \rightarrow B}[C_2] = 0$$

→ Work is trajectory-independent, i.e.  $W_{A \rightarrow B}[C_1] = W_{A \rightarrow B}[C_2]$

We can therefore choose an arbitrary point  $P$  as origin of energies:

$$W_{\mathbf{x} \rightarrow P} = -W_{P \rightarrow \mathbf{x}} = E_{\text{pot}}(\mathbf{x}) \dots \text{potential energy}$$

→ work does not depend on the path

→ for a closed curve, i.e.  $B = A$ :  $W_{A \rightarrow B}[C] = \int_C \mathbf{f} d\mathbf{x} = 0 \dots$  conservative force field

→ this is only true for a conservative force field, where  $\mathbf{f}$  must be defined over a simply connected region without holes

Going from  $A$  to  $B$  via curve  $C_1$  and back via  $C_2$ :

$$0 = \int_{C_1} \mathbf{f} d\mathbf{x} + \int_{C_2} \mathbf{f} d\mathbf{x} = W_{A \rightarrow B}[C_1] + W_{B \rightarrow A}[C_2] = W_{A \rightarrow B}[C_1] - W_{A \rightarrow B}[C_2] = 0$$

→ Work is trajectory-independent, i.e.  $W_{A \rightarrow B}[C_1] = W_{A \rightarrow B}[C_2]$

We can therefore choose an arbitrary point  $P$  as origin of energies:

$$W_{\mathbf{x} \rightarrow P} = -W_{P \rightarrow \mathbf{x}} = E_{\text{pot}}(\mathbf{x}) \dots \text{potential energy}$$

Since  $W_{A \rightarrow B} = \Delta_{A \rightarrow B} E_k$  and  $W_{A \rightarrow B} = -\Delta_{A \rightarrow B} E_{\text{pot}}$ , we can derive the conservation of energy

$$0 = \Delta_{A \rightarrow B}(E_k + E_{\text{pot}}) = \Delta_{A \rightarrow B} E \text{ with } E = E_k + E_{\text{pot}} \dots \text{mechanical energy}$$

Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z}$$

Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z}$$

Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta \mathbf{z} \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta \mathbf{z} \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z}$$

Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} =$$
$$\lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z}$$

Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} =$$
$$\lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow \mathbf{x}_0}}{\Delta z}$$



Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} =$$

$$\lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z}}^{\mathbf{x}_0} \mathbf{f} d\mathbf{x}}{\Delta z}$$

Change of energy if we move a particle upwards:

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} =$$

$$\lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z}}^{\mathbf{x}_0} \mathbf{f} d\mathbf{x}}{\Delta z}$$

Define  $\mathbf{x}(s) = \mathbf{x}_0 + s \cdot \mathbf{z}$  with  $s \in [0, \Delta z]$  :

Change of energy if we move a particle upwards:

$$\begin{aligned}\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} &= \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} = \\ \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z} &= \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z}}^{\mathbf{x}_0} \mathbf{f} d\mathbf{x}}{\Delta z}\end{aligned}$$

Define  $\mathbf{x}(s) = \mathbf{x}_0 + s \cdot \mathbf{z}$  with  $s \in [0, \Delta z]$  :

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\Delta z}^0 \mathbf{f} \cdot \mathbf{z} ds}{\Delta z} = -\lim_{\Delta z \rightarrow 0} \frac{f_z \Delta z}{\Delta z} = -f_z$$

Change of energy if we move a particle upwards:

$$\begin{aligned}\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} &= \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} = \\ \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z} &= \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z}}^{\mathbf{x}_0} \mathbf{f} d\mathbf{x}}{\Delta z}\end{aligned}$$

Define  $\mathbf{x}(s) = \mathbf{x}_0 + s \cdot \mathbf{z}$  with  $s \in [0, \Delta z]$  :

$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\Delta z}^0 \mathbf{f} \cdot \mathbf{z} ds}{\Delta z} = -\lim_{\Delta z \rightarrow 0} \frac{f_z \Delta z}{\Delta z} = -f_z$$

With the same calculation for  $x$  and  $y$ , we find that  $\mathbf{f} = -\nabla E_{\text{pot}}$

Change of energy if we move a particle upwards:

$$\begin{aligned}\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} &= \lim_{\Delta z \rightarrow 0} \frac{E_{\text{pot}}(\mathbf{x}_0 + \Delta z \cdot \mathbf{z}) - E_{\text{pot}}(\mathbf{x}_0)}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} - W_{\mathbf{x}_0 \rightarrow P}}{\Delta z} = \\ \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow P} + W_{P \rightarrow \mathbf{x}_0}}{\Delta z} &= \lim_{\Delta z \rightarrow 0} \frac{W_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z} \rightarrow \mathbf{x}_0}}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\mathbf{x}_0 + \Delta z \cdot \mathbf{z}}^{\mathbf{x}_0} \mathbf{f} d\mathbf{x}}{\Delta z}\end{aligned}$$

Define  $\mathbf{x}(s) = \mathbf{x}_0 + s \cdot \mathbf{z}$  with  $s \in [0, \Delta z]$  :


$$\frac{\partial E_{\text{pot}}(\mathbf{x}_0)}{\partial z} = \lim_{\Delta z \rightarrow 0} \frac{\int_{\Delta z}^0 \mathbf{f} \cdot \mathbf{z} ds}{\Delta z} = -\lim_{\Delta z \rightarrow 0} \frac{f_z \Delta z}{\Delta z} = -f_z$$

With the same calculation for  $x$  and  $y$ , we find that  $\mathbf{f} = -\nabla E_{\text{pot}}$

### In summary:

- Work is related to the kinetic energy:  $W_{A \rightarrow B}[C] = E_k(t_B) - E_k(t_A)$
- Work is related to the potential energy if the forces are conservative:  
 $W_{\mathbf{x} \rightarrow P} = -W_{P \rightarrow \mathbf{x}} = E_{\text{pot}}(\mathbf{x})$
- Mechanical energy is conserved:  $0 = \Delta_{A \rightarrow B}(E_k + E_{\text{pot}}) = \Delta_{A \rightarrow B} E$
- Force is the negative gradient of the potential energy:  $\mathbf{f} = -\nabla E_{\text{pot}}$

## 1.2 System of point particles




For each particle:

$$\mathbf{v}_i(t) = \frac{d\mathbf{x}_i}{dt}$$

$$\mathbf{a}_i(t) = \frac{d^2\mathbf{x}_i}{dt^2}$$

$$\mathbf{f}_i(t) = m_i \frac{d^2\mathbf{x}_i}{dt^2}$$

## 1.2 System of point particles



For each particle:

$$\mathbf{v}_i(t) = \frac{d\mathbf{x}_i}{dt}$$

$$\mathbf{a}_i(t) = \frac{d^2\mathbf{x}_i}{dt^2}$$

$$\mathbf{f}_i(t) = m_i \frac{d^2\mathbf{x}_i}{dt^2}$$


For all particles:

Instead of several  $\mathbf{x}_i$ , make one big vector

$$\mathbf{x}(t) = (x_1, y_1, z_1, \dots, x_N, y_N, z_N)$$

$$\mathbf{f}(t) = (f_1^x, f_1^y, f_1^z, \dots, f_N^x, f_N^y, f_N^z)$$

## 1.2 System of point particles



For each particle:

$$\mathbf{v}_i(t) = \frac{d\mathbf{x}_i}{dt}$$

$$\mathbf{a}_i(t) = \frac{d^2\mathbf{x}_i}{dt^2}$$

$$\mathbf{f}_i(t) = m_i \frac{d^2\mathbf{x}_i}{dt^2}$$

For all particles:

Instead of several  $\mathbf{x}_i$ , make one big vector

$$\mathbf{x}(t) = (x_1, y_1, z_1, \dots, x_N, y_N, z_N)$$


$$\mathbf{f}(t) = (f_1^x, f_1^y, f_1^z, \dots, f_N^x, f_N^y, f_N^z)$$

Work:  $W_{A \rightarrow B}[C] = E_k(t_B) - E_k(t_A) = \int_C \mathbf{f} d\mathbf{x} = \Delta E_k$

with  $E_k = \frac{1}{2} \sum m_i |\mathbf{v}_i|^2$  ... kinetic energy of all particles



## 1.2 System of point particles



For each particle:

$$\mathbf{v}_i(t) = \frac{d\mathbf{x}_i}{dt}$$

$$\mathbf{a}_i(t) = \frac{d^2\mathbf{x}_i}{dt^2}$$

$$\mathbf{f}_i(t) = m_i \frac{d^2\mathbf{x}_i}{dt^2}$$

For all particles:

Instead of several  $\mathbf{x}_i$ , make one big vector

$$\mathbf{x}(t) = (x_1, y_1, z_1, \dots, x_N, y_N, z_N)$$

$$\mathbf{f}(t) = (f_1^x, f_1^y, f_1^z, \dots, f_N^x, f_N^y, f_N^z)$$

Work:  $W_{A \rightarrow B}[C] = E_k(t_B) - E_k(t_A) = \int_C \mathbf{f} d\mathbf{x} = \Delta E_k$

with  $E_k = \frac{1}{2} \sum m_i |\mathbf{v}_i|^2$  ... kinetic energy of all particles


For conservative forcefields:

- $W_{A \rightarrow B} = -\Delta_{A \rightarrow B} E_{\text{pot}}$
- $\mathbf{f} = -\nabla E_{\text{pot}}$  and  $\mathbf{f}_i = -\frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i}$
- $E = E_k + E_{\text{pot}}$

→ same as for single particle, only in 3N-dimensional space

# 1.3 Fundamental symmetries of mechanics

## Translational symmetry



Mechanics observed from  
 $\Sigma$  and  $\Sigma'$  must  
be the same!

Initial conditions:  $\vec{x}_0, \vec{v}_0$  trivially consistent, i.e. offset in  $\vec{x}_0$

Newton's equation of motion, i.e.  $\vec{f}$ :  $\vec{f}$  must transform as a vector for translations  
 $E_{\text{pot}}$  must transform as a scalar, i.e. must be the same for  
 $\Sigma$  and  $\Sigma'$  since  $\vec{f} = -\nabla E_{\text{pot}}$

$$E_{\text{pot}} = E_{\text{pot}}(x_1, y_1, z_1, \dots, x_N, y_N, z_N) = E_{\text{pot}}(\{x_i^\alpha\})$$

where  $i = 1, 2, \dots, N$  (atomic index) and  $\alpha = x, y, z$  (cartesian axis)

$$E_{\text{pot}} = E_{\text{pot}}(x_1, y_1, z_1, \dots, x_N, y_N, z_N) = E_{\text{pot}}(\{x_i^\alpha\})$$

where  $i = 1, 2, \dots, N$  (atomic index) and  $\alpha = x, y, z$  (cartesian axis)

System  $\Sigma$ :  $\{x_i^\alpha\} = \{C_i^\alpha\}$

System  $\Sigma'$ :  $\{x_i^\alpha\} = \{C_i^\alpha + \lambda u^\alpha\}$  with rigid displacement  $\lambda \mathbf{u}$

$$E_{\text{pot}} = E_{\text{pot}}(x_1, y_1, z_1, \dots, x_N, y_N, z_N) = E_{\text{pot}}(\{x_i^\alpha\})$$

where  $i = 1, 2, \dots, N$  (atomic index) and  $\alpha = x, y, z$  (cartesian axis)

System  $\Sigma$ :  $\{x_i^\alpha\} = \{C_i^\alpha\}$

System  $\Sigma'$ :  $\{x_i^\alpha\} = \{C_i^\alpha + \lambda u^\alpha\}$  with rigid displacement  $\lambda \mathbf{u}$

Translational invariance:  $E_{\text{pot}}(\{C_i^\alpha + \lambda u^\alpha\}) = E_{\text{pot}}(\{C_i^\alpha\}) \rightarrow \frac{\partial E_{\text{pot}}}{\partial \lambda} = 0$

$$E_{\text{pot}} = E_{\text{pot}}(x_1, y_1, z_1, \dots, x_N, y_N, z_N) = E_{\text{pot}}(\{x_i^\alpha\})$$

where  $i = 1, 2, \dots, N$  (atomic index) and  $\alpha = x, y, z$  (cartesian axis)

System  $\Sigma$ :  $\{x_i^\alpha\} = \{C_i^\alpha\}$

System  $\Sigma'$ :  $\{x_i^\alpha\} = \{C_i^\alpha + \lambda u^\alpha\}$  with rigid displacement  $\lambda \mathbf{u}$

Translational invariance:  $E_{\text{pot}}(\{C_i^\alpha + \lambda u^\alpha\}) = E_{\text{pot}}(\{C_i^\alpha\}) \rightarrow \frac{\partial E_{\text{pot}}}{\partial \lambda} = 0$

Using the chain rule for all arguments of  $E_{\text{pot}}$ :

$$\frac{\partial E_{\text{pot}}}{\partial \lambda} = \sum_{\alpha} \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} \frac{\partial x_i^\alpha}{\partial \lambda} = \sum_{\alpha} \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} u^\alpha = \sum_{\alpha} u^\alpha \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} = 0$  for all  $\alpha$

$$E_{\text{pot}} = E_{\text{pot}}(x_1, y_1, z_1, \dots, x_N, y_N, z_N) = E_{\text{pot}}(\{x_i^\alpha\})$$

where  $i = 1, 2, \dots, N$  (atomic index) and  $\alpha = x, y, z$  (cartesian axis)

System  $\Sigma$ :  $\{x_i^\alpha\} = \{C_i^\alpha\}$

System  $\Sigma'$ :  $\{x_i^\alpha\} = \{C_i^\alpha + \lambda u^\alpha\}$  with rigid displacement  $\lambda \mathbf{u}$

Translational invariance:  $E_{\text{pot}}(\{C_i^\alpha + \lambda u^\alpha\}) = E_{\text{pot}}(\{C_i^\alpha\}) \rightarrow \frac{\partial E_{\text{pot}}}{\partial \lambda} = 0$

Using the chain rule for all arguments of  $E_{\text{pot}}$ :

$$\frac{\partial E_{\text{pot}}}{\partial \lambda} = \sum_{\alpha} \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} \frac{\partial x_i^\alpha}{\partial \lambda} = \sum_{\alpha} \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} u^\alpha = \sum_{\alpha} u^\alpha \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} = 0$  for all  $\alpha$

$\rightarrow \sum_i f_i^\alpha = 0$  for all  $\alpha$ , thus  $\mathbf{f} = 0$  (=no net force on isolated system)

$$E_{\text{pot}} = E_{\text{pot}}(x_1, y_1, z_1, \dots, x_N, y_N, z_N) = E_{\text{pot}}(\{x_i^\alpha\})$$

where  $i = 1, 2, \dots, N$  (atomic index) and  $\alpha = x, y, z$  (cartesian axis)

System  $\Sigma$ :  $\{x_i^\alpha\} = \{C_i^\alpha\}$

System  $\Sigma'$ :  $\{x_i^\alpha\} = \{C_i^\alpha + \lambda u^\alpha\}$  with rigid displacement  $\lambda \mathbf{u}$

Translational invariance:  $E_{\text{pot}}(\{C_i^\alpha + \lambda u^\alpha\}) = E_{\text{pot}}(\{C_i^\alpha\}) \rightarrow \frac{\partial E_{\text{pot}}}{\partial \lambda} = 0$

Using the chain rule for all arguments of  $E_{\text{pot}}$ :

$$\frac{\partial E_{\text{pot}}}{\partial \lambda} = \sum_{\alpha} \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} \frac{\partial x_i^\alpha}{\partial \lambda} = \sum_{\alpha} \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} u^\alpha = \sum_{\alpha} u^\alpha \sum_i \frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{\partial E_{\text{pot}}}{\partial x_i^\alpha} = 0$  for all  $\alpha$


$\rightarrow \sum_i f_i^\alpha = 0$  for all  $\alpha$ , thus  $\mathbf{f} = 0$  (=no net force on isolated system)

Since  $f_i^\alpha = m \frac{dv_i^\alpha}{dt} = \frac{d}{dt} m_i v_i^\alpha = \frac{dp_i^\alpha}{dt} \rightarrow \frac{d}{dt} \sum_i p_i^\alpha = 0$  and  $\frac{d\mathbf{P}}{dt} = 0$

$\rightarrow$  conservation of linear momentum



# Rotational symmetry



Mechanics observed from  $\Sigma$  and  $\Sigma'$  must be the same

→  $\vec{f}$  must transform as vector with respect to rotation

→  $E_{\text{pot}}$  must transform as scalar (= not change)

$$E_{\text{pot}}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N) = E_{\text{pot}}(+(\vec{x}_1), +(\vec{x}_2), \dots, +(\vec{x}_N))$$

Rotation around  
axis  $\vec{u}$ , angle  $\theta$



using Rodrigues' rotation formula:

$$+(\vec{x}_i) = \underbrace{\vec{x}_i \cos \theta}_{\text{scales vector}} + \underbrace{(\vec{u} \times \vec{x}_i) \sin \theta}_{\text{skews vector toward new position}} + \underbrace{\vec{u} \cdot (\vec{u} \cdot \vec{x}_i) (1 - \cos \theta)}_{\text{re-adds the height that was lost in term 1}}$$

$$\frac{dE_{\text{pot}}}{d\theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} \frac{\partial \mathbf{x}_i}{\partial \theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{f}_i (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{u} (\mathbf{x}_i \times \mathbf{f}_i) = - \mathbf{u} [\sum_i m_i (\mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt})]$$

$$\frac{dE_{\text{pot}}}{d\theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} \frac{\partial \mathbf{x}_i}{\partial \theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{f}_i (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{u} (\mathbf{x}_i \times \mathbf{f}_i) = - \mathbf{u} [\sum_i m_i (\mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt})]$$

Since  $\frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i) = \frac{d\mathbf{x}_i}{dt} \times \mathbf{v}_i + \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt} = \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt}$ , we get:

$$\frac{dE_{\text{pot}}}{d\theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} \frac{\partial \mathbf{x}_i}{\partial \theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{f}_i (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{u} (\mathbf{x}_i \times \mathbf{f}_i) = -\mathbf{u} [\sum_i m_i (\mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt})]$$

Since  $\frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i) = \frac{d\mathbf{x}_i}{dt} \times \mathbf{v}_i + \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt} = \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt}$ , we get:

$$\frac{dE_{\text{pot}}}{d\theta} = -\mathbf{u} [\sum_i m_i \frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i)] = -\mathbf{u} \frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$  and thus  $\sum_i \mathbf{x}_i \times \mathbf{f}_i = 0$

$$\frac{dE_{\text{pot}}}{d\theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} \frac{\partial \mathbf{x}_i}{\partial \theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{f}_i (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{u} (\mathbf{x}_i \times \mathbf{f}_i) = - \mathbf{u} [\sum_i m_i (\mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt})]$$

Since  $\frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i) = \frac{d\mathbf{x}_i}{dt} \times \mathbf{v}_i + \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt} = \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt}$ , we get:

$$\frac{dE_{\text{pot}}}{d\theta} = - \mathbf{u} [\sum_i m_i \frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i)] = - \mathbf{u} \frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$  and thus  $\sum_i \mathbf{x}_i \times \mathbf{f}_i = 0$

→ Total torque  $\tau = \sum_i \mathbf{x}_i \times \mathbf{f}_i$  is zero (=no net torque on isolated system)

$$\frac{dE_{\text{pot}}}{d\theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} \frac{\partial \mathbf{x}_i}{\partial \theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{f}_i (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{u} (\mathbf{x}_i \times \mathbf{f}_i) = - \mathbf{u} [\sum_i m_i (\mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt})]$$

Since  $\frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i) = \frac{d\mathbf{x}_i}{dt} \times \mathbf{v}_i + \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt} = \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt}$ , we get:

$$\frac{dE_{\text{pot}}}{d\theta} = - \mathbf{u} [\sum_i m_i \frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i)] = - \mathbf{u} \frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$  and thus  $\sum_i \mathbf{x}_i \times \mathbf{f}_i = 0$

→ Total torque  $\tau = \sum_i \mathbf{x}_i \times \mathbf{f}_i$  is zero (=no net torque on isolated system)

Angular momentum  $\mathbf{l}_i = m\mathbf{x}_i \times \mathbf{v}_i = \mathbf{x}_i \times \mathbf{p}_i$  and total angular momentum  $\mathbf{L} = \sum_i \mathbf{l}_i$

$$\frac{dE_{\text{pot}}}{d\theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} \frac{\partial \mathbf{x}_i}{\partial \theta} = \sum_i \frac{\partial E_{\text{pot}}}{\partial \mathbf{x}_i} (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{f}_i (\mathbf{u} \times \mathbf{x}_i) = - \sum_i \mathbf{u} (\mathbf{x}_i \times \mathbf{f}_i) = - \mathbf{u} [\sum_i m_i (\mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt})]$$

Since  $\frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i) = \frac{d\mathbf{x}_i}{dt} \times \mathbf{v}_i + \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt} = \mathbf{x}_i \times \frac{d\mathbf{v}_i}{dt}$ , we get:

$$\frac{dE_{\text{pot}}}{d\theta} = - \mathbf{u} [\sum_i m_i \frac{d}{dt}(\mathbf{x}_i \times \mathbf{v}_i)] = - \mathbf{u} \frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$$

This can only be true for all  $\mathbf{u}$  if  $\frac{d}{dt} [\sum_i m_i \mathbf{x}_i \times \mathbf{v}_i] = 0$  and thus  $\sum_i \mathbf{x}_i \times \mathbf{f}_i = 0$

→ Total torque  $\tau = \sum_i \mathbf{x}_i \times \mathbf{f}_i$  is zero (=no net torque on isolated system)

Angular momentum  $\mathbf{l}_i = m\mathbf{x}_i \times \mathbf{v}_i = \mathbf{x}_i \times \mathbf{p}_i$  and total angular momentum  $\mathbf{L} = \sum_i \mathbf{l}_i$

→  $\frac{d}{dt} \sum_i \mathbf{l}_i = 0$  and  $\frac{dL}{dt} = 0$

→ conservation of angular momentum

## Summary:

- Translational symmetry = Conservation of linear momentum  
 $\frac{d\mathbf{p}}{dt} = \frac{d}{dt} \sum_i \mathbf{p}_i = 0$  and  $\mathbf{f} = \sum_i \mathbf{f}_i = 0$
- Rotational symmetry: Conservation of angular momentum  
 $\frac{d\mathbf{L}}{dt} = \frac{d}{dt} \sum_i \mathbf{l}_i = 0$  and  $\tau = \sum_i \tau_i = 0$
- Time translational symmetry: Conservation of mechanical energy  
 $\frac{dE}{dt} = 0$



## Summary:

- Translational symmetry = Conservation of linear momentum  
 $\frac{d\mathbf{p}}{dt} = \frac{d}{dt} \sum_i \mathbf{p}_i = 0$  and  $\mathbf{f} = \sum_i \mathbf{f}_i = 0$
- Rotational symmetry: Conservation of angular momentum  
 $\frac{d\mathbf{L}}{dt} = \frac{d}{dt} \sum_i \mathbf{l}_i = 0$  and  $\tau = \sum_i \tau_i = 0$
- Time translational symmetry: Conservation of mechanical energy  
 $\frac{dE}{dt} = 0$

**Conserved quantities are not an accident but directly stem from symmetries!**

This was discovered by Emmy Noether in 1918, and is one of the most important findings in modern physics!

## Summary:

- Translational symmetry = Conservation of linear momentum  
 $\frac{d\mathbf{p}}{dt} = \frac{d}{dt} \sum_i \mathbf{p}_i = 0$  and  $\mathbf{f} = \sum_i \mathbf{f}_i = 0$
- Rotational symmetry: Conservation of angular momentum  
 $\frac{d\mathbf{L}}{dt} = \frac{d}{dt} \sum_i \mathbf{l}_i = 0$  and  $\tau = \sum_i \tau_i = 0$
- Time translational symmetry: Conservation of mechanical energy  
 $\frac{dE}{dt} = 0$

**Conserved quantities are not an accident but directly stem from symmetries!**

This was discovered by Emmy Noether in 1918, and is one of the most important findings in modern physics!

Spoiler: In MD, we will sometimes break these symmetries and thus induce energy drifts!