

Six Functors and Gestalten

Talk 10 — Gestalten Seminar

after [Sch26], Lecture IX (*up to the extension to stacks*)

Speaker: OU LIU Date: 22 June 2026

Goal of the talk

Starting from a (presentable) six-functor formalism $D: \text{Span}(C) \rightarrow 1\text{Pr}$, we construct *canonically* a functor

$$[-]_D: C \longrightarrow \text{Gest} := \text{StRing}^{\text{op}}, \quad X \longmapsto [X]_D = \text{Gest}(A_{D,X}),$$

the *transmutation* of D , through which D factors as “geometry + quasicoherent sheaves”. The engine is the *category of kernels* $\mathbb{K}_D$; iterating it produces the Stefanich ring $A_{D,X}$. We show $[-]_D$ commutes with finite limits, that every $[f]_D$ is 1-étale and 1-proper, and that $\mathcal{O}(-)_1 \circ [-]_D$ recovers the $(f^*, \otimes)$ -part of D (Theorem 3.4). We stop just before the extension to stacks.

Conventions

Categories. Throughout, “category” means ∞ -category, and C is a small one with finite limits.
The formalism. $D: \text{Span}(C) \rightarrow 1\text{Pr}$ is a *presentable six-functor formalism* — a lax symmetric monoidal functor on the span category — where 1Pr denotes presentable categories and $2\text{Pr} = \text{Mod}_{1\text{Pr}}(1\text{Pr})$.

Kernels and Gestalten. $\mathbb{K}_D$ is the *category of kernels* (Section A) and $\text{Gest} = \text{StRing}^{\text{op}}$. We write $\mathbb{1}$ for the monoidal unit, and $\mathcal{O}(-)_1$ (also written D_{qc}) for “global sections / quasicoherent sheaves” — the bottom floor of a Stefanich ring.

D-properties. A morphism is *D-suave*, *D-prim*, *D-smooth*, *D-étale*, or *D-proper* when it has the corresponding property for D (Section A); the unprefix adjectives (smooth, étale, proper, ...) denote the underlying geometric notions.

1 From motives to six-functor formalisms

In algebraic geometry there are many cohomology theories: the ℓ -adic ones, one for each prime ℓ ; crystalline cohomology; de Rham and Betti cohomology. Grothendieck envisioned a *universal* system of coefficients that would dictate all the rest — an initial cohomology theory out of which every other one is read off. In its original, abelian-categorical form, the idea asks for a category Mot_k

of *motives*, together with an assignment $X \mapsto \text{Mot}(X)$ (the *motive* of X), such that every reasonable “cohomology functor” $h^\bullet: \text{Sch}_k^{\text{op}} \rightarrow \mathbb{A}$ valued in an abelian category $\mathbb{A}$ factors through it:

$$\begin{array}{ccc} \text{Sch}_k^{\text{op}} & \xrightarrow{\text{Mot}} & \text{Mot}_k \\ & \searrow h^\bullet & \downarrow \text{---} \\ & & \mathbb{A}. \end{array}$$

The dashed leg is a *realization*: each Weil cohomology — Betti, de Rham, ℓ -adic — is one realization of the universal motive. All the variation *among* cohomology theories is carried by this choice of realization, while the geometric input $X \mapsto \text{Mot}(X)$ stays fixed. This shape of factorization — a fixed universal object, recovered into each concrete theory by a single leg — is the one we will keep meeting; the goal of the talk is its modern, categorified incarnation.

The first move is to categorify the coefficients. Studying a category C of geometric objects — say schemes of finite type over $\mathbb{Z}$ — on its own is hard; one probes it through linear invariants. The bare association $X \mapsto H^\bullet(X)$ is coarse, since many non-isomorphic spaces share the same cohomology. One remembers far more by attaching to each X a whole category of *coefficients*: *local systems* $\mathcal{L}$, which themselves carry cohomology $H^\bullet(X, \mathcal{L})$ and are transported along a map $f: Y \rightarrow X$ by fiberwise cohomology $f_*\mathcal{L}$. To allow arbitrary maps, not just smooth proper ones, one enlarges these into a category of *sheaves*. Cohomology then acquires a dual companion, *cohomology with compact support* $H_c^\bullet$, already visible in Poincaré duality.

The structure that packages all of this is a *six-functor formalism* (defined carefully in Section A). Informally, it assigns to each $X \in C$ a symmetric monoidal *category of coefficients* $D(X)$ — in our setting $D(X) \in \mathbf{1Pr}$ — and to each map $f: Y \rightarrow X$ two adjoint pairs

$$f^* \dashv f_*, \quad f_! \dashv f^!,$$

encoding pullback, pushforward (cohomology), exceptional pushforward (cohomology with compact support), and exceptional pullback, all tied to the tensor product $\otimes$ by base change and a projection formula.

We now look for the categorified analogue of Mot_k : a fixed target through which *every* six-functor formalism factors. A first candidate is the category AnStck of *analytic stacks*, through which the $(f^*, \otimes)$ -part $D^{*, \otimes}$ often factors. Grothendieck and Scholze are then after the same thing: a universal coefficient through which every theory factors. They differ only in *which leg of the triangle they hold fixed*:

	Grothendieck: motives	Scholze: six functors
factorization	$\begin{array}{ccc} \mathrm{Sch}_k^{\mathrm{op}} & \xrightarrow{\mathrm{Mot}} & \mathrm{Mot}_k \\ & \searrow h^* & \downarrow \mathrm{real.} \\ & & \mathbb{A} \end{array}$	$\begin{array}{ccc} C^{\mathrm{op}} & \xrightarrow{X \mapsto X^?} & \mathrm{AnStck}^{\mathrm{op}} \\ & \searrow D^{*, \otimes} & \downarrow D_{qc} \\ & & \mathrm{CAlg}(\mathrm{1Pr}) \end{array}$
universal coefficient	Mot_k	AnStck (later Gest)
front leg	$\mathrm{Mot} \text{ --- fixed}$	transmutation $X \mapsto X^? \text{ --- varies with } D$
recovering leg	realization $\mathrm{Mot}_k \rightarrow \mathbb{A} \text{ --- varies}$	$D_{qc} = \mathcal{O}(-)_1 \text{ --- fixed}$
variation lives in	the realization	the transmutation

The same triangle, with the fixed and varying legs interchanged. Scholze packages all the realizations into the single theory-independent leg $\mathcal{O}(-)_1 = D_{qc}$ — the same for every D , the leg that reads the $(f^*, \otimes)$ -part of D back off the transmutation (Theorem 3.4) — and studies the front leg instead: the *transmutation* $X \mapsto X^?$, after Bhatt [Bha22, Remark 2.3.8], which converts a geometric object into its universal coefficient and commutes with finite limits (equivalently, its opposite commutes with finite colimits). So the classical Betti and de Rham realizations are just $\mathcal{O}(-)_1$ applied to the transmutations by the Betti and de Rham ring stacks. The one caveat is the dashed top leg: AnStck achieves this only for *some* D .

Before turning to the obstruction, we record when the dashed leg *does* exist. For schemes (separated of finite type over a field, say) such transmutations are produced from a *ring stack*, by the following criterion of Scholze.

Theorem 1.1 ([Sch25, Theorem 10.6]). *Work in the category of analytic stacks over some base analytic stack S . Let k be a ring and let R be a k -algebra stack satisfying the following conditions.*

- (1) *The map $f: R \rightarrow *$ is $!$ -able and D -smooth, and the map $\mathbb{1} \rightarrow f_! f^! \mathbb{1}$ is an equivalence.*
- (2) *The origin $i: * \xrightarrow{0} R$ is a closed immersion, the units $j: R^\times \hookrightarrow R$ form its open complement, and*

$$j_! \mathbb{1} \rightarrow \mathbb{1} \rightarrow i_* \mathbb{1}$$

is a fiber sequence in $D_{qc}(R)$.

This induces a transmutation functor

$$X \mapsto X_R$$

taking any separated scheme X of finite type over k to the static (i.e. 0-truncated) analytic stack over S given by the sheafification of the functor

$$\mathrm{AnRing}_S \ni A \mapsto X(R(A)).$$

The functor $X \mapsto X_R$ commutes with finite limits and Zariski gluing. Moreover, the functor $X \mapsto X_R$ sends étale maps to D-étale maps, proper maps to D-proper maps, and smooth maps to D-smooth maps, and thus $X \mapsto D_{qc}(X_R)$ yields a six-functor formalism with these properties.

For instance, the Betti stack and the solid de Rham stack arise from such ring stacks.

The fixed second leg D_{qc} satisfies the *categorical Künneth formula*: by [Kes25], on analytic stacks over an affine base the external product is an equivalence.¹ Hence, if D factors as above, then D itself satisfies categorical Künneth:

$$D(X) \otimes_{D(*)} D(Y) \xrightarrow{\sim} D(X \times Y) \quad \text{for all } X, Y.$$

For SH, for IndCoh, and for ℓ -adic sheaves, however, the categorical Künneth formula *fails* — so none of these factors through AnStck in this way.

Remark 1.2 (Why the Künneth formula fails for SH, IndCoh, and ℓ -adic sheaves). In each case the external product $D(X) \otimes_{D(*)} D(Y) \xrightarrow{\boxtimes} D(X \times Y)$ fails to be an equivalence — it is not essentially surjective — so the kernel category $\mathbb{K}_D$ is strictly larger than $\text{Mod}_{D(*)}(1\text{Pr})$ (Theorem 2.3). (For ℓ -adic sheaves the external product is moreover fully faithful [Gai25]; for IndCoh and SH the cited sources assert only non-equivalence.)

- *ℓ -adic sheaves* [Gai25]. The functor $\text{Shv}(Y) \otimes \text{Shv}(Y') \xrightarrow{\boxtimes} \text{Shv}(Y \times Y')$ is *never* an equivalence once Y, Y' are positive-dimensional of finite type. For a witness take $Y' = Y$ and the diagonal $\Delta: Y \hookrightarrow Y \times Y$: the constructible sheaf $\Delta_* \mathbb{1}_Y$ is compact, with singular support the conormal $N_\Delta^* \subset T^*(Y \times Y)$. Every external product satisfies $\text{SS}(F \boxtimes G) \subseteq \text{SS}(F) \times \text{SS}(G)$ [KS90, §5.4]. But N_Δ^* is not contained in any finite union of products of conic Lagrangians, the diagonal $Y \subset Y \times Y$ being no product. Hence $\Delta_* \mathbb{1}_Y$ is not a retract of a finite colimit of external products, so it escapes the image. (For ℓ -adic coefficients, singular support is Beilinson's [Bei16].)
- *Ind-coherent sheaves* [Ste25]. For IndCoh on schemes almost of finite presentation over k , the categorification tower reaches module form only late: for an arbitrary map $X \rightarrow Y$ one has $n\text{-IndCoh}(X) \simeq \text{Mod}_{(n-1)\text{-IndCoh}(X/Y)}(n\text{-IndCoh}(Y))$ for $n \geq 3$ (the module form already appears at level 2 for closed immersions). If k is perfect then $\text{Pr}_{\text{IndCoh}} \simeq \text{Pr}$. It is the bottom level that fails, and this is precisely the failure of Künneth: over a base S the formalism $X \mapsto \text{IndCoh}(X)$ takes values in $1\text{Pr}_S = \text{Mod}_{\text{QCoh}(S)}(1\text{Pr})$, the 2-category of $\text{QCoh}(S)$ -linear presentable categories (= 2-QCoh(S)), and its right-lax monoidal structure is the external product

$$\text{IndCoh}(X_1) \otimes_{\text{QCoh}(S)} \text{IndCoh}(X_2) \longrightarrow \text{IndCoh}(X_1 \times_S X_2).$$

¹Precisely, [Kes25, Cor. 1.5.1]: over a base with representable diagonal admitting an affine $!$ -cover — in particular over an affine base — every $!$ -able object is Künneth. Over a general stacky base the *relative* formula may fail, but that is not what is at stake here.

As Gaitsgory–Rozenblyum–Varshavsky note [GRV25], even when S, X_1, X_2 are smooth this is *not* an equivalence as soon as $X_1 \times_S X_2$ fails to be smooth, since IndCoh agrees with QCoh only on smooth targets. Hence Stefanich’s $2\text{-}\mathrm{IndCoh}(S)$ properly enlarges $1\mathrm{Pr}_S$. (Contrast QCoh , which *does* satisfy Künneth on perfect stacks [BFN10, Thm. 4.14].)

- *Motivic spectra* [Aok26]. Aoki shows the kernel 2-category $\mathbb{K}_{\mathrm{SH}}$ of SH is *freely* generated by a ring stack that is homologically trivial, smooth, and sutured — and is emphatically not the unit, giving a moduli description of the genuinely 2-categorical gestalt of SH . Were Künneth to hold, $\mathbb{K}_{\mathrm{SH}}$ would collapse to the single floor $\mathrm{Mod}_{\mathrm{SH}(*)}(1\mathrm{Pr})$.

These three failures are the gap we set out to fill. We enlarge the rigid target AnStck to the category of *Gestalten* $\mathrm{Gest} = \mathrm{StRing}^{\mathrm{op}}$, and factor *every* presentable six-functor formalism D through a canonically associated transmutation:

$$\begin{array}{ccc} C^{\mathrm{op}} & \xrightarrow{\quad [-]_D \quad} & \mathrm{Gest}^{\mathrm{op}} \\ & \searrow \scriptstyle D^{*, \otimes} & \downarrow \scriptstyle \mathcal{O}(-)_1 \\ & & \mathrm{CAlg}(1\mathrm{Pr}). \end{array}$$

In contrast with the analytic-stack picture, which applies only to *some* D , this construction — following Scholze [Sch26] and Stefanich [Ste25] — works for *every* presentable D . We write its front leg $[-]_D$: this is the generic $X \mapsto X^?$ of the table, now produced canonically for every D rather than from an ad hoc ring stack R . *Gestalten* is in this sense the *universal coefficient category* for presentable six-functor formalisms, the categorified counterpart of Grothendieck’s Mot_k . Quasicoherent sheaves on Gest — the fixed functor $\mathcal{O}(-)_1$ — is the *packaged realization* through which every $1\mathrm{Pr}$ -valued D factors. The price is that $[X]_D$ is now a genuinely higher-categorical object, obtained by iterating the category of kernels $\mathbb{K}_D$ into a *Stefanich ring* $A_{D,X}$. The very failure of Künneth that obstructed the factorization through AnStck is exactly what the higher target absorbs (Theorem 2.3) — level by level, as the next section makes precise.

The whole picture is one *ladder of categorical levels*, the skeleton of the rest of the talk: each invariant of X records data one categorical degree higher than the last — and so supports correspondingly richer algebraic operations — while the transmutation $[-]_D$ records every level at once.

Level	Invariant $X \mapsto$	n -categorical algebra	Available operations
0	$H^\bullet(X)$	cohomology ring (numbers)	$+, \times$
1	$D(X)$	symmetric monoidal category	limits, colimits, $\otimes$
2	$\mathbb{K}_{D,X}$	category of kernels	limits, colimits, $\otimes$, adjoints, monads
∞	$[X]_D$	Stefanich ring (a Gestalt)	all of the above, at every level

2 The category of kernels $\mathbb{K}_D$

We now construct the finite-limit-preserving transmutation $[-]_D : C \rightarrow \text{Gest}$ at the heart of the universal-coefficient picture above. The most naive candidate uses D directly — send each X to the Gestalt of its coefficient category, i.e. compose

$$C \hookrightarrow \text{Span}(C) \xrightarrow{D} 1\text{Pr} \longrightarrow \text{Gest}, \quad X \mapsto \text{Gest}(D(X))$$

(precisely, the $(f^*, \otimes)$ -part $D^{*, \otimes} : C^{\text{op}} \rightarrow \text{CAlg}(1\text{Pr})$ postcomposed with the bottom-floor inclusion $\text{CAlg}(1\text{Pr}) \hookrightarrow \text{StRing} = \text{Gest}^{\text{op}}$). But this need not preserve pullbacks: the product $X \times Y$ is sent to $\text{Gest}(D(X \times Y))$, whereas the product of $\text{Gest}(D(X))$ and $\text{Gest}(D(Y))$ in Gest is $\text{Gest}(D(X) \otimes_{D(*)} D(Y))$ — products in $\text{Gest} = \text{StRing}^{\text{op}}$ are relative tensor products of coefficients. The two coincide exactly when D satisfies the *categorical Künneth formula* $D(X) \otimes_{D(*)} D(Y) \xrightarrow{\sim} D(X \times Y)$ — which, as Theorem 1.2 showed, generally fails (Theorem 2.3). The *category of kernels* $\mathbb{K}_D$ is the universal fix, but it acts one categorical degree at a time. Passing from D to $\mathbb{K}_D$ need not make Künneth hold outright; it *lifts* the obstruction one categorification up, so that the kernel theory $X \mapsto \mathbb{K}_{D,X}$ may itself fail Künneth at the next level. Iterating the construction into the Stefanich ring (Section 3) then clears the obstruction level by level, yielding a transmutation that genuinely preserves finite limits.

The kernel 2-category is constructed in Section A (Theorem A.9); here $\mathbb{K}_D = \mathbb{K}_{D,*}$ is its absolute case, over the terminal object of C .

Definition 2.1. Let C be a category with finite limits. Regarding the six-functor formalism D as a commutative algebra object in $\text{Fun}(\text{Span}(C), 1\text{Pr})$, equipped with the Day convolution symmetric monoidal structure, the *2-category of kernels of D* is the module category

$$\mathbb{K}_D := \text{Mod}_D(\text{Fun}(\text{Span}(C), 1\text{Pr})).$$

This comes with a symmetric monoidal functor from $\text{Span}(C)^{\text{op}}$, given as the composite

$$\text{Span}(C)^{\text{op}} \rightarrow \text{Fun}(\text{Span}(C), 1\text{Pr}) \rightarrow \text{Mod}_D(\text{Fun}(\text{Span}(C), 1\text{Pr})) = \mathbb{K}_D,$$

namely the Yoneda embedding followed by the free D -module functor $D \otimes (-)$. We denote it by $X \mapsto [X]_{\mathbb{K}_D}$. Since $[X]_{\mathbb{K}_D}$ is the free D -module on the representable presheaf at X , it is easy to map out of: by the free–forgetful adjunction composed with the Yoneda lemma, for any $X \in \text{Span}(C)^{\text{op}}$ and $M \in \mathbb{K}_D$ one has

$$\text{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, M) = M(X).$$

In particular, as the unit object of $\mathbb{K}_D$ is D itself, taking $M = \mathbb{1}$ gives

$$\text{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, \mathbb{1}) = D(X).$$

Now every object of $\text{Span}(C)$ is canonically self-dual ([HM24, p. 2.3.9]), and the embedding is symmetric monoidal, so $[Y]_{\mathbb{K}_D} \simeq [Y]_{\mathbb{K}_D}^\vee$ and $[X]_{\mathbb{K}_D} \otimes [Y]_{\mathbb{K}_D} \simeq [X \times Y]_{\mathbb{K}_D}$. Hence in general

$$\text{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, [Y]_{\mathbb{K}_D}) \simeq \text{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D} \otimes [Y]_{\mathbb{K}_D}^\vee, \mathbb{1}) \simeq \text{Hom}_{\mathbb{K}_D}([X \times Y]_{\mathbb{K}_D}, \mathbb{1}) = D(X \times Y),$$

which recovers Theorem A.9 (there the mapping object is written $\text{Fun}_Y(X_1, X_2) = D(X_1 \times_Y X_2)$; here Y is the terminal object, so $\text{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, [Y]_{\mathbb{K}_D})$ is the absolute case $\text{Fun}_*(X, Y)$). The direct computation above already matches the two mapping objects; the following theorem of Aoki upgrades this to a canonical equivalence of the whole 2-categories.

Theorem 2.2 (Aoki, [Aok26, Theorem 2.6]). *Let S be a small closed symmetric monoidal category and $V \in \text{CAlg}(\text{Pr}^L)$. Let $D: S \rightarrow V$ be a lax symmetric monoidal functor. Then there is a canonical equivalence*

$$\text{PSh}^V(\tilde{S}^{\text{op}}) \simeq \text{Mod}_D(\text{Fun}(S, V)),$$

where $\tilde{S}$ is the V -enriched category obtained by base changing the self-enrichment of S along D .

Proof. Aoki credits this argument to H. Heine; we expand it, isolating the monadicity step and computing the resulting monad by hand.

Day convolution and the trivial enrichment. Equip $\text{Fun}(S, V)$ with the Day convolution symmetric monoidal structure, for which a lax symmetric monoidal functor $S \rightarrow V$ is the same datum as a commutative algebra; thus $D \in \text{CAlg}(\text{Fun}(S, V))$. Writing $U: \text{An} \rightarrow V$ for the unit functor — the symmetric monoidal left adjoint with $U(*) = \mathbb{1}_V$ — the pointwise category $\text{Fun}(S, V)$ is the V -presheaf category of S for its *trivial* enrichment S^{triv} ,

$$\text{Fun}(S, V) = \text{PSh}^V((S^{\text{triv}})^{\text{op}}), \quad \underline{\text{Hom}}_{S^{\text{triv}}}(a, b) = U(\text{Hom}_S(a, b)).$$

Change of enrichment. As S is closed it is enriched over itself with mapping objects $\underline{\text{Hom}}_S(a, b)$, and a lax monoidal $G: S \rightarrow V$ base-changes this self-enrichment to the V -enrichment $a, b \mapsto G(\underline{\text{Hom}}_S(a, b))$. Taking $G = U(\text{Hom}_S(\mathbb{1}, -))$ recovers S^{triv} (since $\text{Hom}_S(\mathbb{1}, \underline{\text{Hom}}_S(a, b)) \simeq \text{Hom}_S(a, b)$), while $G = D$ yields $\tilde{S}$, with $\underline{\text{Hom}}_{\tilde{S}}(a, b) = D(\underline{\text{Hom}}_S(a, b))$. Now $U(\text{Hom}_S(\mathbb{1}, -))$ is the unit of $\text{CAlg}(\text{Fun}(S, V))$, so it admits a unique algebra map to D ; base changing along it gives a V -enriched, identity-on-objects functor $S^{\text{triv}} \rightarrow \tilde{S}$, whose opposite we call f . Being a base change of enrichment along a map of commutative algebras, f is symmetric monoidal, and passing to enriched presheaves yields an adjunction

$$f_! \dashv f^* \quad \text{in } \text{Mod}_V(\text{Pr}^L),$$

with $f^*: \text{PSh}^V(\tilde{S}^{\text{op}}) \rightarrow \text{Fun}(S, V)$ the restriction of enrichment (lax monoidal, V -linear) and $f_!$ its V -linear, strong monoidal left adjoint, the enriched left Kan extension. Since $\underline{\text{Hom}}_S(\mathbb{1}, a) \simeq a$, restriction of the monoidal unit recovers D :

$$(f^* \mathbb{1})(a) = \underline{\text{Hom}}_{\tilde{S}}(\mathbb{1}, a) = D(\underline{\text{Hom}}_S(\mathbb{1}, a)) \simeq D(a), \quad \text{i.e. } f^* \mathbb{1} \simeq D \text{ in } \text{CAlg}(\text{Fun}(S, V)).$$

Monadicity. Restriction along an identity-on-objects functor is computed objectwise, so f^* preserves all colimits and is conservative — a map of $\tilde{S}$ -presheaves whose restriction to the (unchanged) collection of objects is an equivalence was already an objectwise equivalence. It is moreover V -linear and admits the left adjoint $f_!$, so the monadicity theorem for presentable module categories of Reutter–Zetto applies: the Eilenberg–Moore comparison is fully faithful [RZ25, Thm. 4.7] and, under these hypotheses, an equivalence [RZ25, Thm. 4.12]. Hence f^* is monadic, and with $T := f^* f_!$ the resulting V -linear monad on $\text{Fun}(S, V)$,

$$\text{PSh}^V(\tilde{S}^{\text{op}}) \simeq \text{Mod}_T(\text{Fun}(S, V)).$$

It remains to identify T — first as an endofunctor, then as a monad.

The underlying endofunctor. Because f is the identity on objects, f^* does not move objects and $T(P)(s') = (f_! P)(s')$; the coend formula for the enriched left Kan extension $f_!$ then gives, for $P \in \text{Fun}(S, V)$ and $s' \in S$,

$$T(P)(s') \simeq \int^{a \in S} \underline{\text{Hom}}_S(a, s') \otimes P(a) = \int^a \text{D}(\underline{\text{Hom}}_S(a, s')) \otimes P(a).$$

On the other hand, Day convolution gives

$$(P \otimes \text{D})(s') = \int^{a, b} \text{Hom}_S(a \otimes b, s') \otimes P(a) \otimes \text{D}(b).$$

Integrating over b first (Fubini for coends): for each fixed a , the closed-structure equivalence $\text{Hom}_S(a \otimes b, s') \simeq \text{Hom}_S(b, \underline{\text{Hom}}_S(a, s'))$ followed by the co-Yoneda lemma $\int^b \text{Hom}_S(b, Y) \otimes \text{D}(b) \simeq \text{D}(Y)$ at $Y = \underline{\text{Hom}}_S(a, s')$ collapses the inner coend to $\text{D}(\underline{\text{Hom}}_S(a, s'))$, whence

$$(P \otimes \text{D})(s') \simeq \int^a P(a) \otimes \text{D}(\underline{\text{Hom}}_S(a, s')).$$

The two displays agree, naturally in s' , so $T \simeq (-) \otimes \text{D}$ as endofunctors of $\text{Fun}(S, V)$ — obtained from the closed structure of S alone, with no rigidity or separate projection formula. (Equivalently, by V -linearity it would suffice to compare the two on representables; the coend does it on all of $\text{Fun}(S, V)$ at once.)

The monad structure, and conclusion. We promote this to an equivalence of monads. As f^* is lax monoidal with $f^* \mathbb{1} \simeq \text{D}$, the monoidal adjunction supplies a canonical comparison: for $P \in \text{Fun}(S, V)$,

$$P \otimes \text{D} = P \otimes f^* \mathbb{1} \xrightarrow{\eta_P \otimes \text{id}} f^* f_! P \otimes f^* \mathbb{1} \xrightarrow{\text{lax}} f^* (f_! P \otimes \mathbb{1}) \simeq f^* f_! P = T(P),$$

where η is the unit of $f_! \dashv f^*$. This is a morphism of monads $(-) \otimes \text{D} \rightarrow T$: the unit $\mathbb{1} \rightarrow \text{D}$ of the algebra furnishes its unit, and the multiplication $\text{D} \otimes \text{D} \rightarrow \text{D}$ — the lax constraint of

$D = f^* \mathbb{1}$ assembled over the Day coend — its multiplication. By the previous step its underlying natural transformation is the equivalence $(-) \otimes D \simeq T$, and a monad morphism that is an underlying equivalence is an equivalence of monads. Therefore

$$\mathrm{PSh}^V(\tilde{S}^{\mathrm{op}}) \simeq \mathrm{Mod}_T(\mathrm{Fun}(S, V)) \simeq \mathrm{Mod}_{(-) \otimes D}(\mathrm{Fun}(S, V)) \simeq \mathrm{Mod}_D(\mathrm{Fun}(S, V)),$$

the last identification being $\mathrm{Mod}_{(-) \otimes A} \simeq \mathrm{Mod}_A$ for the free-module monad of an algebra A . Every step is canonical, so the equivalence is. $\square$

Now, taking $V = \mathbf{1Pr}$ and $S = \mathrm{Span}(C)$, we obtain the desired result.

Proposition 2.3 ([Sch26, Prop. 9.3]). *The natural functor*

$$\mathrm{Mod}_{D(*)}(\mathbf{1Pr}) \longrightarrow \mathbb{K}_D$$

is fully faithful, with essential image the “restricted²” objects: $M \in \mathbb{K}_D$ lies in it iff $D(X) \otimes_{D()} M(*) \rightarrow M(X)$ is an equivalence for all X . In particular,*

$$\mathbb{K}_D = \mathrm{Mod}_{D(*)}(\mathbf{1Pr}) \iff D(X) \otimes_{D(*)} D(Y) \xrightarrow{\sim} D(X \times Y) \text{ for all } X, Y.$$

Proof. We work with the description $\mathbb{K}_D = \mathrm{Mod}_D(\mathrm{Fun}(\mathrm{Span}(C), \mathbf{1Pr}))$ of Theorem 2.1 and the formula $\mathrm{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, M) = M(X)$ established there.

Evaluation is atomic. For each X , that formula says the corepresentable of $[X]_{\mathbb{K}_D}$ is the evaluation functor

$$\mathrm{ev}_X = \mathrm{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, -) : \mathbb{K}_D \longrightarrow \mathbf{1Pr}, \quad M \longmapsto M(X).$$

Colimits in $\mathbb{K}_D = \mathrm{Mod}_D(\mathrm{Fun}(\mathrm{Span}(C), \mathbf{1Pr}))$ are created by the forgetful functor to $\mathrm{Fun}(\mathrm{Span}(C), \mathbf{1Pr})$, where they are computed pointwise; hence ev_X preserves all colimits, and it is $\mathbf{1Pr}$ -linear. Thus every $[X]_{\mathbb{K}_D}$ is $\mathbf{1Pr}$ -atomic. In particular the unit $\mathbb{1} = [*]_{\mathbb{K}_D}$ is $\mathbf{1Pr}$ -atomic, with $\mathrm{End}_{\mathbb{K}_D}(\mathbb{1}) = \mathrm{ev}_*(\mathbb{1}) = D(*)$.

The natural functor and its right adjoint. Since $\mathrm{ev}_*(M) = M(*)$ is naturally a module over $\mathrm{End}_{\mathbb{K}_D}(\mathbb{1}) = D(*)$, the functor ev_* refines to a colimit-preserving functor

$$G : \mathbb{K}_D \longrightarrow \mathrm{Mod}_{D(*)}(\mathbf{1Pr}), \quad M \longmapsto M(*).$$

The functor of the statement is its left adjoint — induction along $D(*) = \mathrm{End}_{\mathbb{K}_D}(\mathbb{1}) \rightarrow \mathbb{K}_D$,

$$L : \mathrm{Mod}_{D(*)}(\mathbf{1Pr}) \longrightarrow \mathbb{K}_D, \quad N \longmapsto N \otimes_{D(*)} \mathbb{1},$$

the unique $\mathbf{1Pr}$ -linear colimit-preserving functor with $L(D(*)) = \mathbb{1}$; one has $L \dashv G$.

²in the sense of [Gai25]

Full faithfulness. L is fully faithful iff the unit $N \rightarrow GL(N)$ of the adjunction is an equivalence for every N . Here

$$GL(N) = (N \otimes_{D(*)} \mathbb{1})(*) \simeq N \otimes_{D(*)} \mathbb{1}(*) = N \otimes_{D(*)} D(*) = N.$$

The middle equivalence is where atomicity enters: ev_* is colimit-preserving and 1Pr -linear, so it commutes with the relative tensor product $N \otimes_{D(*)} (-)$ (a geometric realization of the bar construction); and $\mathbb{1}(*) = D(*)$. Hence $GL \simeq \text{id}$ and L is fully faithful.

The essential image. As L is fully faithful with right adjoint G , an object $M \in \mathbb{K}_D$ lies in the essential image of L iff the counit $\varepsilon_M: LG(M) \rightarrow M$ is an equivalence. Evaluating $LG(M) = M(*) \otimes_{D(*)} \mathbb{1}$ at X — again using that ev_X commutes with $M(*) \otimes_{D(*)} (-)$, together with $\mathbb{1}(X) = D(X)$ —

$$(LG(M))(X) \simeq M(*) \otimes_{D(*)} \mathbb{1}(X) = D(X) \otimes_{D(*)} M(*),$$

and ε_M at X is precisely the canonical map $D(X) \otimes_{D(*)} M(*) \rightarrow M(X)$. So M is in the image — i.e. *restricted* — iff this map is an equivalence for all X , as claimed.

The Künneth criterion. The restricted objects are closed under colimits: LG and id both preserve colimits, so the locus where the natural transformation ε is invertible is closed under colimits. Since $\mathbb{K}_D$ is generated under colimits by the representables $[Y]_{\mathbb{K}_D}$, the functor L is essentially surjective — equivalently an equivalence, i.e. $\mathbb{K}_D = \text{Mod}_{D(*)}(1\text{Pr})$ — iff every $[Y]_{\mathbb{K}_D}$ is restricted. Finally, by the $\text{Hom}_{\mathbb{K}_D}$ -formula,

$$[Y]_{\mathbb{K}_D}(*) = \text{Hom}_{\mathbb{K}_D}(\mathbb{1}, [Y]_{\mathbb{K}_D}) = D(Y), \quad [Y]_{\mathbb{K}_D}(X) = \text{Hom}_{\mathbb{K}_D}([X]_{\mathbb{K}_D}, [Y]_{\mathbb{K}_D}) = D(X \times Y),$$

so $[Y]_{\mathbb{K}_D}$ is restricted iff $D(X) \otimes_{D(*)} D(Y) \rightarrow D(X \times Y)$ is an equivalence for all X . Ranging over Y as well, this is exactly the categorical Künneth formula. $\square$

3 Transmutation

We now construct the transmutation functor $C^{\text{op}} \rightarrow \text{Gest}^{\text{op}} = \text{StRing}$. Note that for any $X \in C$, we can repeat Theorem 2.1 with the slice $C_{/X}$, defining

$$\mathbb{K}_{D,X} := \text{Mod}_D(\text{Fun}(\text{Span}(C_{/X}), 1\text{Pr})) \in \text{CAlg}(2\text{Pr}).$$

The pullback functor $\text{Span}(C) \rightarrow \text{Span}(C_{/X})$ induces, by passing to free presentable 2-categories, a symmetric monoidal functor

$$\text{Fun}(\text{Span}(C), 1\text{Pr}) \rightarrow \text{Fun}(\text{Span}(C_{/X}), 1\text{Pr}), \quad F \mapsto F|_{C_{/X}}.$$

Because D is compatible with this restriction, the two routes to D -modules — restrict then take modules, or take modules then restrict — agree; there is only one sense of “passing to modules over D ”.

For any map $f: Y \rightarrow X$ in C , the pullback functor $f^*: C/X \rightarrow C/Y$ induces a symmetric monoidal functor

$$f_1^*: \mathbb{K}_{D,X} \rightarrow \mathbb{K}_{D,Y},$$

so this assembles into a functor

$$C^{\text{op}} \rightarrow \text{CAlg}(2\text{Pr}).$$

Proposition 3.1 ([Sch26, Prop. 9.4]). *For every $f: Y \rightarrow X$ the functor $f_1^*: \mathbb{K}_{D,X} \rightarrow \mathbb{K}_{D,Y}$ admits isomorphic left and right adjoints $f_{1!} \cong f_{1*}$, whose formation commutes with base change.*

Proof. Existence of the adjoints. Forgetting the Mod_D -structure to $\text{Fun}(\text{Span}(C/X), 1\text{Pr})$, the functor f_1^* is restriction (precomposition) along the span functor $\text{Span}(u): \text{Span}(C/Y) \rightarrow \text{Span}(C/X)$, where $u: C/Y \rightarrow C/X$ is the forgetful functor left adjoint to base change ($u \dashv f^*$); both u and f^* preserve pullbacks, so $\text{Span}(u)$ and $\text{Span}(f^*)$ exist. Restriction along $\text{Span}(u)$ has as left and right adjoints the Kan extensions $f_{1!} = \text{Span}(u)_!$ and $f_{1*} = \text{Span}(u)_*$, which exist because the target 1Pr is presentable. These adjoints are D -linear and survive $\text{Mod}_D(-)$ — passing to modules over D is compatible with base change — giving $f_{1!} \dashv f_1^* \dashv f_{1*}$ between $\mathbb{K}_{D,X}$ and $\mathbb{K}_{D,Y}$.

The ambient 3-category. We prove $f_{1!} \cong f_{1*}$ by ambidexterity, and the correct ambient for this is the 3-category $2\text{Pr} = \text{Mod}_{1\text{Pr}}(1\text{Pr})$: its objects are the presentable 2-categories such as the $\mathbb{K}_{D,X}$, its 1-morphisms the 1Pr -linear functors such as f_1^* , its 2-morphisms the 1Pr -linear transformations, its 3-morphisms the modifications. Crucially we do *not* work inside a single $\mathbb{K}_{D,X}$, which is only a 2-category, where the unit and counit of $f_1^* \dashv f_{1*}$ would have no room to acquire adjoints; in 2Pr they are 2-morphisms and can.

The criterion, and which unit/counit it consumes. Recall the general ambidexterity criterion (after Hopkins–Lurie; [Sch26, Prop. 4.4]): a 1-morphism F of a 3-category with a chosen right adjoint F^R admits a left adjoint F^L with $F^L \simeq F^R$ as soon as the unit $\eta: \text{id} \rightarrow F^R F$ and counit $\varepsilon: F F^R \rightarrow \text{id}$ of the adjunction $F \dashv F^R$ themselves admit right adjoints. We apply it with $F = f_1^*$ and $F^R = f_{1*}$ in 2Pr ; the relevant 2-morphisms are thus the unit and counit of $f_1^* \dashv f_{1*}$ — *not* those of $f_{1!} \dashv f_1^*$.

Discharging the hypotheses. That η and ε admit right adjoints is checked by descent on the categorical level. By evaluation at the monoidal unit, the unit and counit of the level-2 adjunction $f_1^* \dashv f_{1*}$ are identified with those of the analogous restriction/Kan-extension adjunction one categorical level down, inside $\text{Fun}(\text{Span}(C/_), 1\text{Pr})$, where their right-adjointability is the adjointability of the relevant span correspondences ($\text{Span}(u)$ and the legs of the spans). At the bottom level this holds by the self-duality of span categories: every object of $\text{Span}(C/Y)$ is self-dual, every correspondence admits both adjoints (its transpose), and the transposition equivalence fixes $\text{Span}(u)$, so the data is compatible with $\text{Span}(u)$ and propagates back up through $\text{Mod}_D(-)$. Hence η, ε admit right adjoints in 2Pr .

Ambidexterity. The criterion therefore applies and yields a left adjoint F^L of f_1^* with $F^L \simeq f_{1*}$; a left adjoint being unique, $F^L \simeq f_{1!}$, whence the canonical equivalence $f_{1!} \simeq f_{1*}$.

Base change. For a pullback square in C the induced square of slice forgetful functors is adjointable and the mate of the base-change square is an equivalence; passing through $\text{Span}(-)$ and $\text{Mod}_D(-)$ (both base-change compatible) transports this to the Kan-extension adjoints, so the formation of $f_{1!}$ and of f_{1*} commutes with base change. Since $f_{1!} \simeq f_{1*}$ is the canonical equivalence extracted from the same adjunction data, it is natural in f and commutes with base change as well. $\square$

This already yields a functor $C \rightarrow \text{Gest}$, $X \mapsto \text{Gest}(\mathbb{K}_{D,X})$, valued in 2-affine Gestalten. Being 2-affine, every map is automatically 2-proper; by Theorem 3.1 it is moreover 1-prim, the right adjoint f_{1*} supplying the primness datum. It is not, however, evidently 1-suave: that would need left adjoints for the pullback $\text{Mod}_{\mathbb{K}_{D,X}}(2\text{Pr}) \rightarrow \text{Mod}_{\mathbb{K}_{D,Y}}(2\text{Pr})$ *one level up*, whose existence is not clear. And promoting 1-prim/1-suave to 1-proper/1-étale would mean applying Theorem 3.1 to the iterated diagonals of f , which requires the very Künneth formula that may fail (Theorem 2.3). Either obstruction blocks us at this level — which is what forces us to iterate the construction.

Remark 3.2 (Why $\text{Gest}(\mathbb{K}_{D,X})$ is 2-affine). By [Sch26, Def. 1.18] a Stefanich ring is a sequence $A = (A_1, A_2, \dots)$ with $A_n \in \text{CAlg}(n\text{Pr})$ and $A_n \simeq \text{End}_{A_{n+1}}(\mathbb{1})$ — where $0\text{Pr} = \text{An}$, $1\text{Pr} = 1\text{Pr}$ and $(n+1)\text{Pr} = \text{Mod}_{n\text{Pr}}(1\text{Pr})$ — and $\mathcal{O}(\text{Gest}(A))_n = A_n$. On the *unit* Stefanich ring its terms are the units $\mathbb{1}_{n\text{Pr}} = (n-1)\text{Pr}$, so the relevant term is $A_3 = 2\text{Pr}$.

By [Sch26, Def. 6.5], B is *n-affine* if it lies in the essential image of $\text{CAlg}(A_{n+1}) \rightarrow \text{StRing}_A$; equivalently, B is generated by its single component $B_n \in \text{CAlg}(A_{n+1})$ through $B_{m+1} \simeq \text{Mod}_{B_m}(A_{m+1})$ for $m \geq n$, the lower terms being recovered by looping. In words: an n -affine Gestalt carries no new categorical data above level n .

Now $\mathbb{K}_{D,X} = \text{Mod}_D(\text{Fun}(\text{Span}(C/X), 1\text{Pr}))$ is a commutative algebra in 2Pr , hence $\mathbb{K}_{D,X} \in \text{CAlg}(2\text{Pr}) = \text{CAlg}(A_3)$ — exactly such a generator, at level 2. So $\text{Gest}(\mathbb{K}_{D,X})$ is Gest applied to

$$(D(X), \mathbb{K}_{D,X}, \text{Mod}_{\mathbb{K}_{D,X}}(2\text{Pr}), \dots), \quad D(X) = \text{End}_{\mathbb{K}_{D,X}}(\mathbb{1}),$$

whose only genuine datum is $\mathbb{K}_{D,X}$: the higher terms are iterated module categories and $D(X)$ is the looping. This is 2-affineness — the higher analogue of an affine scheme $\text{Spec } R$, whose tower $(D(R), \text{Mod}_{D(R)}(1\text{Pr}), \dots)$ is likewise iterated modules over one ring.

The level is sharp, and reconnects to Theorem 2.3. $\text{Gest}(\mathbb{K}_{D,X})$ would already be 1-affine — that is, $\mathbb{K}_{D,X} \simeq \text{Mod}_{D(X)}(1\text{Pr})$ — precisely when the categorical Künneth formula holds over C/X . When it fails, $\mathbb{K}_{D,X} \supsetneq \text{Mod}_{D(X)}(1\text{Pr})$ and $\text{Gest}(\mathbb{K}_{D,X})$ is genuinely 2-affine, a 2-stack morally at the level of $B^2\mathbb{G}_m$. The full transmutation $[X]_D = \text{Gest}(A_{D,X})$ is, by contrast, n -affine for no finite n : the genuine data added at each higher level, wherever Künneth fails, is exactly what the iteration records.

The failure diagnosed before the remark — that $\text{Gest}(\mathbb{K}_{D,X})$ is only 1-prim, not yet 1-suave or 1-proper — is repaired by climbing one categorical level and repeating the construction. The assignment

$$C^{\text{op}} \longrightarrow \text{CAlg}(2\text{Pr}), \quad X \mapsto \mathbb{K}_{D,X},$$

itself admits, by Theorem 3.1, base-change-compatible left adjoints $f_1!$ ($\cong f_{1*}$) for every f ; it therefore upgrades to a once-categorified six-functor formalism

$$\mathrm{Span}(C) \longrightarrow 2\mathrm{Pr} = \mathrm{Mod}_{1\mathrm{Pr}}(1\mathrm{Pr}),$$

valued one categorical degree above D . Applying the kernel construction to this sheaf theory produces the level-3 data, and iterating level by level yields — following Stefanich [Ste25] — a *Stefanich ring*

$$A_{D,X} = (D(X), \mathbb{K}_{D,X}, \dots) \in \mathrm{StRing}, \quad \text{functorial in } X, \text{ i.e. a functor } C^{\mathrm{op}} \rightarrow \mathrm{StRing}.$$

This is the level- ∞ row of the ladder of categorical levels from the introduction: the single object that records every categorical degree of X at once.

Definition 3.3 (Transmutation). With $\mathrm{Gest} := \mathrm{StRing}^{\mathrm{op}}$, the *transmutation* of D is

$$[-]_D : C \longrightarrow \mathrm{Gest}/_{\mathrm{Gest}(A_D)}, \quad X \longmapsto [X]_D = \mathrm{Gest}(A_{D,X}).$$

The target is the slice over $\mathrm{Gest}(A_D) = [*]_D$, the transmutation of the point; forgetting it reads $[-]_D$ simply as the functor $C \rightarrow \mathrm{Gest}$ of the factorization diagram above.

Theorem 3.4 ([Sch26, Prop. 9.5]). *For every presentable six-functor formalism D , the transmutation $[-]_D$ satisfies:*

- (1) *it commutes with finite limits;*
- (2) *for each $f : Y \rightarrow X$ in C , the map $[f]_D : [Y]_D \rightarrow [X]_D$ is 1-étale and 1-proper, with $f_1! \cong f_{1*}$ (the left and right adjoints to the kernel pullback $f_1^* : \mathbb{K}_{D,X} \rightarrow \mathbb{K}_{D,Y}$ agreeing);*
- (3) *the composite $O(-)_1 \circ [-]_D$ recovers the $(f^*, \otimes)$ -functoriality of D .*

Moreover the construction is, in Scholze’s words, “completely canonical”.

Proof. For each $f : Y \rightarrow X$ the iteration furnishes, at every categorical level, the kernel pullback together with *both* of its adjoints, so Theorem 3.1 applies level by level: $[f]_D$ is 1-prim and 1-suave, with $f_1! \cong f_{1*}$. The extra dualizability demanded by n -suaveness is automatic, both adjoints being 1Pr-linear [Sch26, Lemma 6.22]. Promoting these to 1-properness and 1-étaleness only requires applying Theorem 3.1 to the iterated diagonals of f , legitimate once $[-]_D$ preserves finite limits.

For that it suffices to treat binary products, since a fibre product reduces to one after passing to a slice. A product in $\mathrm{Gest} = \mathrm{StRing}^{\mathrm{op}}$ is a relative tensor product of Stefanich rings, so the claim is a Künneth formula for the lower-shriek functors, one categorical level at a time. The base-change compatibility of the adjoints of Theorem 3.1 is exactly what makes each

$$C^{\mathrm{op}} \longrightarrow \mathrm{CAlg}(n\mathrm{Pr}), \quad X \longmapsto O(A_{D,X})_n,$$

a six-functor formalism (with respect to the right adjoints), and it is there that Künneth supplies the product.

Finally $\mathcal{O}(A_{D,X})_1 = D(X)$ is the bottom term of the Stefanich ring, with its $(f^*, \otimes)$ -functoriality intact, so $\mathcal{O}(-)_1 \circ [-]_D = D^{*, \otimes}$. $\square$

Warning 3.5. $\mathcal{O}(-)_1 \circ [-]_D$ recovers D only at the level of $(f^*, \otimes)$; the shriek functors are not seen by the target $\mathbf{CAlg}(1\mathbf{Pr})$. Whether the *full* formalism — the $!$ -functors included — can be reconstructed from $[-]_D$, by singling out a class of $!$ -able maps of Gestalten and a six-functor formalism on them, is the subject of the appendix to *Scholze’s Lecture IX* — not the present appendix — and is where Talk 11 takes over; the results there are, in Scholze’s words, “not definitive”.

4 Dictionary: six-functor notions vs. Gestalten

Having built $[-]_D$, we can now read the geometric vocabulary of a map f directly off its image $[f]_D$ in Gest: each six-functor property of f becomes a Gestalt-level property of $[f]_D$.

By Theorem 3.4 every $[f]_D$ is automatically 1-étale and 1-proper, with $f_{1!} \cong f_{1*}$; the categorical layers above the bottom are forced by ambidexterity and separate no maps. The whole dictionary therefore lives at layer 0, where the four Stefanich-ring conditions unwind directly to the six-functor notions of Section A. Concretely, the layer-0 datum of $[f]_D$ is the unit map $\mathbb{1} \rightarrow (B/A)_1$ of the underlying map of Stefanich rings. This is a 1-morphism $[X] \rightarrow [Y]$ in $A_2 = \mathbb{K}_{D,X}$, carrying the Fourier–Mukai kernel of f . Asking it to admit a right adjoint is 0-primness [Sch26, Def. 6.9]; asking a left adjoint is 0-suaveness [Sch26, Def. 6.16]. By Theorem A.14 these say exactly that f is D-prim, resp. D-suave. Imposing the same on all iterated diagonals of f gives 0-properness and 0-étaleness [Sch26, Def. 6.12 and 6.19], which is the recursive Theorem A.18; that recursion terminates only for truncated f , whence the extra truncatedness clause for the cohomological notions. All four equivalences then follow directly from the definitions.

Remark 4.1 ([Sch26, Rem. 9.6]). In tabular form:

f in C	$[f]_D$ in Gest
suave	0-suave
prim	0-prim
cohomologically étale	truncated $\wedge$ 0-étale
cohomologically proper	truncated $\wedge$ 0-proper

Remark 4.2 (Stack-theoretic perspective). The suave and prim notions above have a clean stack-level form, due to Aoki [Aok26, Def. 4.5]: a morphism $[X] \rightarrow [Y]$ in the kernel 2-category $\mathbb{K}_{D,Y}$ (Theorem A.9) is *weakly suave* if it admits an $[X]$ -linear left adjoint, *prim* if it admits an $[X]$ -linear right adjoint, and *suave* if it is weakly suave and $[Y]$ is dualizable over $[X]$. Aoki’s *weakly suave*

matches the object-level Theorem A.11 up to the standard reformulation “is a left adjoint” versus “admits an $[X]$ -linear left adjoint” (via the self-duality $\mathbb{K}_{D,Y} \simeq \mathbb{K}_{D,Y}^{\text{op}}$), while his *suave* adds Aoki’s dualizability of $[Y]$ over $[X]$ — equivalently, by Theorem A.16, of the suave dual — exactly the extra input promoting suaveness to cohomological smoothness. This is the stack-level reading behind the dictionary above: suaveness and primness of f become properties of $[f]_D$ in Gest.

Outlook: towards the extension to stacks

One last question, and the most satisfying. The transmutation was built object by object — but does it see the *topology*? Tracking covers through $[-]_D$ reveals where the D-topology really comes from. Recall from [Sch25, Def. 5.3] the D-topology on C : a family $\{f_i: Y_i \rightarrow X\}$ (with each f_i in E and forming a cover for the canonical topology) is a D-cover iff it moreover satisfies *universal !-descent*, i.e. after any base change $X' \rightarrow X$ the upper-shriek functor

$$D(X') \xrightarrow{\sim} \lim \left(\prod_i D(Y_i \times_X X') \rightrightarrows \prod_{i,j} D(Y_i \times_X Y_j \times_X X') \cdots \right)$$

is an equivalence.

Proposition ([Sch26, Prop. 9.7]). If $\{f_i: Y_i \rightarrow X\}$ satisfies universal !-descent, then $\coprod_i [Y_i]_D \rightarrow [X]_D$ is a cover (effective epimorphism) in Gest.

Corollary ([Sch26, Cor. 9.8]). Hence $[-]_D$ extends to a functor

$$\text{Shv}_D(C) \longrightarrow \text{Gest}/[*_]_D, \quad X \longmapsto [X]_D,$$

from D-sheaves on C (with values in 1-étale Gestalten), commuting with countable colimits and finite limits.

The upshot is that the D-topology on C is nothing other than the restriction along $[-]_D$ of the canonical topology on Gest: in distilling a Grothendieck topology from a six-functor formalism one was already seeing — if only in secret — the canonical topology on Gestalten. Constructing the six-functor formalism *on* Gest itself — and pinning down the sense in which D can be recovered from it — is where Talk 11 takes over.

One-line summary

$D \rightsquigarrow \mathbb{K}_D \rightsquigarrow A_{D,X} \rightsquigarrow [X]_D = \text{Gest}(A_{D,X})$: the six-functor formalism *is* quasicohherent sheaves on its transmutation, canonically and for all D — so Gest is the universal coefficient category for presentable six-functor formalisms.

Acknowledgements

I thank Bastiaan Cnossen and Sil Linskens for organizing this Oberseminar. I am also grateful to Claude Code (Opus 4.8) for its help in preparing these notes, and to Rethlas, which relayed my queries to Opus 4.8 and helped me track down several errors of detail in the original proofs. While in Regensburg I developed dry-eye syndrome and serious psychological difficulties, brought on by the dry climate and prolonged screen use; for this reason I relied throughout on the AI assistant Claude Code to help complete the text.

A Six-functor formalisms, smoothness, and suaveness

A modern viewpoint, originating in Grothendieck’s school, organizes “cohomology with coefficients” on a class of geometric objects C as a single coherent package: an assignment $X \mapsto D(X)$ of a category of “coefficient systems” to each object, together with six functors relating these categories under morphisms.

An informal definition. A *six-functor formalism* on C assigns to each $X \in C$ a presentable symmetric monoidal stable category $D(X)$, and to each morphism $f: Y \rightarrow X$ in C four functors

$$f^* \dashv f_*: D(Y) \rightleftarrows D(X), \quad f_! \dashv f^!: D(X) \rightleftarrows D(Y),$$

together with the symmetric monoidal product $\otimes$ and internal hom $\underline{\mathrm{Hom}}$ on each $D(X)$, such that:

- f^* is colimit-preserving and symmetric monoidal, and $f_!$ is colimit-preserving and satisfies the projection formula;
- base change: for every pullback square in C ,

$$\begin{array}{ccc} Y' & \xrightarrow{g'} & Y \\ f' \downarrow & \lrcorner & \downarrow f \\ X' & \xrightarrow{g} & X \end{array}$$

the canonical maps $g^* f_! \rightarrow f'_! g'^*$ and $f^! g_* \rightarrow g'_* f'^!$ are equivalences;

- projection formula: $f_!(A \otimes f^* B) \simeq f_!(A) \otimes B$ for $A \in D(Y)$, $B \in D(X)$.

These data, with all their higher coherences, are most cleanly encoded as a 2-functor out of a span 2-category — the precise formulation we take up in Section A.1. First we record the geometric content of each functor.

The six functors carry distinct geometric content:

- f^* is *pullback*, f_* the *direct image*, $\otimes$ the *symmetric monoidal product* encoding cup products, and $\underline{\mathrm{Hom}}$ the right adjoint to $\otimes$.
- $f_!$ is the *exceptional / proper-supported pushforward*: it agrees with f_* when f is proper and with $j_#$ (extension by zero) when f is an open immersion.
- $f^!$ is the *exceptional pullback*, encoding orientation and duality data; in many examples, for f smooth of relative dimension d , $f^! \simeq f^* \otimes \omega_f$ with ω_f the relative dualizing complex.

The three operations f_* , $f^!$, $\underline{\mathrm{Hom}}$ are *right adjoints* of f^* , $f_!$, $\otimes$ respectively, hence determined by the others. Modern formulations therefore typically specify only the three left functors $(f^*, f_!, \otimes)$ (a so-called *three-functor formalism*) and recover the rest by adjunction.

Three representative examples are:

- *Locally compact Hausdorff topological spaces* (Heyer–Mann [HM24]; see also [Sch25, Lecture VII]): here X ranges over (condensed) topological spaces and the coefficient category $D(X)$ is the derived category of sheaves of Λ -modules on X , for a fixed coefficient ring Λ .
- *ℓ -adic étale cohomology of schemes* [Sch25, Appendix to Lecture VII].
- *The motivic formalism SH* [Sch25, Lecture XI] — the universal $\mathbb{A}^1$ -invariant one, by Drew–Gallauer.

For pedagogical introductions complementing the present appendix, we recommend Scholze’s lecture notes [Sch25], Gallauer’s mini-course [Gal21] (the source of much of this section’s exposition), and the textbook treatment in Cisinski–Déglise [CD19]. The encoding via span 2-categories adopted here is closest to Mann–Scholze [Man22; Sch25] and Cnossen–Lenz–Linskens [CLL25], to which we turn in the next subsection.

A.1 Span categories and suitable decompositions

The economical encoding of a six-functor formalism, initiated by Liu–Zheng [LZ24] and brought to its present form by Mann [Man22] and Scholze [Sch25], packages all the operations $(f^*, f_!, \otimes)$ and their coherences into a single lax symmetric monoidal functor on a span category. Given a wide subcategory $E \subset C$ closed under composition and base change, the *span category* $\mathrm{Span}(C, E)$ has the same objects as C and morphisms $X \rightarrow Y$ given by spans

$$X \leftarrow Z \xrightarrow{e} Y, \quad e \in E,$$

composed by pullback; when C has finite products, $\mathrm{Span}(C, E)$ inherits a symmetric monoidal structure induced by the finite products of C , with unit the terminal object. (This monoidal structure is

not Cartesian: the tensor $X \otimes Y \simeq X \times Y$ is induced by the *product* of C , and is in general neither the categorical product nor the coproduct of $\text{Span}(C, E)$. Independently, $\text{Span}(C, E)$ is semiadditive; but its biproduct $X \oplus Y \simeq X \sqcup Y$ comes from the *coproduct* of C , a different operation from the tensor.) A *three-functor formalism* on (C, E) is then a lax symmetric monoidal functor

$$D: \text{Span}(C, E) \longrightarrow \text{Cat},$$

encoding f^* (backward legs), $f_!$ (forward legs), and $\otimes$ (monoidal structure) in a single coherent package; the right adjoints f_* , $f^!$, $\underline{\text{Hom}}$ are recovered automatically and are therefore *properties* of D rather than additional data.

The challenge is that constructing a lax monoidal functor out of $\text{Span}(C, E)$ directly is difficult. In practice, one starts from a much simpler piece of data, namely a functor

$$D_0: C^{\text{op}} \longrightarrow \text{CAlg}(\text{Cat})$$

encoding only f^* and $\otimes$, and asks when it extends to $\text{Span}(C, E)$. The key input is a *suitable decomposition* of E in the sense of Mann [Man22, Definition A.5.9]: a pair of wide subcategories $I, P \subset E$ (thought of as “étale-like” and “proper-like” morphisms respectively) such that every $e \in E$ factors as $e = p \circ i$ with $i \in I$ and $p \in P$, subject to mild cancellation and truncation axioms. The extension theorem then reads as follows. Suppose $i^* = D_0(i)$ admits a left adjoint $i_\#$ and $p^* = D_0(p)$ admits a right adjoint p_* , both satisfying the expected base change and projection formulae — in particular the mixed Beck–Chevalley equivalence $i_\# p'_* \simeq p_* i'_\#$. Then D_0 admits an essentially unique extension to a three-functor formalism D , in which

$$f_! = p_* i_\# \quad \text{for any factorization } f = p \circ i.$$

Recently, Cnossen–Lenz–Linskens [CLL25] gave a conceptually streamlined account of this extension theorem. They enhance $\text{Span}(C, E)$ to a 2-category whose 2-morphisms encode the adjunction data of $i_\#$ and p_* in a fully coherent way, and they prove that the inclusion $C^{\text{op}} \hookrightarrow \text{Span}_2(C, E)_{I, P}$ is a universal (I, P) -biadjointable functor. In this setup, Mann’s extension theorem becomes a formal corollary; as a byproduct, the unit and counit witnessing $i_\# \dashv i^*$ are part of the structure of D , rather than non-canonical choices recoverable only after the fact.


Construction A.1 (CLL 2-categorical span category). The 2-category $\text{Span}_2(C, E)_{I, P}$ has:

- *objects*: objects in C ;
- *1-morphisms* $X \rightarrow Z$: spans

$$\begin{array}{ccc} & Y & \\ f \swarrow & & \searrow g \\ X & & Z \end{array}$$

with $g \in E$;

- 2-morphisms between two such spans: diagrams



with $a \in P$ and $b \in I$.

Theorem A.2 ([CLL25, Theorem 5.17]). *Let C be a category with finite products and let $I, P \subseteq E \subseteq C$ be wide subcategories closed under base change such that*

- (1) *I and P are left cancellable;*
- (2) *every map in E factors as a map in I followed by a map in P ;*
- (3) *every map in $I \cap P$ is truncated.*

Let $D_0: C^{\text{op}} \rightarrow \text{CAlg}(\text{Cat})$ be a functor satisfying:

- (1) *for every $i: a \rightarrow b$ in I , the functor $i^*: D_0(b) \rightarrow D_0(a)$ admits a left adjoint $i_{\#}$ satisfying the Beck–Chevalley condition with respect to pullbacks in C and the projection formula $i_{\#}(- \otimes i^*(-)) \simeq i_{\#}(-) \otimes (-)$;*
- (2) *for every $p: a \rightarrow b$ in P , the functor $p^*: D_0(b) \rightarrow D_0(a)$ admits a right adjoint p_* satisfying the dual Beck–Chevalley condition and dual projection formula;*
- (3) *for every pullback square in C with vertical maps in P and horizontal maps in I , the mixed Beck–Chevalley map $j_{\#}p_* \rightarrow q_*i_{\#}$ is an equivalence.*

Then the lax symmetric monoidal functor $(C^{\times})^{\text{op}} \rightarrow \text{Cat}^{\times}$ corresponding to D_0 admits a unique extension to a lax symmetric monoidal 2-functor

$$D: \text{Span}_2(C, E)_{I, P}^{\otimes} \longrightarrow \text{Cat}^{\times}.$$

Moreover, for any $e \in E$ with factorization $e = p \circ i$ ($i \in I$, $p \in P$), the exceptional pushforward is $e_! \simeq p_ i_{\#}$.*

The above is the Mann-style extension formulation; equivalently, the inclusion $C^{\text{op}} \hookrightarrow \text{Span}_2(C, E)_{I, P}$ is the universal (I, P) -biadjointable functor (left base change for I , right base change for P).

Thus, to construct a six-functor formalism it suffices to verify the three biadjointability conditions: left base change for I , right base change for P , and the mixed Beck–Chevalley equivalence. All coherences come for free from the universal property.

Throughout this subsection, (C, E) denotes a geometric setup and $D: \text{Span}(C, E) \rightarrow \text{Cat}$ a three-functor formalism (or a six-functor formalism, when stated). We write Cat_2 for the 2-category of categories.

Cohomological smoothness

With the formalism in hand, we turn to the geometric properties of a morphism that D can detect, beginning with smoothness. The motivating example is Poincaré duality.

Theorem A.3 (Poincaré duality, classical). *Let $f: X \rightarrow Y$ be a smooth submersion of topological manifolds of relative dimension d (i.e. locally of the form $Y \times \mathbb{R}^d \rightarrow Y$). Then $f_!$ admits a right adjoint $f^!$, and there is a sheaf $\omega_f := f^!(\mathbb{1}_Y) \in D(X, \mathbb{Z})$, locally isomorphic to $\mathbb{Z}[d]$, with*

$$f^! \simeq f^* \otimes \omega_f.$$

The sheaf ω_f is the dualizing complex of f . (For f proper one has instead $f_! \simeq f_$; the two phenomena are independent.)*

A morphism in E which satisfies the analogue of Poincaré duality in D will be called *cohomologically smooth* (or *D-smooth*). Two requirements are immediate:

- There is an isomorphism $f^! \simeq f^* \otimes \omega_f$ for some sheaf ω_f ;
- $\omega_f := f^!(\mathbb{1}_Y)$ is invertible.

But this is not yet enough. Consider a pullback square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Y. \end{array}$$

We expect f' to be D-smooth as well, and the dualizing complexes to be compatible with restriction:

$$g'^* \omega_f \simeq \omega_{f'}.$$

This compatibility is not automatic from pure category theory; it must be imposed.

Definition A.4 (Cohomologically smooth morphism). A morphism $f: X \rightarrow Y$ in E is *cohomologically smooth* (or *D-smooth*) if it satisfies:

(1) The natural transformation

$$f^!(\mathbb{1}_Y) \otimes f^*(-) \longrightarrow f^*(-)$$

is an equivalence; we write $\omega_f := f^!(\mathbb{1}_Y)$ and call it the *dualizing complex* of f .

(2) The dualizing complex ω_f is $\otimes$ -invertible in $D(X)$.

(3) For every pullback square as above with $g \in E$, the morphism f' also satisfies (1)–(2), and the canonical map $g'^* f^!(\mathbb{1}_Y) \rightarrow f'^!(\mathbb{1}_{Y'})$ is an equivalence.

Remark A.5. Cohomological smoothness in the sense of Theorem A.4 is weaker than the geometric notion of a smooth morphism. For instance, in the étale six-functor formalism $X \mapsto \mathrm{Shv}(X_{\text{ét}}, D_{\text{pf}}(\mathbb{Z}))$, every universal homeomorphism between schemes (a map of underlying topological spaces that remains a homeomorphism after pullback) is cohomologically smooth. The étale topology is insensitive to such maps, so they look like isomorphisms at the level of D — and isomorphisms are trivially D -smooth.

The kernel 2-category and Fourier–Mukai transforms

Verifying Theorem A.4 directly is awkward because of the base-change clause (3). The next definitions package D -smoothness into a single 2-categorical statement, after which the base-change clause is automatic.

Fix $Y \in C$, and let $C_E \subseteq C$ be the wide subcategory whose morphisms are those in E . The slice $(C_E)_{/Y}$, paired with the class of *all* morphisms, is itself a geometric setup, and D restricts to a functor

$$D_Y: \mathrm{Span}((C_E)_{/Y}, \mathrm{all})^{\otimes} \rightarrow \mathrm{Span}(C, E)^{\otimes} \xrightarrow{D} \mathrm{Cat}^{\times}.$$

Definition A.6 (Fourier–Mukai transform). Given $X_1, X_2 \in (C_E)_{/Y}$ and $K \in D(X_1 \times_Y X_2)$, the *Fourier–Mukai transform with kernel K* is the functor

$$F_K: D(X_1) \rightarrow D(X_2), \quad A \longmapsto (\pi_2)_!(\pi_1^* A \otimes K).$$

A span $X_1 \xleftarrow{h} Z \xrightarrow{k} X_2$ over Y induces a Fourier–Mukai transform whose kernel is $K = (h, k)_! \mathbb{1}_Z$.

Remark A.7. The terminology is borrowed from the classical analytic transform $T: C^\infty(X) \rightarrow C^\infty(Y)$, $f \mapsto \int_X K(x, y) f(x) dx$, determined by its kernel K . Theorem A.6 categorifies this construction: $\pi_1^* A$ pulls A back to a “function of two variables”, $\otimes K$ multiplies pointwise by the kernel, and $(\pi_2)_!$ is integration along the first variable.

Specializing to $f: X \rightarrow Y$ in E :

- Setting $X_1 = X$, $X_2 = Y$, and $K_1 = \mathbb{1}_X \in D(X \times_Y Y) = D(X)$, the Fourier–Mukai transform is $A \mapsto f_! A$.

- Setting $X_1 = Y$, $X_2 = X$, and $K_2 = \omega_f = f^!(\mathbb{1}_Y) \in D(Y \times_Y X) = D(X)$, the transform is $B \mapsto f^*B \otimes \omega_f$.

Idea A.8. The desired identity $f^! \simeq f^* \otimes \omega_f$ should arise by exhibiting the kernels K_1 and K_2 as an adjoint pair in a 2-category whose objects are schemes over Y and whose 1-morphisms are kernels. Since 2-functors preserve adjunctions, the resulting adjunction is automatically compatible with arbitrary base change $g: Y' \rightarrow Y$, taking care of clause (3) of Theorem A.4 for free.

The invertibility of ω_f is a separate, relatively independent condition; this is what motivates the notion of suaveness below.

Definition A.9 (2-category of kernels). Let $Y \in C$. The 2-category of kernels $\mathbb{K}_{D,Y}$ is the 2-category whose:

- *Objects* are objects of $(C_E)_{/Y}$, i.e., morphisms $X \rightarrow Y$ in E .
- *1-morphism category* from X_1 to X_2 is $\text{Fun}_Y(X_1, X_2) := D(X_1 \times_Y X_2)$.
- *Composition* of $M \in \text{Fun}_Y(X_1, X_2)$ and $N \in \text{Fun}_Y(X_2, X_3)$ is

$$N \circ M := (\pi_{13})_!(\pi_{12}^* M \otimes \pi_{23}^* N) \in D(X_1 \times_Y X_3),$$

where the π_{ij} are the projections from $X_1 \times_Y X_2 \times_Y X_3$.

The identity 1-morphism on X is $\Delta_f!(\mathbb{1}_X) \in D(X \times_Y X)$, where $\Delta_f: X \rightarrow X \times_Y X$ is the relative diagonal.

Conceptually, $\mathbb{K}_{D,Y}$ is obtained from $\text{Span}((C_E)_{/Y}, \text{all})$, which is closed (being a span category over a category with finite limits) and hence self-enriched, by base-changing the self-enrichment along D_Y ; see [Aok26, Definition 2.4].

There is a canonical 2-functor

$$\Psi_{D,Y} = \text{Fun}_Y(Y, -) : \mathbb{K}_{D,Y} \longrightarrow \text{Cat}_2,$$

sending an object X to $D(X)$ and a 1-morphism $M \in \text{Fun}_Y(X_1, X_2)$ to the Fourier–Mukai transform $(\pi_2)_!(\pi_1^*(-) \otimes M)$.

For the reader’s convenience, we record three facts about adjunctions in 2-categories that we use freely (proofs are standard, see [HM24, Appendix D]):

- Any 2-functor $F: \mathbb{C} \rightarrow \mathbb{D}$ preserves adjunctions; in fact, it preserves the entire mate correspondence [HM24, Lemma D.2.7].

- Adjunctions are unique up to a contractible space of choices: any two right adjoints to a given 1-morphism are canonically equivalent.
- A pair (f, g) with candidate unit η and counit ε such that the triangle composites are isomorphisms (not necessarily identities) can be corrected to a strict adjunction by modifying η .

Theorem A.10 (Smoothness via kernels). *A morphism $f: X \rightarrow Y$ in E is D-smooth if and only if*

- (1) *The 1-morphism $\mathbb{1}_X \in \text{Fun}_Y(X, Y) = \text{D}(X)$ is a left adjoint in $\mathbb{K}_{\text{D}, Y}$. Denote its right adjoint by $\omega_f \in \text{Fun}_Y(Y, X) = \text{D}(X)$.*
- (2) *ω_f is $\otimes$ -invertible in $\text{D}(X)$.*

When these hold, $\omega_f \simeq f^!(\mathbb{1}_Y)$.

Proof sketch. $(\Rightarrow)$ Suppose f is D-smooth. By Theorem A.4(1), $f^!(-) \simeq f^*(-) \otimes \omega_f$. Combined with the adjunction $f_! \dashv f^!$ this becomes $f_! \dashv f^*(-) \otimes \omega_f$ at the level of functors. Translating to $\mathbb{K}_{\text{D}, Y}$, the kernel $\mathbb{1}_X$ corresponds to $f_!$ and ω_f to $f^*(-) \otimes \omega_f$, so $\mathbb{1}_X \dashv \omega_f$ in $\mathbb{K}_{\text{D}, Y}$. Invertibility is clause (2).

$(\Leftarrow)$ Apply $\Psi_{\text{D}, Y}$ to the adjunction $\mathbb{1}_X \dashv \omega_f$. Since 2-functors preserve adjunctions, the resulting functors $\Psi_{\text{D}, Y}(\mathbb{1}_X) \simeq f_!$ and $\Psi_{\text{D}, Y}(\omega_f) \simeq f^*(-) \otimes \omega_f$ form an adjunction in Cat_2 . By uniqueness of right adjoints, $f^*(-) \otimes \omega_f \simeq f^!(-)$, which is clause (1) of Theorem A.4; setting $- = \mathbb{1}_Y$ gives $\omega_f \simeq f^!(\mathbb{1}_Y)$. The base-change clause (3) follows because $S \mapsto \mathbb{K}_{\text{D}, S}$ assembles into a 2-functor $\text{Span}(C, E)^\otimes \rightarrow \text{Cat}_2^\times$ [HM24, Theorem 4.2.4], so the adjunction $\mathbb{1}_X \dashv \omega_f$ is preserved by base change. $\square$

Suaveness and primness

Theorem A.10 decomposes cohomological smoothness as “left-adjointness in $\mathbb{K}_{\text{D}, Y}$ ” plus “invertibility of the dualizing complex”. As foreshadowed, dropping the invertibility yields broader classes of objects.

Definition A.11 (Suave and prim objects). Let $f: X \rightarrow Y$ be in E and $A \in \text{D}(X)$.

- (1) A is *f -suave* if, viewed as a 1-morphism $X \rightarrow Y$ in $\mathbb{K}_{\text{D}, Y}$, it is a left adjoint. Its right adjoint is the *f -suave dual* $\text{SD}_f(A) \in \text{D}(X)$. Write $\text{Suave}_f(X) \subseteq \text{D}(X)$ for the full subcategory of f -suave objects.
- (2) A is *f -prim* if, viewed as a 1-morphism $Y \rightarrow X$ in $\mathbb{K}_{\text{D}, Y}$, it is a left adjoint. Its right adjoint is the *f -prim dual* $\text{PD}_f(A) \in \text{D}(X)$. Write $\text{Prim}_f(X) \subseteq \text{D}(X)$ for the full subcategory of f -prim objects.

(Equivalently, via the self-duality $\mathbb{K}_{D,Y} \simeq \mathbb{K}_{D,Y}^{\text{op}}$, A is f -prim iff it is a *right* adjoint 1-morphism $X \rightarrow Y$; this is the form used by Heyer–Mann [HM24].)

Remark A.12 (Origin of the terminology). The terms are due to Scholze (suave) and Hansen (prim); applications typically require D to be a six-functor formalism, although the definition makes sense for three-functor formalisms.

- *Suave (relaxed smoothness)*: when $A = \mathbb{1}_X$ is f -suave, applying $\Psi_{D,Y}$ to the adjunction $\mathbb{1}_X \dashv \text{SD}_f(\mathbb{1}_X)$ gives $f^*(-) \otimes \text{SD}_f(\mathbb{1}_X) \simeq f^!(-)$. More generally, every f -suave A produces a relation between f^* and $f^!$.
- *Prim (primitive properness)*: when $A = \mathbb{1}_X$ is f -prim, the dual statement gives $f_!(\text{PD}_f(\mathbb{1}_X) \otimes -) \simeq f_*(-)$. Primness is the phenomenon that f_* is recovered from $f_!$ by twisting; this is strictly weaker than the proper-base-change identity $f_! \simeq f_*$, so it is “primitive” in this sense.

In a precise sense, A is f -suave (or f -prim) iff it is “finite-dimensional relative to f ”: it admits a reflexive duality.

Proposition A.13 (Reflexive duality). *Let D be a six-functor formalism on (C, E) and $f: X \rightarrow Y$ in E .*

- (1) *If $A \in D(X)$ is f -suave, then $\text{SD}_f(A)$ is also f -suave (with right adjoint A in $\mathbb{K}_{D,Y}$), and there is a natural isomorphism*

$$\text{SD}_f(A) \otimes f^*(-) \simeq \underline{\text{Hom}}(A, f^!(-)): D(Y) \rightarrow D(X).$$

Setting $- = \mathbb{1}_Y$ yields $\text{SD}_f(A) \simeq \underline{\text{Hom}}(A, f^!(\mathbb{1}_Y))$, exhibiting $\text{SD}_f(A)$ as a relative Verdier dual of A . The bidual identity $\text{SD}_f(\text{SD}_f(A)) \simeq A$ then holds, and SD_f defines a contravariant equivalence

$$\text{SD}_f: \text{Suave}_f(X)^{\text{op}} \xrightarrow{\sim} \text{Suave}_f(X).$$

- (2) *Dually, if $A \in D(X)$ is f -prim, then $\text{PD}_f(A)$ is also f -prim, with*

$$f_!(\text{PD}_f(A) \otimes -) \simeq f_* \underline{\text{Hom}}(A, -): D(X) \rightarrow D(Y),$$

and the bidual $\text{PD}_f(\text{PD}_f(A)) \simeq A$ holds. Consequently PD_f defines a contravariant equivalence

$$\text{PD}_f: \text{Prim}_f(X)^{\text{op}} \xrightarrow{\sim} \text{Prim}_f(X).$$

Suave morphisms and prim morphisms

Specializing Theorem A.11 to $A = \mathbb{1}_X$ yields classes of *morphisms*.

Definition A.14 (Suave and prim morphisms). Let $f: X \rightarrow Y$ be in E .

- (1) f is *D-suave* if $\mathbb{1}_X \in \text{Suave}_f(X)$. As before we write $\omega_f := \text{SD}_f(\mathbb{1}_X) \in \text{D}(X)$ for its *dualizing complex*.
- (2) f is *D-prim* if $\mathbb{1}_X \in \text{Prim}_f(X)$. We then write $\delta_f := \text{PD}_f(\mathbb{1}_X) \in \text{D}(X)$ for the *co-dualizing complex* of f .

Remark A.15. Combining Theorem A.10 with Theorem A.14 gives

$$f \text{ is D-smooth} \iff f \text{ is D-suave and } \omega_f \text{ is invertible.}$$

By Theorem A.13, the dualizing complex is the relative Verdier dual $\omega_f \simeq f^!(\mathbb{1}_Y)$. The co-dualizing complex is computed directly in the kernel 2-category (Theorems A.9 and A.11) as the right adjoint of the identity kernel,

$$\delta_f \simeq \pi_{2*} \Delta_f! \mathbb{1}_X,$$

where $\pi_2: X \times_Y X \rightarrow X$ is the projection onto the second factor.

For D-suave f , the formula $\omega_f \otimes f^*(-) \xrightarrow{\sim} f^!(-)$ is equivalent (by adjunction) to $f^*(-) \xrightarrow{\sim} \underline{\text{Hom}}(\omega_f, f^!(-))$. Plugging in $\mathbb{1}_Y$ yields $\mathbb{1}_X \xrightarrow{\sim} \underline{\text{Hom}}(\omega_f, \omega_f)$, and for dualizable ω_f this gives $\mathbb{1}_X \simeq \omega_f^\vee \otimes \omega_f$. We record this as an observation.

Observation A.16. For a D-suave morphism f , the dualizing complex ω_f is $\otimes$ -invertible if and only if it is dualizable. Consequently the verification of D-smoothness reduces to checking suaveness and dualizability of ω_f .

We close with the base-change identities for suave and prim morphisms; they will guide the definitions of cohomologically étale and proper morphisms in the next subsection.

Lemma A.17 (Base change for suave and prim morphisms, [HM24, Lemma 4.5.13]). *Let D be a six-functor formalism on (C, E) and consider a pullback square*

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

in C_E .

(1) If f is D-suave, the natural transformations

$$g'_* f'^* \simeq f^* g_*, \quad g'_! f'^! \simeq f^! g_!, \quad f'^* g'^! \simeq g'^! f'^*, \quad g'^* f'^! \simeq f'^! g'^*$$

are all isomorphisms.

(2) If f is D-prim, the natural transformations

$$g^* f_* \simeq f'_* g'^*, \quad g^! f_! \simeq f'_! g'^!, \quad f_! g'_* \simeq g_* f'_!, \quad f_* g'_! \simeq g'_! f_*$$

are all isomorphisms.

Cohomological étaleness and properness

We now ask: under what conditions are there genuine isomorphisms $f^! \simeq f^*$ and $f_! \simeq f_*$, rather than mere twists by ω_f or δ_f ?

A candidate map $f^! \rightarrow f^*$ is constructed using the diagonal. Consider

$$\begin{array}{ccccc} X & & \xrightarrow{\text{id}_X} & & X \\ & \searrow \Delta_f & & \searrow \downarrow & \\ & X \times_Y X & \xrightarrow{\pi_2} & & X \\ & \swarrow \text{id}_X & \downarrow \pi_1 & \lrcorner & \downarrow f \\ & X & \xrightarrow{f} & & Y \end{array}$$

We have $f\pi_1 = f\pi_2$ and $\text{id}_X = \pi_2\Delta_f$. Suppose inductively that Δ_f satisfies $\Delta_f^! \simeq \Delta_f^*$. Then

$$\begin{aligned} f^! &\simeq \text{id}_X^* f^! \simeq \Delta_f^* \pi_2^* f^! \xrightarrow{\sim} \Delta_f^! \pi_2^* f^! \\ &\rightarrow \Delta_f^! \pi_1^! f^* \simeq (\pi_1 \Delta_f)^! f^* \simeq f^*, \end{aligned}$$

where $\pi_2^* f^! \rightarrow \pi_1^! f^*$ is induced (via right adjoints) from the base-change isomorphism $\pi_{1!} \pi_2^* \simeq f^* f_!$. By Theorem A.17, when f is D-suave the latter is an isomorphism, and so $f^! \simeq f^*$.

A dual construction gives $f_! \rightarrow f_*$, an isomorphism when Δ_f satisfies $\Delta_{f!} \simeq \Delta_{f*}$ and f is D-prim. To make the recursion well-founded we impose a truncatedness hypothesis: the recursion terminates at -2 -truncated morphisms (i.e., isomorphisms), where the conclusion is trivial.

Definition A.18 (Cohomologically étale and proper morphisms). Let D be a six-functor formalism on (C, E) , and let $f: X \rightarrow Y$ be a truncated morphism in E (i.e., n -truncated for some $n \in \mathbb{N}$).

- f is D-étale (or cohomologically étale) if f is D-suave and Δ_f is either D-étale or an isomorphism.

- f is *D-proper* (or *cohomologically proper*) if f is D-prim and Δ_f is either D-proper or an isomorphism.

These definitions are well-founded because n -truncated morphisms have $(n - 1)$ -truncated diagonals.

Proposition A.19 (Characterization of étaleness and properness). *Let D be a six-functor formalism on (C, E) and $f: X \rightarrow Y$ in E .*

(1) *Suppose Δ_f is D-étale. There is a natural map $f^! \rightarrow f^*$, and the following are equivalent:*

- (a) *f is D-étale;*
- (b) *$f^! \mathbb{1}_Y \simeq f^* \mathbb{1}_Y$ in $D(X)$;*
- (c) *$f^! \xrightarrow{\sim} f^*$ as functors.*

(2) *Suppose Δ_f is D-proper. There is a natural map $f_! \rightarrow f_*$, and the following are equivalent:*

- (a') *f is D-proper;*
- (b') *$f_! \mathbb{1}_X \simeq f_* \mathbb{1}_X$ in $D(Y)$;*
- (c') *$f_! \xrightarrow{\sim} f_*$ as functors.*

This completes the foundational vocabulary — cohomological smoothness, suaveness, primness, étaleness, properness — whose translation under the transmutation $[-]_D$ is the subject of the dictionary in the main text.

References

-
- [Aok26] K. Aoki. “Berkovich 2-motives and normed ring stacks” (2026). arXiv: [2603.01877](#).
 - [Bei16] A. Beilinson. “Constructible sheaves are holonomic”. *Selecta Math. (N.S.)* 22.4 (2016), pp. 1797–1819.
 - [BFN10] D. Ben-Zvi, J. Francis, and D. Nadler. “Integral transforms and Drinfeld centers in derived algebraic geometry”. *J. Amer. Math. Soc.* 23.4 (2010), pp. 909–966.
 - [Bha22] B. Bhatt. “Prismatic F -gauges”. Lecture notes, IAS, course MAT 549, Fall 2022. 2022.
 - [CD19] D.-C. Cisinski and F. Déglise. *Triangulated Categories of Mixed Motives*. Springer Monographs in Mathematics. Springer, 2019.
 - [CLL25] B. Cnossen, T. Lenz, and S. Linskens. “Universality of span 2-categories and the construction of 6-functor formalisms” (2025). arXiv: [2505.19192](#).
 - [Gai25] D. Gaitsgory. “Local and global Langlands conjecture(s) over function fields” (2025). ICM 2026 plenary report. arXiv: [2509.24902](#).

- [Gal21] M. Gallauer. *An introduction to six-functor formalisms*. Lecture notes. 2021. arXiv: [2112.10456](#).
- [GRV25] D. Gaiitsgory, N. Rozenblyum, and Y. Varshavsky. “Applications of (higher) categorical trace I: the definition of AGCat” (2025). arXiv: [2512.21726](#).
- [HM24] C. Heyer and L. Mann. “6-Functor Formalisms and Smooth Representations” (2024). arXiv: [2410.13038](#).
- [Kes25] Y. Kesting. *Categorical Künneth formulas for analytic stacks*. 2025. arXiv: [2507.08566](#).
- [KS90] M. Kashiwara and P. Schapira. *Sheaves on Manifolds*. Vol. 292. Grundlehren der mathematischen Wissenschaften. Springer-Verlag, Berlin, 1990.
- [LZ24] Y. Liu and W. Zheng. “Enhanced six operations and base change theorem for higher Artin stacks” (2024). arXiv: [1211.5948](#).
- [Man22] L. Mann. “A p -adic 6-functor formalism in rigid-analytic geometry”. PhD thesis. Universität Bonn, 2022. arXiv: [2206.02022](#).
- [RZ25] D. Reutter and M. Zetto. “Enriched ∞ -categories as marked module categories” (2025). arXiv: [2501.07697](#).
- [Sch25] P. Scholze. “Six-Functor Formalisms” (2025). arXiv: [2510.26269](#).
- [Sch26] P. Scholze. “Geometry and Higher Category Theory”. Lecture notes (with G. Stefanich), Universität Bonn, Winter 2025/26. 2026.
- [Ste25] G. Stefanich. “Categorification of sheaf theory” (2025). arXiv: [2511.09553](#).