

Calculating Subset Weighted Analysis Using PROC SURVEYFREQ and GENMOD

Jessica J. Hale, MS¹, David M. Thompson, PhD², and Paul M. Darden, MD¹

¹ Department of Pediatrics, School of Medicine;

² Department of Biostatistics and Epidemiology, School of Public Health; University of Oklahoma Health Sciences Center, Oklahoma City, OK, 73104

Why Subset?

- May not need to analyze all the observations collected for analysis
- Allows analyst to use a smaller, condensed dataset that contains only the necessary observations
- Less code to write

National Immunization Survey (NIS)-Teen

- Center for Disease Control (CDC) began collecting the NIS-Teen survey in 2008
- Immunization related data collected from parents and providers
- Examples and results shown are from the NIS-Teen 2010 data
 - Outcome of interest is being up-to-date (UTD) for the Tetanus (TET) vaccine for females only
 - Parent collected information used
 - Weight variable is the provider weight variable (PROVWT)
 - Clusters are represented by the unique identifier, SEQNUMT, which is assigned to each teen that participated in the survey
 - Estimation area or stratum are identified by the ESTIAPT variable
 - Domain variable is sex

Methods

- Three SAS® procedure examples, SURVEYFREQ, FREQ and GENMOD are displayed
- Domain variables can be used in the survey procedures available in SAS® for subset analysis
- Subset weighted analysis can be performed outside of the survey procedures using standardized weights or specifying the survey parameters when available
- Individual standardized weights (ISW) (Program 6: SAS® Code)

ISW Formula:

$$ISW = (w_{PROVWT} / \sum w_{PROVWT}) * n$$

Step 1: Calculate the sum of PROVWT

Step 2: Divide each respondent’s PROVWT value by the sum of PROVWT

Step 3: Multiply by the total sample size of respondents with a PROVWT value

Results

Results from SAS® programs 1 to 5

SAS® Procedure	Program	Sample Subset	Survey Parameters Used in Analysis	Format of Parameter Estimate (Est) from Procedure	Est (percent)	Standard Error (percent)
SURVEYFREQ	1	All records	ESTIAPT, SEQNUMT, PROVWT	Percentage	78.4456	0.9346
	2	Females only	PROVWT	Percentage	78.4456	0.9374
FREQ	3	Females only	PROVWT	Proportion	78.4456	0.0149
	4	Females only	ISW	Proportion	78.4456	0.4881
GENMOD	5	Females only	ESTIAPT, SEQNUMT, PROVWT	Proportion	78.4456	0.9373

SAS® Programming

Program 1: Designating a domain variable in the TABLE statement for calculating subset specific analysis using PROC SURVEYFREQ

```
proc surveyfreq data=nis10;  
tables sex*r_tet_not_utd/col  
row;  
strata estiapt10;  
cluster seqnumt;  
weight provwt;  
run;
```

Program 2: Designating a subset using the WHERE statement and calculating subset specific analysis using PROC SURVEYFREQ and only the weight variable; ignoring other survey parameters.

```
proc surveyfreq data=nis10;  
tables r_tet_not_utd/row cl;  
weight provwt;  
where sex = 2;  
run;
```

Program 3: Designating a subset using the WHERE statement and calculating subset specific analysis using PROC FREQ and only the weight variable while ignoring other survey parameters.

```
proc freq data=nis10;  
exact binomial;  
tables r_tet_not_utd;  
weight provwt;  
where sex = 2;  
run;
```

Program 4: Using Individual Provider Weight in PROC FREQ to Calculate Subset Weighted Proportions

```
proc freq data=nis10_new;  
exact binomial;  
tables r_tet_not_utd;  
weight new_provwt;  
where sex = 2;  
run;
```

Program 5: Using PROC GENMOD to Calculate Subset Weighted Proportions

```
proc genmod data=nis10;  
class seqnumt estiapt10;  
model r_tet_not_utd = /  
dist=binomial link=identity;  
weight provwt;  
repeated subject  
=seqnumt(estiapt10);  
where sex = 2;  
run;
```

Program 6: Calculating the ISW

```
*Step 1;  
proc means data=nis10 sum n;  
var provwt;  
output out = provwt_2 sum =  
sum_provwt n = n_provwt;  
run;  
*Steps 2 and 3;  
data nis_10_new;  
if _n_ = 1 then set provwt_2;  
set nis_10;  
new_providerwt_SAS =  
((provwt/sum_provwt)*n_provwt);  
run;
```

Discussion

- Researchers and analysts need to be aware that various methods produce variable estimates and standard errors
- SAS® survey procedures use all survey parameters, which can be difficult if the focus is on analyzing multiple subsets within a dataset
- The SAS® manual advises against using the WHERE statement to perform weighted analysis of subsets of survey data
- The examples shown analyze a subset of survey data while calculating similar or more conservative SE
 - When a SAS® procedure does not permit specifying the stratum or cluster variables, the ISW is an alternative method to performing the analysis
 - Using the original weights without the capability of specifying the stratum and cluster variables yields very small standard errors

Conclusions

- Multiple options are available for calculating weighted subset analysis
 - PROC SURVEYFREQ
 - PROC FREQ
 - PROC GENMOD
- Estimates across the various methods remain consistent, but the standard errors vary depending on the method used
- Stratum and cluster variables for a survey can be applied in PROC GENMOD using the REPEATED statement and SUBJECT option
- Ability to utilize the capabilities of GENMOD , such as GEE, allows researchers and analysts the opportunity to discover potential new findings previously unavailable for survey data

Caution

- The methods presented here apply to the NIS-Teen survey, but may produce different results on other complex survey designs
- In order to determine if these methods apply to a complex design survey other than the NIS-Teen, tests similar to the various examples shown could be used

Acknowledgements

SAS® and all other SAS® Institute Inc. product or service names are registered trademarks or trademarks of SAS® Institute Inc. in the USA and other countries. ® indicates USA registration. Other brand and product names are trademarks of their respective companies

Contact Us

Jessica J. Hale
David M. Thompson
Paul M. Darden

jessica-j-hale@ouhsc.edu
david-thompson@ouhsc.edu
paul-darden@ouhsc.edu

