
Estimated Precision (PPV) by Gene Barcode
1.00
0.95
0.90
0.85
0.80
Estimated PPV: Median with 0.95 CI
Gene barcodes by lower 0.025 quantile of PPV confidence inte


Simulated vs Run Counts
1000
100
10
Spot count
Barcodes sorted by run count
Count type
Gene (sim)
Gene (run)
Blank (sim)
Blank (run)


MCMCSA step diagnostics
MSE (log)
5
4
3
2
2.5
5.0
7.5
10.0
12.5
step


MCMCSA Flip Rates
Flip Rate
0.15
0.10
0.05
2.5
5.0
7.5
10.0
12.5
step
Type
0 to 1
1 to 0


MCMCSA bit−noise correlation parameters
0.20
0.15
0.10
0.05
0.00
Value
2.5
5.0
7.5
10.0
12.5
step
Type
bias
sd


Estimated Precision (PPV) by Gene Barcode
1.00
0.95
0.90
0.85
0.80
Estimated PPV: Median with 0.95 CI
Gene barcodes by lower 0.025 quantile of PPV confidence inte


Simulated vs Run Counts
1000
100
10
1
Spot count
Barcodes sorted by run count
Count type
Gene (sim)
Gene (run)
Blank (sim)
Blank (run)


MCMCSA step diagnostics
MSE (log)
5
4
3
2.5
5.0
7.5
10.0
12.5
step


MCMCSA Flip Rates
Flip Rate
0.15
0.10
0.05
2.5
5.0
7.5
10.0
12.5
step
Type
0 to 1
1 to 0


MCMCSA bit−noise correlation parameters
0.20
0.15
0.10
0.05
0.00
Value
2.5
5.0
7.5
10.0
12.5
step
Type
bias
sd


Estimated Precision (PPV) by Gene Barcode
1.00
0.95
0.90
0.85
0.80
Estimated PPV: Median with 0.95 CI
Gene barcodes by lower 0.025 quantile of PPV confidence inte


Simulated vs Run Counts
1000
100
10
1
Spot count
Barcodes sorted by run count
Count type
Gene (sim)
Gene (run)
Blank (sim)
Blank (run)


MCMCSA step diagnostics
MSE (log)
5
4
3
2
1
2.5
5.0
7.5
10.0
12.5
step


MCMCSA Flip Rates
0.16
0.12
0.08
0.04
Flip Rate
2.5
5.0
7.5
10.0
12.5
step
Type
0 to 1
1 to 0


MCMCSA bit−noise correlation parameters
0.20
0.15
0.10
0.05
0.00
Value
2.5
5.0
7.5
10.0
12.5
step
Type
bias
sd