

ALGEBRA Volumen I y II



Asesoría Bimestral







Reduce:
$$G = (-3) + (-4) - (-4) - (-10) - (18) - (-20)$$

Resolución

Reduciendo:

$$G = -3 - 4 + 4 + 10 - 18 + 20$$

Separando:

$$G = (+ 4 + 10 + 20) - (3 + 4 + 18)$$
 $G = (+ 34) - (25)$
 $G = +34 - 25$

Rpta:

9





Reduce

$$M = \left(\frac{\frac{14}{3} - \frac{7}{3} + \frac{5}{3}}{1 + \frac{1}{3}}\right) + 3$$

$$M = \left(\frac{\frac{12}{3}}{1 + \frac{1}{3}}\right) + 3 = \left(\frac{\frac{12}{3}}{\frac{4}{3}}\right) + 3$$



$$M = \frac{12}{4} + 3 = 3 + 3 = 6$$
 Rpta:



Determine el valor de x en:

$$\frac{2x-1}{4} - \frac{5x}{8} = \frac{x-2}{2}$$

mcm(4; 8; 2) = 8

Resolución

$$8\left(\frac{2x-1}{4}\right) - 8\frac{(5x)}{8} = 8\left(\frac{x-2}{2}\right)$$

$$2(2x-1) - 5x = 4(x-2)$$

$$4x - 2 - 5x = 4x - 8$$

$$-x - 2 = 4x - 8$$

$$-2 + 8 = 4x + x$$

$$6 = 5x$$

$$6/5 = x$$



Rpta:

6/5



Efectúe
$$R = \left(\frac{1}{5}\right)^{-2} + \left(\frac{1}{3}\right)^{-4} + \left(\frac{1}{5}\right)^{-3}$$

$$R = (5)^2 + (3)^4 + (5)^3$$

$$R = 25 + 81 + 125$$





Reduce
$$A = \frac{2^{a+3}}{2} + \frac{5^{a-4}}{5^{a-5}} + \frac{3^{a-2}}{3^{a-4}}$$

$$A = 2^{a+3-(a)} + 5^{a-4-(a-5)} + 3^{a-2-(a-4)}$$

$$A = 2^{\hat{a}+3-\hat{a}} + 5^{\hat{a}-4-\hat{a}+5} + 3^{\hat{a}-2-\hat{a}+4}$$

$$A=2^3+5^1+3^2$$

$$A = 8 + 5 + 9$$





Calcula el exponente final de x en

$$B = \sqrt[2]{x^7 \cdot \sqrt[4]{x^3 \cdot \sqrt[2]{x}}}; x \neq 0$$

Resolución

Multiplicación de índices

$$B = \sqrt[16]{x^{65}}$$

$$B = \chi_{16}^{65}$$

Rpta

El exponente es
$$\frac{65}{16}$$





Reduce la expresión:

$$M = \sqrt[2]{x^1 \cdot \sqrt[2]{x^1 \cdot \sqrt[3]{x}}} \cdot \sqrt[8]{x} \quad ; \quad x \neq 0$$

$$\frac{1}{por} \int_{más}^{1} \frac{por}{más} = 7$$

Multiplicación de índices
$$M = \sqrt[8]{x^7} \cdot \sqrt[8]{x}$$

$$M = \chi^{\frac{1}{8}} = \chi$$



Rpta
$$M = x$$



Determine el valor de x en:

$$\frac{2x-5}{3} - \frac{x-3}{6} = \frac{3x-2}{2} \quad mcm(3;6;2) = 6$$

$$6\left(\frac{2x-5}{3}\right) - 6\frac{(x-3)}{6} = 6\left(\frac{3x-2}{2}\right) \quad 2(2x-5) - (x-3) = 3(3x-2) \\ 4x - 10 - x + 3 = 9x - 6$$

$$2(2x - 5) - (x - 3) = 3(3x - 2)$$
$$4x - 10 - x + 3 = 9x - 6$$



$$3x - 7 = 9x - 6$$
$$-1 = 6x$$

Rpta:
$$x = -1/6$$



Determine el valor de 100AB, si:

A=
$$\left(\frac{4}{10}\right) \left(\frac{-15}{16}\right) \left(\frac{-2}{25}\right)$$
 y B= $\left(\frac{25}{32}\right) \div \left(\frac{35}{14}\right) + 1$

$$A = \begin{pmatrix} \frac{4}{10} \end{pmatrix} \begin{pmatrix} \frac{-15}{16} \end{pmatrix} \begin{pmatrix} \frac{-2}{25} \\ \frac{1}{25} \end{pmatrix} = \frac{3}{100}$$

$$B = \begin{pmatrix} \frac{25}{32} \end{pmatrix} \begin{pmatrix} \frac{14}{35} \end{pmatrix} + 1 = \begin{pmatrix} \frac{5}{16} \end{pmatrix} + 1 = \begin{pmatrix} \frac{21}{16} \end{pmatrix}$$

100AB=
$$100 \left(\frac{3}{100}\right) \left(\frac{21}{16}\right)$$
100AB= $\left(\frac{63}{16}\right)$



Efectúe
$$R = 4^{1^3} + 1^{5^{4^3}} + 2^{3^{1^{35}}} - 5^{2^{0^{351}}}$$

$$R = (4)^{1} + (1)^{5^{64}} + (2)^{3} - (5)^{2^{0}}$$

$$R + 1 + 8 - 5$$

