

# 1 Introduction

We will review the definition of a Springer fiber and define, for a nilpotent  $Y \in \mathfrak{gl}_m$ , the Springer fiber at the  $n$ -Slodowy slice at  $Y$ . For every  $n$  and every nilpotent  $Y \in \mathfrak{gl}_m$ , we will find the irreducible components of the Springer fiber at the  $n$ -Slodowy slice at  $Y$ . Finally, we will use our results about Springer fibers at  $n$ -Slodowy slices to find the irreducible components of some other variety (which probably needs a name), and show that they all have the same dimension.

## 2 Springer fibers

Let  $G \subseteq \mathrm{GL}_m(\mathbb{C})$  be a connected semisimple Lie group, and let  $\mathfrak{g} \subseteq \mathfrak{gl}_m(\mathbb{C})$  be its Lie algebra. Let  $\mathcal{N} \subseteq \mathfrak{g}$  be the subset consisting of nilpotent elements. Let  $\mathcal{B}$  be the variety of Borel subalgebras of  $\mathfrak{g}$ . Let  $\tilde{\mathcal{N}} = \{(\mathfrak{b}, n) : n \in \mathfrak{b}\} \subseteq \mathcal{B} \times \mathcal{N}$ . Let  $\pi : \tilde{\mathcal{N}} \rightarrow \mathcal{N}$  be the projection onto the second coordinate. For  $n \in \mathcal{N}$ , we call  $\pi^{-1}(n)$  the *Springer fiber at  $n$* .

We mention some results about Springer fibers, which we will use later in this paper. TODO: mention them.

## 3 Slodowy slice

A basis for  $\mathfrak{sl}_2(\mathbb{C})$  is

$$e' := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, h' := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f' := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Given a Lie algebra  $\mathfrak{g}$ , and a homomorphism  $\phi : \mathfrak{sl}_2 \rightarrow \mathfrak{g}$  sending  $(e', h', f')$  to  $(e, h, f)$ , we say that  $(e, h, f)$  is an  *$\mathfrak{sl}_2$ -triple*. Observe that  $e, f$  must be nilpotent, and  $h$  must be Cartan (???). If  $\mathfrak{g}$  is semisimple, then given any nilpotent  $e \in \mathfrak{g}$ , the Jacobson-Morozov theorem [1, 3.7.1] says that there exist  $h, f \in \mathfrak{g}$  such that  $(e, h, f)$  is an  $\mathfrak{sl}_2$ -triple.

Given  $(e, h, f)$ , we define the *Slodowy slice at  $e$*  as  $\mathcal{S}_e := e + \ker \mathrm{ad}_f$ . By the Jacobson-Morozov theorem, we can always find a Slodowy slice at any nilpotent  $e \in \mathfrak{g}$ , when  $\mathfrak{g}$  is semisimple.

## 4 Finding $\mathfrak{sl}_2$ -triples $(E, H, F)$ in $\mathfrak{gl}_{m+n}$ with a particular $E$ .

Let

$$e = \begin{pmatrix} 0 & & & \\ 1 & 0 & & \\ & 1 & \ddots & \\ & & \ddots & 0 \\ & & & 1 & 0 \end{pmatrix} \in \mathfrak{gl}_n,$$

and let  $E = \begin{pmatrix} 0 & 0 \\ 0 & e \end{pmatrix} \in \mathfrak{gl}_{m+n}$ . We will show that there is exactly one  $\mathfrak{sl}_2$ -triple  $(E, H, F)$ , and we will find what it looks like. First we solve the case  $m = 0$  (so  $E = e$ ), and then we use this to solve the case of arbitrary  $m$ .

**Lemma 4.1.** *There is exactly one way to choose  $h, f \in \mathfrak{gl}_n$  so that  $(e, h, f)$  is an  $\mathfrak{sl}_2$ -triple.*

*Proof.* Note that  $[h', e'] = 2e'$ , and  $[e', f'] = h'$ , and  $[h', f'] = -2f'$ . Thus  $e, h, f$  must obey the same relations. In particular,  $he - eh = 2e$ . The matrix  $eh$  is  $h$  shifted down one, and  $he$  is  $h$  shifted left one. Thus,  $2e_{ij} = (he - eh)_{ij} = h_{i,j+1} - h_{i-1,j}$ . We can use this to show that  $h_{ij} = 0$  when  $i \neq j$ . Then we can use it to show that  $h_{ii} = h_{i-1,i-1} + 2$ , so that  $h_{ii} = h_{11} + 2(i-1)$ .

Similarly, from  $[e, f] = h$  we get that  $h_{ij} = (ef - fe)_{ij} = f_{i-1,j} - f_{i,j+1}$ . We can use this to show that  $f_{ij} = 0$  when  $j \neq i+1$ . Then we can use it to show that  $f_{i,i+1} = f_{i+1,i+2} + h_{i+1,i+1}$ , that  $f_{1,2} = -h_{1,1}$ , and that  $f_{n-1,n} = h_{n,n}$ . From the first equation we find that

$$-h_{1,1} = f_{1,2} = f_{n-1,n} + \sum_{i=1}^{n-1} h_{ii} = \sum_{i=1}^n h_{ii} \implies \sum_i h_{ii} = 0.$$

Remark: this is just the statement that  $h \in \mathfrak{sl}_n$ ; in other words, we will see that every choice of  $\mathfrak{sl}_2$ -triple in  $\mathfrak{gl}_{m+n}$  is also an  $\mathfrak{sl}_2$ -triple in  $\mathfrak{sl}_{m+n}$ . This shows that  $h_{11} = n-1, h_{22} = n-3, \dots, h_{nn} = 1-n$ . So we have determined

$h$ ; it is

$$h = \begin{pmatrix} n-1 & & & & \\ & n-3 & & & \\ & & \ddots & & \\ & & & 3-n & \\ & & & & 1-n \end{pmatrix}.$$

Now, we can use our expression for  $f$  in terms of  $h$  to obtain

$$f = \begin{pmatrix} 0 & 1-n & & & \\ & 0 & (1-n) + (3-n) & & \\ & & \ddots & \ddots & \\ & & & 0 & (1-n) + \cdots + (n-1) \\ & & & & 0 \end{pmatrix} =$$

$$\begin{pmatrix} 0 & 1(1-n) & & & \\ & 0 & 2(2-n) & & \\ & & 0 & (n-2)(-2) & \\ & & & 0 & (n-1)(-1) \\ & & & & 0 \end{pmatrix}.$$

□

**Lemma 4.2.** *There is exactly one way to choose  $H, F \in \mathfrak{gl}_{m+n}$  so that  $(E, H, F)$  is an  $\mathfrak{sl}_2$ -triple.*

*Proof.* Suppose we have  $H, F$  so that  $(E, H, F)$  is an  $\mathfrak{sl}_2$ -triple. Writing  $F =: \begin{pmatrix} F_{11} & F_{12} \\ F_{21} & f \end{pmatrix}$ , and similarly for  $H$ , we have

$$2E = [H, E] = \begin{pmatrix} 0 & H_{12}e \\ 0 & he \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ eH_{21} & eh \end{pmatrix},$$

$$H = [E, F] = \begin{pmatrix} 0 & 0 \\ eF_{21} & ef \end{pmatrix} - \begin{pmatrix} 0 & F_{12}e \\ 0 & fe \end{pmatrix}.$$

From these we observe that  $(e, h, f)$  must also be an  $\mathfrak{sl}_2$ -triple, so  $h, f$  must be as in Lemma 4.1. We also see that  $H_{11} = 0$ . Recalling that left multiplication by  $e$  is a down-shift, and right multiplication is a left-shift, we see that  $H_{12}$

is all zeroes except for the leftmost column, and  $H_{21}$  is all zeroes except for the bottom row. Then our final constraint is that

$$-2F = [H, F] =$$

$$\begin{pmatrix} H_{12}F_{21} & H_{12}F_{22} \\ H_{21}F_{11} + H_{22}F_{21} & H_{21}F_{12} + H_{22}F_{22} \end{pmatrix} - \begin{pmatrix} F_{12}H_{21} & F_{11}H_{12} + F_{12}H_{22} \\ F_{22}H_{21} & F_{21}H_{12} + F_{22}H_{22} \end{pmatrix}.$$

Now  $H_{12} = F_{12}e$ , and  $H_{21} = eF_{21}$ , from the equation  $H = [E, F]$ . Substituting in the equation above then,

$$\begin{aligned} -2F &= \begin{pmatrix} F_{12}eF_{21} & F_{12}ef \\ eF_{21}F_{11} + hF_{21} & eF_{21}F_{12} + hf \end{pmatrix} - \begin{pmatrix} F_{12}eF_{21} & F_{11}F_{12}e + F_{12}h \\ feF_{21} & F_{21}F_{12}e + fh \end{pmatrix} = \\ &= \begin{pmatrix} 0 & F_{12}(fe + h) \\ eF_{21}F_{11} + hF_{21} & eF_{21}F_{12} + hf \end{pmatrix} - \begin{pmatrix} 0 & F_{11}F_{12}e + F_{12}h \\ (ef - h)F_{21} & F_{21}F_{12}e + fh \end{pmatrix} = \\ &= \begin{pmatrix} 0 & F_{12}fe \\ eF_{21}F_{11} & eF_{21}F_{12} + hf \end{pmatrix} - \begin{pmatrix} 0 & F_{11}F_{12}e \\ efF_{21} & F_{21}F_{12}e + fh \end{pmatrix}. \end{aligned}$$

Now we see that  $F_{11} = 0$ , and consequently that  $F_{12} = F_{21} = 0$  as well. This shows that  $H_{12} = H_{21} = 0$ . We conclude that  $H$  and  $F$  just have  $h$  and  $f$  in their bottom-right corners, respectively.  $\square$

## 5 Finding the Slodowy slices with the same $E$

First we find  $\ker \text{ad}_f$ . We have  $(fX)_{ij} = i(n-i)A_{i+1,j}$ , and  $(Xf)_{ij} = (j-1)(n-(j-1))A_{i,j-1}$ . So, for all  $i, j \in \{1, \dots, n\}$ , we have

$$i(n-i)A_{i+1,j} = (j-1)(n-(j-1))A_{i,j-1}.$$

Taking  $j = 1$ , we find that  $A_{i,1} = 0$  for  $i \geq 2$ . Then, taking  $j > 1$ , we find that for  $i, j \in \{1, \dots, n-1\}$ ,

$$A_{i+1,j+1} = \frac{(j-1)(n-(j-1))}{i(n-i)}A_{ij}.$$

So,  $\ker \text{ad}_f$  is the set of matrices which are upper triangular and satisfy the above condition.

To find the Slodowy slice associated to the previous  $\mathfrak{sl}_2$ -triple  $(E, H, F)$ , we just need to find  $\ker \operatorname{ad}_F$ . We have

$$[F, X] = \begin{pmatrix} 0 & 0 \\ fX_{21} & fX_{22} \end{pmatrix} - \begin{pmatrix} 0 & X_{12}f \\ 0 & X_{22}f \end{pmatrix}.$$

Thus,  $X_{21}$  must be all zeroes except for the first row, and  $X_{12}$  must be all zeroes except for the last column, and  $X_{22} \in \ker \operatorname{ad}_f$ . There is no restriction on  $X_{11}$ . This describes  $\ker \operatorname{ad}_F$ .

For  $X \in \mathcal{S}_{m,n}$ , define  $u(X) := X_{11}$ .

### 5.1 Finding $\tilde{\mathcal{N}}_{m,n}$

Let  $\mathcal{N}_m \subseteq \mathfrak{gl}_m$  be the nilpotent elements. Let  $\mathcal{S}'_{m,n}$  be the set of  $X \in \mathcal{S}_{m,n}$  such that both  $X$  and  $u(X)$  are nilpotent. Let  $\tilde{\mathcal{N}}_{m,n} = \{(\mathfrak{b}, X) : X \in \mathfrak{b}\} \subseteq \mathcal{B}_{m+n} \times \mathcal{S}'_{m,n}$ . Define  $\pi_{m,n} : \tilde{\mathcal{N}}_{m,n} \rightarrow \mathcal{N}_m$  by  $(\mathfrak{b}, X) \mapsto X_{11}$ . For  $Y \in \mathfrak{gl}_m$ , we call  $\pi_{m,n}^{-1}(Y)$  the *Springer fiber at the  $n$ -Slodowy slice at  $Y$* .

**Lemma 5.1.** *Let  $J$  be a jordan block with zeroes along the diagonal, and let  $A$  be upper triangular and nonzero. Then  $J + A$  is not nilpotent.*

*Proof.* It is straightforward to show by induction that if  $v_i = 0$  for  $i < j$ , and  $v_j \neq 0$ , then  $((J + A)^k v_j)_{j+k} = v_j$ . Let  $i$  be such that  $Ae_i \neq 0$ . Then  $(J + A)^{i-1}e_1$  has nonzero  $e_i$ -component. Then  $(J + A)^i e_1$  has some nonzero  $e_{i'}$ -component for some  $i' \leq i$ . Then  $(J + A)^{i+(n-i')}e_1$  has some nonzero  $e_n$ -component. And  $i + (n - i') \geq n$ , so we're done.  $\square$

**Lemma 5.2.** *Let  $X \in \mathfrak{gl}_m$ , and let*

$$Y = \begin{pmatrix} y_{11} & y_{12} & y_{13} & \cdots & y_{1,n-1} & y_{1n} \\ d_1 & y_{22} & y_{23} & \cdots & y_{2,n-1} & y_{2n} \\ & d_2 & y_{33} & \cdots & y_{3,n-1} & y_{3n} \\ & & \ddots & \cdots & \vdots & \vdots \\ & & & d_{n-2} & y_{n-1,n-1} & y_{n-1,n} \\ & & & & d_{n-1} & y_{nn} \end{pmatrix} \in \mathfrak{gl}_n.$$

For any  $a, b \in \mathbb{C}^m$ ,

$$\det \left( \begin{array}{c|c} X & \begin{array}{c} | \\ b \\ | \end{array} \\ \hline \begin{array}{c} - & a & - \end{array} & Y \end{array} \right) =$$

$$\det X \det Y + \left( \prod_i d_i \right) \det \left( \begin{array}{c|c} X & \begin{array}{c} | \\ b \\ | \end{array} \\ \hline \begin{array}{c} - & a & - \end{array} & 0 \end{array} \right)$$

*Proof.* By induction on  $n$ . In the case  $n = 1$ , expanding along the last row (taking the usual interpretation of the empty product) gives the desired result.

Now suppose  $n > 1$ . Expanding along the last row, we get

$$d_{n-1} \det \left( \begin{array}{c|c} X & \begin{array}{c} | \\ b \\ | \end{array} \\ \hline \begin{array}{c} - & a & - \end{array} & Y_{n,n-1} \end{array} \right) - y_{nn} \det \left( \begin{array}{c|c} X & \begin{array}{c} | \\ | \\ | \end{array} \\ \hline \begin{array}{c} - & a & - \end{array} & Y_{n,n} \end{array} \right).$$

Using our inductive hypothesis for the first determiniant, and using that  $\det \left( \begin{array}{c|c} A_{11} & 0 \\ \hline A_{21} & A_{22} \end{array} \right) = \det A_{11} \det A_{22}$  for the second, the expression becomes

$$d_{n-1} \left( \det X \det Y_{n,n-1} + \left( \prod_{i \leq n-2} d_i \right) \det \left( \begin{array}{c|c} X & \begin{array}{c} | \\ b \\ | \end{array} \\ \hline \begin{array}{c} - & a & - \end{array} & 0 \end{array} \right) \right) - y_{nn} \det X \det Y_{nn} =$$

$$(d_{n-1} Y_{n,n-1} - y_{nn} \det Y_{nn}) \det X + \left( \prod_i d_i \right) \det \left( \begin{array}{c|c} X & \begin{array}{c} | \\ b \\ | \end{array} \\ \hline \begin{array}{c} - & a & - \end{array} & 0 \end{array} \right) =$$

$$\det Y \det X + \left( \prod_i d_i \right) \det \left( \begin{array}{ccc|c} & & & b \\ & & & | \\ & & & | \\ \hline - & a & - & 0 \end{array} \right).$$

□

**Corollary 5.3.** *If  $X$  is nilpotent, and*

$$\left( \begin{array}{ccc|c} & & & b \\ & & & | \\ & & & | \\ \hline - & a & - & \\ & & & Y \end{array} \right)$$

*is nilpotent as well, then  $Y$  is nilpotent (TODO: and that other determinant is zero).*

*Proof.* By the previous lemma, the characteristic polynomial of the big matrix is

$$g_X(\lambda)g_Y(\lambda) + f(\lambda),$$

where  $g_X(\lambda) = \lambda^m$  is the characteristic polynomial of  $X$ , and  $f(\lambda)$  is some polynomial of degree at most  $m - 1$ . □

Now, taking the previous corollary and the first lemma together, we see that

$$\mathcal{S}'_{m,n} = \left\{ \left( \begin{array}{ccc|cccc} & & & & & & b \\ & & & & & & | \\ & & & & & & | \\ \hline - & a & - & 0 & & & \\ & & & 1 & 0 & & \\ & & & & & \ddots & \\ & & & & & & \ddots & \\ & & & & & & & 1 & 0 \end{array} \right) : a, b \in \mathbb{C}^m, X \in \mathfrak{gl}_m \text{ is nilpotent} \right\}.$$

## 6 Simplifying the definition of a Springer fiber at a Slodowy slice

Let  $\mathcal{N}_m \subseteq \mathfrak{gl}_m(\mathbb{C})$  be the subset consisting of nilpotent elements. For  $X \in \mathcal{N}_m$  and  $a, b \in \mathbb{C}^m$ , let

$$A_{X,a,b} = \left( \begin{array}{c|ccc} X & & & & b \\ \hline \text{---} & a & \text{---} & & \\ & & 0 & & \\ & & 1 & 0 & \\ & & & 1 & \ddots \\ & & & & \ddots & 0 \\ & & & & & 1 & 0 \end{array} \right) \in \mathfrak{gl}_{m+n}(\mathbb{C}).$$

By the definition given in the previous section, we have

$$\tilde{\mathcal{N}}_{m,n} = \{(\mathfrak{b}, X) : X \in \mathfrak{b}\} \subseteq \mathcal{B}_{m+n} \times \mathcal{S}'_{m,n},$$

and the Springer fiber at the  $n$ -Slodowy slice at a nilpotent  $X \in \mathfrak{gl}_m$  is

$$\begin{aligned} \pi_{m,n}^{-1}(X) &= \{(\mathfrak{b}, A_{X,a,b}) : A_{X,a,b} \in \mathfrak{b}, A_{X,a,b} \in \mathcal{S}'_{m,n}\} \\ &= \{(\mathfrak{b}, A_{X,a,b}) : A_{X,a,b} \in \mathfrak{b}, A_{X,a,b} \text{ nilpotent}\} \\ &\cong \{(\mathfrak{b}, a, b) : A_{X,a,b} \in \mathfrak{b}, A_{X,a,b} \text{ nilpotent}\}. \end{aligned}$$

We will make one last simplification to this by using a correspondence between Borel subalgebras and complete flags. TODO: fix the next few paragraphs, they are out of context.

We have  $\mathcal{M} = \{AHA^{-1} : A \in \text{GL}_{m+n}(\mathbb{C})\}$ , where  $H \subseteq \mathfrak{gl}_{m+n}(\mathbb{C})$  is the set of upper triangular matrices. We say a map  $X : \mathfrak{gl}_{m+n} \rightarrow \mathfrak{gl}_{m+n}$  preserves a flag  $V_0 \subseteq \cdots \subseteq V_{m+n}$  if  $XV_i \subseteq V_i$  for each  $i$ . Let  $E_0 \subseteq \cdots \subseteq E_{m+n}$  be the standard flag of  $\mathbb{C}^{m+n}$ . Since  $H$  is the set of  $X$  which preserve  $E$ ,

$$\mathcal{M} = \{\{X : \forall i. X(AE_i) \subseteq AE_i\} : A \in \text{GL}_n(\mathbb{C})\}.$$

So, for  $X \in \mathcal{N}$ ,

$$\pi^{-1}(X) \cong \{(V, a, b) : \forall i. A_{X,a,b}V_i \subseteq V_{i-1}\}.$$



## 7 Strategy and setup for finding the irreducible components of a Springer fiber at a Slodowy slice

Fix any  $X \in \mathcal{N}$ . As we have fixed  $X$ , we now write  $A_{a,b} := A_{X,a,b}$ . Let  $(e_{ij})_{1 \leq i \leq k, 1 \leq j \leq \lambda_i}$  be a Jordan basis for  $X$ . For convenience we define  $e_{i0} := 0$ ; now we may express the fact that  $(e_{ij})$  is a Jordan basis by writing  $\forall i. X e_{ij} = e_{i,j-1}$ .

In this section we find the irreducible components of

$$\pi^{-1}(X) \cong V := \{(A_{a,b}, U) : \forall i. A_{a,b} U_{i+1} \subseteq U_i\}.$$

For  $1 \leq w \leq k$  and  $0 \leq r \leq \lambda_w$  (note that we allow  $r = 0$ ), define

$$V_{w,r} := \{(A_{a,b}, U) \in V : \exists P \in \mathrm{GL}_m. \exists b'. (P^{-1}, I_n) A_{a,b} (P, I_n) = A_{e_{wr}, b'}\}.$$

**Lemma 7.1.**  $V = \bigcup_{1 \leq w \leq k, 0 \leq r \leq \lambda_w} V_{w,r}$ . Further,  $V_{w_1 r_1} = V_{w_2 r_2}$  exactly when either  $r_1 = r_2 = 0$ , or  $\lambda_{w_1} = \lambda_{w_2} \wedge r_1 = r_2$ . When  $V_{w_1 r_1} \neq V_{w_2 r_2}$ , we have  $V_{w_1 r_1} \cap V_{w_2 r_2} = \emptyset$ .

*Proof.* TODO □

Now, fix any  $w$  and  $r$ . We will find the irreducible components of  $V_{w,r}$ . These will all happen to be equidimensional (with dimensions independent of  $w$  and  $r$ ), so their closures in  $V$  will be the irreducible components of  $V$ .

Let

$$G := \{P \in \mathrm{GL}_m : P^{-1} X P = X\},$$

and

$$G_{wr} = \{A \in G : e_{wr} A = e_{wr}\}.$$

Now, define

$$U_{wr} = \{(A_{e_{wr}, b}, U) \in V_{wr}\}.$$

Let  $G$  act on  $V_{wr}$  by

$$P \cdot (A_{e_{wr}, b}, U) := ((P, I_n) A_{e_{wr}, b} (P, I_n)^{-1}, (P, I_n) U) = (A_{e_{wr} P^{-1}, P b}, (P, I_n) U).$$

Consider the map  $\varphi : U_{wr} \times G \rightarrow V_{wr}$  defined by

$$(x, P) \mapsto P \cdot x.$$

By restriction of  $G$  to  $G_{wr}$  and  $V_{wr}$  to  $U_{wr}$ , we obtain an action of  $G_{wr}$  on  $U_{wr}$ . Then, letting  $G_{wr}$  act on  $G$  by  $g \cdot h := hg^{-1}$ , we obtain an action of  $G_{wr}$  on  $U_{wr} \times G$ .

**Lemma 7.2.** *As an algebraic variety,  $G$  is irreducible.*

*Proof.* It's just  $\mathbb{C}^{\text{something}}$ , since its blocks are the lower-left-toeplitz matrices.  $\square$

**Lemma 7.3.** *The map  $\varphi$  is some sort of quotient by the action of  $G_{wr}$ .*

*Proof.* Let  $y \in V_{wr}$ . By definition of  $V_{wr}$ , there is  $P_y \in G$  with  $P_y \cdot y \in U_{wr}$ . We have

$$\begin{aligned} \varphi^{-1}(y) &= \{(x, P) : P \cdot x = y = P_y^{-1} \cdot (P_y \cdot y)\} = \\ &= \{(x, P) : x = (P^{-1}P_y^{-1}) \cdot (P_y \cdot y)\} = \\ &= \{((P^{-1}P_y^{-1}) \cdot (P_y \cdot y), P) : (P^{-1}P_y^{-1}) \in G_{wr}\} = \\ &= (\text{taking } Q := P^{-1}P_y^{-1}, \text{ so } P = (QP_y)^{-1} = P_y^{-1}Q^{-1}) \\ &= \{(Q \cdot (P_y \cdot y), P_y^{-1}Q^{-1}) : Q \in G_{wr}\} = \\ &= \{Q \cdot (P_y \cdot y, P_y^{-1}) : Q \in G_{wr}\}. \end{aligned}$$

So we see that the fibers are exactly the  $G_{wr}$ -orbits.

Now, what kind of quotient is this? All I need is something that allows me to deduce the dimension of the quotient...  $\square$

Our strategy is to find the irreducible components  $X \subseteq U_{wr}$ , and we will then argue that the irreducible components of  $V_{wr}$  are of the form  $\varphi(X \times G)$ . So, we will now find the irreducible components of  $U_{wr}$ .

Actually this will be unnecessarily difficult to think about; it is easiest in the case  $r = 0$ . So, we will change basis to make  $r = 0$ . Let  $\lambda' = (\lambda_1, \dots, \lambda_{w-1}, \lambda_w - r, \lambda_{w+1}, \dots, \lambda_k)$ . Let  $X'$  be in Jordan normal form with shape  $\lambda'$ . Let  $U'_{wr} = \{(A_{X',0,b}, U) : \forall i. A_{X',0,b}U_{i+1} \subseteq U_i\}$ . Let  $e'_{ij}$  be a Jordan basis for  $X'$ . Let  $f_1, \dots, f_n$  be a Jordan basis for the restriction of  $A_{X,0,b}$  to ... Set  $m' := m - r$ , and  $n' := n + r$ . Let  $f'_1, \dots, f'_{n'}$  blah. Define the linear map  $Q_{wr} : \mathbb{C}^{m'+n'} \rightarrow \mathbb{C}^{m+n}$  by:

- For all  $i$ ,  $e'_{ij} \mapsto e_{ij}$ .
- For  $j = 1, \dots, r$ ,  $f'_{n+j} \mapsto e_{w,(\lambda_w-r)+j}$ .

- For  $j = 1, \dots, n$ ,  $f'_j \mapsto f_j + e_{w,(\lambda_w-r)-n+j}$ .

Observe that conjugation by  $Q_{wr}$  maps  $U_{wr}$  to  $U'_{wr}$ , and conjugation by  $Q_{wr}^{-1}$  maps  $U'_{wr}$  to  $U_{wr}$ . We conclude that  $U_{wr} \cong U'_{wr}$ ; so to find the irreducible components of  $U_{wr}$  we just need to find the irreducible components of  $U'_{wr}$ . To clear the context, which is rather cluttered by now, and to avoid writing primes everywhere, we move to a new section.

## 8 Finding the irreducible components of $U_0$

Let  $X$  be nilpotent with Jordan basis  $(e'_{ij})_{1 \leq i \leq k, 1 \leq j \leq \lambda'_i}$ . Let  $V = \{(A_{a,b}, U) : \forall i. A_{a,b}U_{i+1} \subseteq U\}$ . Let

$$U_0 = \{(A_{0,b}, U) \in V\}.$$

We will find the irreducible components of  $U_0$ .

We write  $b'_{ij}$  to denote the projection of  $b \in \mathbb{C}^m$  onto  $e'_{ij}$ . For each row  $i$ , let  $p'_i(b) = \max\{j : b'_{ij} \neq 0\}$  (the maximum of the empty set is zero). Then set  $q'_i(b) = \lambda'_i - p'_i(b)$ . When it is clear enough from context, we will just write  $p'_i$  and  $q'_i$  instead of  $p'_i(b)$  and  $q'_i(b)$ . Then, let  $\{n_1(b), \dots, n_k(b)\} = \{1, \dots, k\}$  be such that the list

$$(q'_{n_1}, n_1), \dots, (q'_{n_k}, n_k)$$

is in reverse lexicographical order; that is, for any  $\alpha < \beta$ , we have either  $q'_{n_\alpha} > q'_{n_\beta}$  or  $q'_{n_\alpha} = q'_{n_\beta} \wedge n_\alpha > n_\beta$ . The point is that we want the list  $q'_{n_1}, \dots, q'_{n_k}$  to be nonincreasing. To break ties in an arbitrary canonical way, we say that if  $q'_\gamma = q'_\delta$ , then we put the one with the bigger subscript first.

Now, in the rest of this section it will be convenient for us to have the values  $q'_i$  in nonincreasing order. To avoid death by double subscript, we define  $e_{ij}(b) := e'_{n_{ij}}$ ,  $\lambda_i(b) := \lambda'_{n_i}$ ,  $p_i(b) := p'_{n_i}$ ,  $q_i(b) := q'_{n_i}$ .

For any  $I \subseteq \{1, \dots, k\}$  and any values  $(\rho_{i_\alpha})_{\alpha \in I'}$  such that (1)  $\rho_{i_\alpha} \leq \lambda_{i_\alpha}$  and (2)  $\rho_{i_\alpha}$  is decreasing with  $\alpha$ , define

$$U_{I',(\rho_i)} := \{(A_{0,b}, U) \in U_0 : I' = \{i : p_i > \max_{j: q_j > q_i} p_j\} \wedge \forall i \in I'. p_i = \rho_i\}.$$

**Lemma 8.1.**  $U_0 = \bigcup_{I',(\rho_i)} U_{I',(\rho_i)}$ , where  $I'$  ranges over all possible subsets of  $\{1, \dots, k\}$ , and  $(\rho_i)$  ranges over all possible maps  $I' \rightarrow \mathbb{N}_{>0}$  such that  $\rho_i$  is decreasing as  $i$  increases.

*Proof.* Let  $(A_{0,b}, U) \in U_0$ . Define  $I' = \{i : p_i > \max_{j: q_j > q_i} p_j\}$ , and for  $i \in I'$ ,  $\rho_i := p_i$ . All we need to do is show that each  $\rho_i$  is positive and that  $\rho_i$  is decreasing. Then we get that  $(A_{0,b}, U) \in U_{I', (\rho_i)}$ , and we are done.

Recall that by convention we take that the maximum of the empty set is zero. So, the fact that  $\rho_i > \max_{j: q_j > q_i} p_j$  tells us that  $\rho_i$  is positive. Now, let  $i < i'$ , with  $i, i' \in I'$ . If we had  $p_i \leq p_{i'}$ , then  $q_i \geq q_{i'}$

□

We note that even finding the irreducible components of  $\{A_{0,0}, U \in V\}$  is a pretty hard problem. Not ambitious enough to try to re-solve that problem, we will use its solution.

To find the irreducible components of  $U_0$ , we will use a known result about the irreducible components of the usual Springer fiber at an element  $J(\mu) \in \mathfrak{gl}_{m+n}$ . (I should define this somewhere else, but  $J(\mu)$  just has the blocks in order of nonincreasing size.)

**Theorem 8.2.** *(Needs citation!) The irreducible components of the Springer fiber at  $J(\mu)$  are in bijection with the standard Young tableaux of shape  $\mu$ . Further, the irreducible components are equidimensional, of dimension  $\sum_{i \neq j} \min(\mu_i, \mu_j) = \sum_i (i-1)\mu_i$ .*

Let  $W_\mu$  denote the Springer fiber at  $J(\mu)$ , and let  $(W_{\mu,d})_d$  be the irreducible components. We remark that the Springer fiber at  $PJ(\mu)P^{-1}$  is  $PW_\mu$ , and its irreducible components are  $(PW_{\mu,d})_d$ . We will use this result; to find the regular Springer fiber at any  $X \in \mathfrak{gl}_{m+n}$ , we just need to find  $P$  such that  $P^{-1}XP$  is in Jordan normal form; equivalently, we just need to find a Jordan basis for  $X$ .

So, our strategy will be to find a subset of the possible values of  $b$  such that  $A_{0,b}$  has some fixed Jordan form  $\mu$ , and we have an algebraic map  $b \mapsto P_b$  such that for  $A_{0,b}$  in the subset,  $P_b J(\mu) P_b^{-1} = A_{0,b}$ . Then we obtain an irreducible component by picking one of these subsets of the possible values of  $b$  and picking a Springer fiber at  $J(\mu)$ .

Finding the map  $b \mapsto P$  amounts to finding a Jordan basis for  $A_{0,b}$ . So, with this as our motivation, we find a Jordan basis for  $A_{0,b}$ .

Normally we index  $e_{ij}$  so that  $\lambda_i$  is nonincreasing with  $i$ . Now, it will be more convenient to assume our indices are such that  $q_i$  is nonincreasing, so we will do that: from now on, we assume that  $q_i$  is nonincreasing.

We list the chains of the Jordan basis in (roughly) order of decreasing length. First, for each  $i$  such that  $q_i \geq n$ , we take the chain of length  $p_i + q_i$

beginning with  $e_{i,p_i+q_i}$ .

Now we handle  $f_n$ . Let  $P = \max_{i:q_i < n} p_i$ . Set

$$v = f_n - (A^{n+P} f_n \gg n + P).$$

Note that  $A^n f_n = b$ , so  $A^{n+P} f_n = A^P b = b \ll P$ . By shifting  $b$  left  $P$  times, we zero out all the rows  $i$  where  $q_i < n$ . This ensures that the operation of shifting  $b' \ll P$  right  $n + P$  times is invertible by applying  $A$ , and thus shifting left,  $n + P$  times. So, we take the chain of length  $n + P$  beginning with  $v$ .

Now, it is clear that each  $f_i$  is in the span of the chains we have listed so far. It is also clear that  $b \ll k$  is in the span, for each  $k \geq 0$ .

Now, we handle the  $i$  with  $q_i < n$ . We take as an inductive hypothesis that for all  $i' < i$  and all  $j$ , we have  $e_{i'j}$  in the span of the chains we have already listed.

If  $p_i \leq \max_{k>i} p_k$ , then we take the chain beginning with  $e_{i,p_i+q_i}$  of length  $p_i + q_i$ . Clearly we then have  $e_{i'j}$  in the span of the chains we have listed, for  $i' \leq i$ .

Otherwise, we set

$$v_i = A^{n-q_i} f_n - \sum_{k=1}^{p_i} b_{i,k} e_{i,k+q_i}.$$

Note that  $A^{q_i} v_i$  is just  $b$  with row  $i$  zeroed out. Then,  $A^{q_i + \max_{k>i} p_k} v_i$  will have, in addition, rows  $k$  zeroed out, for all  $k > i$ . This ensures that shifting  $A^{q_i + \max_{k>i} p_k} v_i$  right  $q_i + \max_{k>i} p_k$  times will be inverted by applying  $A$ , and thus shifting left,  $q_i + \max_{k>i} p_k$  times. So, we take the chain beginning with  $v_i - (A^{q_i + \max_{k>i} p_k} v_i \gg q_i + \max_{k>i} p_k)$ , which has length  $q_i + \max_{k>i} p_k$ .

Now, we want to show that for each  $j$ ,  $e_{ij}$  is in the span of the chains we've listed so far. Since  $A^{q_i + \max_{k>i} p_k} v_i$  has rows  $k$  zeroed out for  $k \geq i$ , our inductive hypothesis tells us that  $A^{q_i + \max_{k>i} p_k} v_i \gg q_i + \max_{k>i} p_k$  is in the span of the chains already listed. So it suffices to show that the closure of  $\langle v_i \rangle$  under action by  $A$  contains the span of  $e_{ij}$ . And  $A^{n-q_i} f_n$  is also in the span of chains already listed, so it suffices to show that the closure of  $\langle \sum_k b_{ik} e_{i,k+q_i} \rangle$  under action by  $A$  contains  $\langle e_{ij} \rangle$ . And this just follows from the fact that  $b_{i,p_i}$  is nonzero.

Now, I've shown that my purported Jordan basis has a large enough span; to show that it is indeed a Jordan basis I just need to count the number of vectors and show that we obtain  $m + n$ . I list the purported basis here:

- TODO

Indeed, the sum of the lengths is

$$\left[ \sum_{i:q_i \geq n+r} (p_i + q_i) \right] + (r+n + \max_{i:q_i < n+r} p_i) + \sum_{i:q_i < n+r} \begin{cases} p_i + q_i, & p_i \leq \max_{k>i} p_k \\ q_i + \max_{k>i} p_k, & p_i > \max_{k>i} p_k \end{cases} =$$

$$\left[ \sum_{i:q_i \geq n+r} (p_i + q_i) \right] + (r+n + \max_{i:q_i < n+r} p_i) + \sum_{i:q_i < n+r} (q_i + \min(p_i, \max_{k>i} p_k)).$$

Let  $i_1 < \dots < i_s$  be the ‘peaks’: that is, the set  $I_1 = \{i : q_i < n+r \wedge p_i > \max_{k>i} p_k\}$ . (As I should’ve mentioned before, we take the max of the empty set to be zero.) Let  $I_2 = \{i : q_i < n+r\} \setminus I_1$ . Then

$$\sum_{i:q_i < n+r} (q_i + \min(p_i, \max_{k>i} p_k)) = \sum_{i \in I_2} (p_i + q_i) + \sum_{k=1}^s (q_{i_k} + p_{i_{k+1}}).$$

Clearly then, the whole sum is  $m+n$ .

## 9 Finding the irreducible components of a springer fiber at a slodowy slice

Apply things from the previous two sections, and conclude.

## 10 Finding a centralizer

For nilpotent  $X$  in Jordan form and  $a$  in ‘normalized’ form (i.e., with at most one nonzero element, which is a one), we will find the centralizer of  $A_{X,a,b}$  in

$$\{(A, I) : A \in \mathfrak{gl}_m\} \subseteq \mathfrak{gl}_{m+n}.$$

Note that an element of the form  $(A, I)$  commutes with  $A_{X,a,b}$  if and only if  $A$  commutes with  $X$ , and  $aA = a$ , and  $Ab = b$ . So, we just have to find

$$\{A \in \mathfrak{gl}_m : AX = XA, aA = a, Ab = b\}.$$

We begin by finding

$$\{A \in \mathfrak{gl}_m : AX = XA\}.$$

We write the shape of  $X$  as  $\lambda = (\lambda_1, \dots, \lambda_k)$ , so that

$$X = \begin{pmatrix} J_{\lambda_1} & & \\ & \ddots & \\ & & J_{\lambda_k} \end{pmatrix}.$$

Then we write  $A$  as the block matrix

$$A =: \begin{pmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & & \vdots \\ A_{k1} & \cdots & A_{kk} \end{pmatrix},$$

where  $A_{ij}$  is a block of size  $\lambda_i \times \lambda_j$ . We have

$$XA = \begin{pmatrix} J_{\lambda_1} A_{11} & \cdots & J_{\lambda_1} A_{1k} \\ \vdots & & \vdots \\ J_{\lambda_k} A_{k1} & \cdots & J_{\lambda_k} A_{kk} \end{pmatrix}, \text{ and } AX = \begin{pmatrix} A_{11} J_{\lambda_1} & \cdots & A_{1k} J_{\lambda_k} \\ \vdots & & \vdots \\ A_{k1} J_{\lambda_1} & \cdots & A_{kk} J_{\lambda_k} \end{pmatrix}.$$

So, the constraint that  $XA = AX$  is simply saying that

$$\forall i, j. J_{\lambda_i} A_{ij} = A_{ij} J_{\lambda_j}.$$

It's easy to see that left multiplication by  $J_{\lambda_i}$  is just a down-shift by one, and right multiplication by  $J_{\lambda_j}$  is a left-shift by one. This means that  $A_{ij}$  is constrained to be a “lower-left-Toeplitz” matrix. A Toeplitz matrix is one which is constant along diagonal bands:  $\forall ijk. x_{ij} = x_{i+k, j+k}$ . A lower-left-Toeplitz matrix is one which is all zeroes except for the bottom-left corner; that is, the bands start in the bottom-left corner, and continue until hitting the band which includes the lower-right corner or the band which includes the upper-left corner (whichever comes first). In other words, an  $n \times m$  lower-left-Toeplitz matrix is one which satisfies  $x_{ij} = 0$  for all  $i, j$  with  $i < j$  and all  $i, j$  with  $i - j < m - n$ . In yet other words, a matrix is lower-left-Toeplitz if it is Toeplitz, satisfies  $x_{1j} = 0$  for  $j \neq 1$ , and satisfies  $x_{in} = 0$  for  $i \neq m$ . Anyway, it is easy to see that the lower-left-Toeplitz matrices are exactly those matrices  $A$  such that left-shifting  $A$  by one has the same result as down-shifting by one.

So, we have found the set

$$C_1 := \{A \in \mathfrak{gl}_m : AX = XA\}.$$

It is just the set of  $A = [A_{ij}]_{ij}$ , where each  $A_{ij}$  is lower-left-Toeplitz.

Let  $v_{ij}$  be the leftmost column of  $A_{ij}$ , so that

$$A_{ij} = \begin{pmatrix} v_{ij} \mathbin{\mathbb{V}} 0 & \cdots & v_{ij} \mathbin{\mathbb{V}} [\lambda_j - 1] \end{pmatrix}.$$

Since  $A_{ij}$  is lower-left-Toeplitz,  $v_{ij}$  is of the form

$$v_{ij} = \begin{pmatrix} 0 \\ v'_{ij} \end{pmatrix},$$

where  $v'_{ij} \in \mathbb{C}^{\min(\lambda_i, \lambda_j)}$  can be chosen freely. Now we determine which matrices of this form satisfy  $aA = a$ . For simplicity, we will instead find the matrices  $A$  such that  $aA = 0$  (so that  $a(A + I) = a$ ). (Note that  $I \in C_1$ , and  $C_1$  is closed under addition, so this is really the same thing.)

In the case that  $a = 0$ , clearly every  $A$  works. Otherwise, let  $i_0, j_0$  be such that  $a_{i_0, j_0} = 1$ . Now, clearly the constraint that  $aA = 0$  is just saying that the  $(i_0, j_0)$ th row of  $A$  must be zero. That is, for each  $j$  the  $j_0$ th row of  $A_{i_0 j}$  must be zero. This just requires that for each  $j$ , we must have

$$v_{ij} = \begin{pmatrix} 0 \\ v'_{ij} \end{pmatrix},$$

where  $v'_{ij} \in \mathbb{C}^{\min(\lambda_i - j_0, \lambda_j)}$  can be chosen freely. So, we have now found the set

$$C_2 := \{A \in \mathfrak{gl}_m : AX = XA, aA = a\}.$$

It is the matrices of the form  $I + A$ , where  $A_{ij}$  is a block matrix of size  $\lambda_i \times \lambda_j$  with

$$A_{ij} = T(v_{ij}) = T \begin{pmatrix} 0 \\ v'_{ij} \end{pmatrix},$$

where  $v'_{ij} \in \mathbb{C}^{\min(\lambda_i, \lambda_j)}$  in the case  $i \neq i_0$  and  $v'_{ij} \in \mathbb{C}^{\min(\lambda_i - j_0, \lambda_j)}$  if  $i = i_0$ .

## 11 A Jordan basis for $A_{X,a,b}$

### 11.1 A ‘normalization’ fact about Jordan bases

Let  $V$  be a finite-dimensional vector space,  $A : V \rightarrow V$  a nilpotent operator, and  $f : V \rightarrow \mathbb{C}$  a linear map. Let  $(e_{ij} : i \leq m, j \leq \lambda_i)$  be a Jordan basis for  $A$ .



**Lemma 11.1.** *There is a Jordan basis  $e_{ij}$  for  $A$  such that there is at most one pair  $(i, j)$  with  $f(e_{ij}) \neq 0$ .*

*Proof.* For any change of basis  $P : V \rightarrow V$  commuting with  $A$ , we obtain a new Jordan basis  $(P(e_{ij}) : i \leq m, j \leq \lambda_i)$ . For any such  $P$ , define

$$S_P = \sum_i \begin{cases} -1, & \forall j. f \circ P(e_{ij}) = 0 \\ \lambda_i - \min\{j : f \circ P(e_{ij}) \neq 0\}, & \text{otherwise} \end{cases}.$$

Let  $P$  be any operator, among all invertible operators commuting with  $A$ , which minimizes  $S_P$ . Write  $e'_{i,j} := P(e_{ij})$ .

Suppose for contradiction that there are two distinct  $i$ 's (and some  $j$ 's) with  $f(e'_{ij}) \neq 0$ . Then we can take  $e'_{i_1,j_1}$  and  $e'_{i_2,j_2}$ , where for  $k \in \{1, 2\}$  we have  $f(e'_{i_k,j_k}) \neq 0$ , and  $\forall j < j_k. f(e'_{i_k,j}) = 0$ . Wlog, assume  $\lambda_{i_1} - j_1 \leq \lambda_{i_2} - j_2$ . Then, we can define  $Q : V \rightarrow V$  by

- $Q(e'_{i_1,\lambda_1}) := e'_{i_1,\lambda_1} - \frac{f(e'_{i_1,j_1})}{f(e'_{i_2,j_2})} e'_{i_2,j_2+(\lambda_{i_1}-j_1)}$
- For  $j < \lambda_1$ ,  $Q(e'_{i_1,j}) := A^{\lambda_{i_1}-j} Q(e'_{i_1,\lambda_1})$
- For  $i \neq i_1$ ,  $Q(e'_{ij}) = e'_{ij}$ .

Clearly  $Q$  is invertible, and it commutes with  $A$ . Further, I claim that  $S_{QP} < S_P$ . It suffices to show that  $\forall j \leq j_1. f \circ Q(e'_{i_1,j}) = 0$ . We have

$$\begin{aligned} f \circ Q(e'_{i_1,j}) &= f \left( A^{\lambda_{i_1}-j} \left( e'_{i_1,\lambda_1} - \frac{f(e'_{i_1,j_1})}{f(e'_{i_2,j_2})} e'_{i_2,j_2+(\lambda_{i_1}-j_1)} \right) \right) = \\ &= f \left( e'_{i_1,j} - \frac{f(e'_{i_1,j_1})}{f(e'_{i_2,j_2})} e'_{i_2,j_2+(j-j_1)} \right) = f(e'_{i_1,j}) - \frac{f(e'_{i_1,j_1})}{f(e'_{i_2,j_2})} f(e'_{i_2,j_2+(j-j_1)}). \end{aligned}$$

Clearly (by design), this expression is zero when  $j = j_1$ . And for  $j < j_1$ , we have  $f(e'_{i_1,j}) = f(e'_{i_2,j_2+(j-j_1)}) = 0$ , so it is zero then as well. Thus we see that  $S_{QP} < S_P$ , contradicting that  $S_P$  is minimal. So there must be at most one  $i$  such that there exists  $j$  such that  $f(e'_{ij}) \neq 0$ .

If there is no such  $i$ , we have found the desired basis. So, suppose there is such an  $i_0$ . Let  $U = \langle e'_{i_0,j} \rangle$ . Simply write  $e_j := e'_{i_0,j}$ . Let  $j_0 = \min\{j : f(e_j) \neq 0\}$ . We just have to change basis to zero out  $e_j$  for  $j \neq j_0$ . Let  $P_1(e_{\lambda_{i_0}}) := e_{\lambda_{i_0}} - \frac{f(e_{j_0+1})}{f(e_{j_0})} e_{\lambda_{i_0}-1}$ , and notice that  $f \circ P_1(e_j) = 0$  for all  $j \leq j_0 + 1$

except for  $j_0$ . Then we define  $P_2(e_{\lambda_{i_0}}) := P_1(e_{\lambda_{i_0}}) - \frac{f \circ P_1(e_{j_0+2})}{f \circ P_1(e_{j_0})} e_{\lambda_{i_0}-2}$ , and notice that  $f \circ P_2(e_j) = 0$  for all  $j \leq j_0 + 2$  except for  $j_0$ . Eventually we get  $P_{\lambda_{i_0}-j_0}$ , and by applying this to the  $e_j$ 's we obtain a basis of  $U$ , in which there is exactly one  $j$  with  $f \circ P_{\lambda_{i_0}-j_0}(e_j) \neq 0$ .  $\square$

## 11.2 When $X$ is all zeroes

In this case we have  $A(y, z) = (z_n b, (a \cdot y, z_1, \dots, z_{n-1}))$ , and the condition becomes

$$\sum_i b_i a_i = 0.$$

In this case we will be able to write down explicitly the irreducible components of  $F := \{(V, a, b) : \forall i. A_{X,a,b} V_i \subseteq V_i\} \cong \pi^{-1}(X)$ . For any nonnegative  $\delta_0, \delta_1, \dots, \delta_n, \delta_{n+1}$  summing to  $m$ , define the corresponding sequence  $i_0 = \delta_0$ ,  $i_n = \delta_{n+1} + i_n$ , and for  $j \in \{1, \dots, n\}$ ,  $i_j = i_{j-1} + 1 + \delta_j$ . Let  $E$  be the span of the  $e_i$ 's, and let  $E' = \{(x, 0) \in E : x \cdot a = 0\}$ , where the dot is the  $m$ -dimensional dot product. Then define  $F_\delta$  as the set of  $(V, a, b) \in F$  such that

- $b \in V_{i_0} \subseteq E'$
- for all  $j \in \{1, \dots, n\}$ , we have  $f_j \in V_{i_j} \subseteq E' + \langle f_1, \dots, f_j \rangle$

I claim that the  $F_\delta$ 's are the irreducible components of  $F$ . To begin, I show that their union is  $F$ .

**Lemma 11.2.** *Let  $(V, a, b) \in \mathcal{F}$ . Write  $f_0 = b$ , and  $F = \langle f_0, f_1, \dots, f_n \rangle$ . For each  $i$ , either  $F \subseteq V_i$ , or else there exists  $j$  such that  $e_j \notin V_i$ , but  $\langle e_0, \dots, e_{j-1} \rangle \subseteq V_i \subseteq E' + \langle e_1, \dots, e_j \rangle$ .*

*Proof.* For  $i = 0$ , we may take  $j = 0$ . Now assume the statement holds for  $i$ , and we will prove it for  $i + 1$ . If  $F \subseteq V_i$ , then  $F \subseteq V_{i+1}$ , and we are done.

So, suppose there is  $j$  such that  $e_j \notin V_i$ , but  $\langle e_0, \dots, e_{j-1} \rangle \subseteq V_i \subseteq E' + \langle e_0, \dots, e_j \rangle$ . We have two cases: either  $V_{i+1} = V_i + \langle e_j \rangle$ , or not.

- If so, then either  $j = n$ , in which case  $F \subseteq V_{i+1}$ , or else  $j \neq n$ , in which case  $e_{j+1} \notin V_{i+1}$ , but  $\langle e_0, \dots, e_j \rangle \subseteq V_{i+1} \subseteq E' + \langle e_0, \dots, e_{j+1} \rangle$ .
- If not, then  $e_j \notin V_{i+1}$ . Let  $v_{i+1}$  be so that  $V_{i+1} = V_i + \langle v_{i+1} \rangle$ . I just need to show that  $v_{i+1} \in Y + \langle e_1, \dots, e_j \rangle$ . It suffices to show that  $v_{i+1}^\top e_k = 0$

for  $k > j$ . And to do this, it suffices to show that  $(Av_{i+1})^\top e_{k-1} = 0$  for  $k > j$ .

Note that  $Av_{i+1} \in V_i \cap A\mathbb{C}^{2n} \subseteq \langle e_1, \dots, e_j \rangle$ . So, for  $k > j + 1$  it is clear that  $(Av_{i+1})^\top e_{k-1} = 0$ . Now, suppose for contradiction that  $(Av_{i+1})^\top e_j \neq 0$ . Then  $Av_{i+1}$  is linearly independent of  $e_1, \dots, e_{j+1}$ . Since  $Av_{i+1} \in \langle e_1, \dots, e_j \rangle$ , it follows that  $e_j \in \langle e_1, \dots, e_{j-1}, Av_{i+1} \rangle \subseteq V_i$ , a contradiction.

□

**Corollary 11.3.**  $\mathcal{F} = \bigcup_\delta \mathcal{F}_\delta$

*Proof.* Let  $(V, a, b) \in F$ . Let  $i_0 = \min\{i : b \in V_i\}$ , and let  $i' = \max\{i : V_i \subseteq E'\}$ . We want that  $b \in V_{i_0} \subseteq E'$ , so we want that  $i_0 \leq i'$ . For contradiction, suppose  $i' < i_0$ .

□

**Lemma 11.4.** Each  $\mathcal{F}_\delta$  is a closed subvariety of  $\mathcal{F}$ .

*Proof.*

□

**Lemma 11.5.** Each  $\mathcal{F}_\delta$  is irreducible of dimension  $m$ .

*Proof.* Let  $\mathcal{E}$  be the variety of partial flags of  $E$  of shape  $(\delta_0, \dots, \delta_{n+1})$ . Define  $g : \mathcal{F}_\delta \rightarrow \mathcal{E}$  by

$$(V, a, b) \mapsto 0 \subseteq V_{i_0} \cap E \subseteq V_{i_1} \cap E \subseteq \dots \subseteq V_{i_{n+1}} \cap E = E.$$

It is clear that  $g$  is surjective, as the definition of  $\mathcal{F}_\delta$  places no restriction on the intersections  $V_i \cap E$ . Now let's look more closely at the fibers  $g^{-1}(U)$ .

First let's find the flags  $V$  such that there exist  $a, b$  with  $g(V, a, b) = U$ . We see that, for instance,  $V_{i_0} \cap E = U_1$ . In fact  $V_{i_0} \subseteq E$ , so  $V_{i_0} = U_1$ . But we are free to choose the vector spaces between 0 and  $V_{i_0}$  however we wish, so we get some degrees of freedom like  $\mathcal{F}_{\delta_0}$ , the complete flag variety on  $\mathbb{C}^{\delta_0} \cong V_{i_0}/0$ . Similarly, for every  $j = 1, \dots, n + 1$ , we can choose the vector spaces between  $V_{i_{j-1}}$  and  $V_{i_j}$  arbitrarily, so we get degrees of freedom like  $\mathcal{F}_{\delta_{i_j}}$ , the complete flag variety on  $\mathbb{C}^{\delta_{i_j}} \cong V_{i_j}/V_{i_{j-1}}$ . Finally, to meet the constraint of  $(V, a, b)$  being in  $\mathcal{F}_\delta$ , we can choose any  $b \in U_1$  and any  $a$  such that  $\bar{a} \in U_n^\perp$ . Thus we get an isomorphism

$$g^{-1}(U) \cong \mathcal{F}_{\delta_0} \times \dots \times \mathcal{F}_{\delta_{i_{n+1}}} \times \mathbb{C}^{\delta_0} \times \mathbb{C}^{\delta_{n+1}}.$$

□

**Theorem 11.6.** The  $\mathcal{F}_\delta$ 's are the irreducible components of  $\mathcal{F}$ .

### 11.3 In the case that $X$ is a Jordan block

This case seems harder to work with explicitly than the case that  $X$  is zero, so our strategy is to reuse

## 12 A different variety

Define  $R = \{(X, \mathfrak{b}_1, \mathfrak{b}_2) : X \in \mathfrak{b}_1 \wedge u(X) \in \mathfrak{b}_2\} \subseteq \mathcal{S}'_{m,n} \times \mathcal{B}_{m+n} \times \mathcal{B}_m$ . We can obtain a subvariety of  $R$  by requiring that  $u(X)$  is in some fixed similarity class. (TODO: why is this a subvariety? Is it? Is this even the right way of explaining the significance of the  $m^2$ ?) We expect that each of these subvarieties is an irreducible component of dimension  $m^2$ . We will verify these things using our previous computations of springer fibers.

### 12.1 TODO

- Why are SOn flags what they are.

## References

- [1] N. Chriss and victor ginzburg. *Representation Theory and Complex Geometry*. Modern Birkhäuser Classics. Birkhäuser Boston, 2009.