

School of Physics



The Fast Transient Sky

Stage Transfer Report

Owen Johnson[†]
Astrophysics Group
January, 2024

Abstract

The abstract is a short concise outline of your project area, **of no more than 100 words.**

Supervisor: Assoc. Prof. Evan Keane

[†]ojohnson@tcd.ie

Contents

1	A Prelude to Pulsars	1
1.1	The Population of Pulsars	1
1.2	Pulsar Subclasses	2
1.3	Spider Pulsars	3
1.3.1	4FGL J0523-2529	4
1.3.2	4FGL J2054-6904	4
2	Work This Far	4
2.1	Search for Extraterrestrial Intelligence	4
2.2	M-dwarf Radio Flares	4
3	Summary	4

Declaration

I hereby declare that this report is entirely my own work and that it has not been submitted as an exercise for a degree at this or any other university.

I have read and I understand the plagiarism provisions in the General Regulations of the University Calendar for the current year, found at <http://www.tcd.ie/calendar>.

Signed: _____

Date: _____

Publications and Presentations

Publications

Johnson, O.A., Gajjar, V., Keane, E.F., et. al (2023). Simultaneous dual-site SETI with LOFAR international stations. Manuscript accepted for publication to AJ. arXiv:2310.15704

Presentations

1. Low Frequency's Place in SETI, January, 2024, PSETI Symposium, Penn State.
[Invited]
2. Technosignatures with NenuFAR, December, 2023, Science at Low Frequencies IX, UvA.
3. SETI Science at 30 - 190 MHz, November, 2023, BLUK Workshop, SKAO.
[Invited]
4. Technosignature Science at Low Frequencies, November, 2023, NASA Goddard Flight Center.
[Invited]
5. Dual Site SETI Searches, 2023, International Astronautical Congress, Baku.

1 A Prelude to Pulsars

When stars with a mass of at least $8 M_{\odot}$ reach the end of their evolutionary stage and experience a depletion of nuclear fuel will undergo a core collapse and explode as a Supernovae. Depending on the mass of the host star the Supernova will form a Black Hole or a Neutron Star. Based on the electron degeneracy pressure limit (Chan 1967) stars that fall in the range of $20 - 30 M_{\odot}$ form neutron stars (need citation).

Neutron stars are supported against further collapse by the presence of neutron degeneracy pressure which arises from the Pauli exclusion principle. Strong Nuclear forces between the neutrons also provides additional support against gravitational collapse. With these two opposing forces a stable equilibrium is formed.

In turn, this makes neutron stars exceptionally dense, they become the densest known objects in the universe that emit light. The average density of a neutron star is 10^{17}kg/m^3 (need citation). The radius of neutron stars are comparable to the size of cities, with radii of $10 - 20 \text{ km}$.

During collapse the conservation of magnetic flux plays a crucial role in the large strength magnetic fields that are observed in neutron stars along with contributions from the dynamo effect and frozen-in magnetic fields. The magnetic field of a neutron star are of the order of $10^{12} - 10^{15} \text{ Gauss}$ (Need citation).

Charged particles accelerate along the magnetic field lines in the magnetosphere of the neutron star. The particles emit electromagnetic radiation in a cone shape along the magnetic axis. If the magnetic axis is not aligned with the rotational axis of the neutron star, the radiation beam will sweep across the sky. This is known as a pulsar and are analogous to cosmic lighthouses.

1.1 The Population of Pulsars

As of writing there are currently more than 3380 known pulsars. Since their discovery in 1967 by Jocelyn Bell Burnell the population has grown immensely but there remains many questions about pulsar evolution and the subclasses that lie within the population as a whole. Similarly to how exoplanet populations are shown using the mass-radius diagram and stellar populations are shown using the Hertzsprung-Russell diagram, pulsar populations are shown using what is known as the $P - \dot{P}$ diagram.




Figure 1: The $P - \dot{P}$ diagram showing the population of pulsars. The the millisecond pulsar subclasses are colour coded. The red region represents the death line, where pulsars are theoretically no longer able to emit radio waves.

P representing the Pulsar's rotational period and \dot{P} it's derivative. These are key ways that pulsars are classified and study in context of their evolution. An example of a $P - \dot{P}$ diagram is shown in fig. 1. Different values on the plot indicate the roughly the Pulsars age and magnetic field strength. Figure 1 shows the vastly different values between pulsars in the millisecond range and pulsars in the second range.

Theoretically it has been shown that pulsars exhibit a death line in the $P - \dot{P}$ diagram. This is the line where pulsars are no longer able to emit radio waves. This is due to the fact that the pulsar's magnetic field is no longer strong enough to accelerate particles along the magnetic field lines. However it has been shown that pulsars do exist below this line. The area below this line is commonly referred to as the "graveyard".

1.2 Pulsar Subclasses

The population of pulsars can be broken down into subclasses based on unique patterns in their properties. The main subclasses¹ are as follows:

1. Normal Pulsars: These are the most common type of pulsars. They are character-

¹This is a non-exhaustive list.

ized by their regular pulses and are often observed in radio wavelengths. They are also known as radio pulsars.

2. Rotating Radio Transients (RRATs): These are a subclass of pulsars that were initially discovered through their sporadic radio bursts rather than regular pulses. They exhibit irregular and infrequent radio emission.
3. Magnetars: While not exclusively pulsars, magnetars are highly-magnetized neutron stars that can also emit pulsed radiation. They are characterized by extremely strong magnetic fields, much more intense than typical pulsars.
4. Binary Pulsars: These are pulsars that are in orbit around another star, usually a normal (non-neutron) star. The interaction with the companion star can have significant effects on the pulsar's behavior, and studying binary pulsars has provided important tests of theories of gravitation.
5. Millisecond Pulsars (MSPs): These are pulsars with very short rotation periods, typically less than 10 milliseconds. They are believed to be old pulsars that have been spun up by the accretion of mass from a companion star in a binary system.
6. Gamma-ray Pulsars: Pulsars that emit pulsed gamma-ray radiation are known as gamma-ray pulsars. These are often detected by space-based telescopes like the Fermi Gamma-ray Space Telescope.
7. X-ray Pulsars: Pulsars that emit pulsed X-ray radiation fall into this category. These pulsars are typically observed in binary systems where the pulsar accretes matter from its companion star, leading to X-ray emission.
8. Anomalous X-ray Pulsars (AXPs) and Soft Gamma-ray Repeaters (SGRs): These are closely related to magnetars and are characterized by their intense and variable X-ray and gamma-ray emission. They are believed to be neutron stars with extremely strong magnetic fields.

1.3 Spider Pulsars

The type of pulsar that is of interest to this project is a subclass of transional miliscond pulsars known as Spider Pulsars. Spider pulsars fall into two categories depending on their orbiting companion. The first category are known as Black Widow pulsars segreated based on their companion mass falling in the range of $0.01 - 0.05 M_{\odot}$ with a compantion orbital period (P_B) of less than 10 hours (citation). The second category are known as Redback pulsars and have a companion mass of $0.2 M_{\odot}$ or greater with a P_B of less than 1 day (citation).

1.3.1 4FGL J0523-2529

1.3.2 4FGL J2054-6904

2 Work This Far

2.1 Search for Extraterrestrial Intelligence

2.2 M-dwarf Radio Flares

3 Summary

References