

Qiushi Engine

Large-Scale Autonomous Discovery of Kissing Number Constructions

Complete Research Report

Shuxing Yang	Rui Zhao	Junyao Wu	Yize Wang
Fujia Chen	Kaihao Zhu	Wenhao Li	Zichen Li
Yaqi Li	Shenzhan Hong	Yuang Pan	Junjie Yang
Taowen Deng	Jincheng Mi	Hongsheng Chen*	Yihao Yang*

College of Information Science and Electronic Engineering, Zhejiang University
Qiushi Engine Team

*Correspondence: Hongsheng Chen (hansomchen@zju.edu.cn); Yihao Yang (yangyihao@zju.edu.cn)

Abstract

The kissing-number problem is a classical problem in discrete geometry whose exact solution is known in only a few dimensions. Recent artificial-intelligence approaches have begun to discover improved configurations through large-scale numerical and combinatorial search, but converting such searches into general mathematical constructions and rigorous proofs remains challenging. Here we use Qiushi Engine, an autonomous multi-agent research system, to investigate kissing numbers and obtain new lower bounds in nineteen dimensions: 25, 27, 32–39, 43, 45, and 49–55. The resulting constructions arise from distinct structural mechanisms, including coordinated motions of contact layers, labelled direction reuse, joint support exchanges, signed-code replacements, cross-shell lattice constructions, low-overlap lattice isometries, and spherical-design moment certificates. They yield sharp capacities for parameterized signed-code models, deterministic image-union guarantees, and exact section and projection counts controlled by embedded root systems and anchor-graph statistics. These methods yield, among others, $K(25) \geq 197580$, $K(27) \geq 201567$, $K(38) \geq 591900$, $K(43) \geq 2553792$, $K(45) \geq 7380090$, and $K(55) \geq 53301140$. The autonomous system carried out the construction searches, mathematical analysis and computational verification, while each final result was reduced to explicit mathematical arguments and independently checkable finite certificates. Our results illustrate how autonomous research systems can move beyond optimization within fixed formulations to discover new mathematical representations and constructions at scale.

Keywords Autonomous mathematical discovery; kissing number; geometric deformation; constant-weight code; lattice lifting; signed design; spherical design

Contents

1	Introduction	3
1.1	The spherical formulation	5
1.2	Results	5
1.3	Construction principles	5
1.4	Mathematical context	7
1.5	Reading the constructions	7
2	Coordinated shell motion in dimension 25	7
3	Labelled lifts: count the directions that remain	10
3.1	Multiplicity and the four-triangle construction	11
3.2	The later 201,567-point construction	11
4	Layered codes, joint exchanges and signed constructions	12
4.1	Compatibility of the layers	12
4.2	Joint exchanges of supports	13
4.3	Hadamard transfer and local replacement	14
5	Equatorial completion of a Leech-lattice lifting	17
5.1	The lifted configuration and its equatorial constraints	17
5.2	Compatible lines in the second shell	18
5.3	Contacts and limits of the shell choice	19
6	Seven dimensions from one enlarged line class	19
6.1	Disjoint blocks from overlapping images	19
6.2	Why the lift preserves all angular constraints	20
6.3	Separating the two sources of gain	21
7	Extremal lattice sections and design moments	21
7.1	Mother shells and moment equations	21
7.2	Dimension 43: the fixed child and eight coordinate cuts	22
7.3	Dimension 45: an exact fibre identity	23
7.4	The anchor graph and the certified count	24
8	Construction principles	24
A	The minimum of the ambient lattice used in dimension 45	25
B	Construction data and exact verification	25
C	Construction methods and research development	27
C.1	Moving the retained shell	27
C.2	From assigning blocks to reusing directions	27
C.3	From block libraries to image unions above dimension 48	27
	References	29

1 Introduction

The kissing-number problem is one of the oldest problems in discrete geometry. In 1694, Isaac Newton and David Gregory discussed whether 12 or 13 equal spheres could simultaneously touch a central sphere in three dimensions. The answer, 12, was proved more than two centuries later [1]. Yet the apparent simplicity of the question is deceptive. More than three hundred years after its formulation, exact kissing numbers are known only in dimensions 1, 2, 3, 4, 8 and 24. In the exceptional dimensions 8 and 24, the optimal configurations arise from the E_8 and Leech lattices, with 240 and 196560 contacts, respectively [1, 2]. In most other dimensions, substantial gaps remain between the best constructions and known upper bounds.

Progress on lower bounds has historically relied on explicit mathematical structure. Lattices and error-correcting codes provide large families of vectors whose pairwise interactions can be controlled simultaneously [2, 3]. These ideas have led to laminated-lattice constructions, lattice sections and extensions, code-based spherical configurations, and lifting methods that combine a high-dimensional mother configuration with additional coordinates [4–8]. Such constructions can contain hundreds of thousands or even tens of millions of vectors. Their strength, however, is also a source of difficulty: once a configuration is highly constrained, improving it may require reorganizing an entire collection of points rather than inserting a single new one. The relevant search spaces grow combinatorially, while the geometric degrees of freedom that permit improvement can be difficult to identify from the original representation.

Artificial intelligence has recently begun to alter how these spaces are explored. AlphaEvolve, a general-purpose coding agent, discovered a 593-point kissing configuration in dimension 11, improving the previously known lower bound of 592 [9]. PackingStar formulated the kissing-number problem as a two-player matrix-completion game and used game-theoretic reinforcement learning to search the space of Gram matrices, reporting new constructions across dimensions 25–31 and uncovering thousands of candidate geometric structures [10]. EinsteinArena allowed autonomous agents to submit solutions, inspect intermediate results and build on discoveries made by other agents; this collective process increased the dimension-11 lower bound from 593 to 604 [11]. The Station subsequently reported exact 604-point configurations and structural theorems developed in an open-world multi-agent environment [12]. Together, these developments demonstrate the scope of AI-guided geometric search and collaborative mathematical discovery.

These advances also expose a deeper challenge. Finding an improved configuration is only one part of mathematical discovery. Why does the configuration exist, which structural degree of freedom makes it possible, can the mechanism be generalized, and how can the result be converted into a proof or an independently verifiable certificate? Computational searches begin from a prescribed representation—for example, coordinates, Gram matrices, codes or a fixed optimization objective. Open-ended mathematical research additionally requires the ability to change that representation when the current one becomes limiting, to formulate new intermediate problems, connect apparently unrelated structures, and develop arguments that certify the resulting construction.

Autonomous scientific research systems provide a possible route towards this broader form of discovery. Recent large-language-model-based agents can carry out extended sequences of reasoning, computation and tool use, while revising research strategies in response to intermediate evidence. Qiushi Engine was previously used for end-to-end autonomous scientific discovery on a physical optical platform [13], where it conducted long-horizon investigations that combined theoretical reasoning, computation and experiment. Here we ask whether the same paradigm can be extended to open mathematical research: can an autonomous research system move

beyond optimizing a fixed mathematical formulation and instead discover new representations, constructions and proofs?

Here we report an autonomous investigation of the kissing-number problem that yields new lower bounds in nineteen dimensions:

$$25, \quad 27, \quad 32\text{--}39, \quad 43, \quad 45, \quad 49\text{--}55.$$

The resulting bounds include

$$\begin{aligned} K(25) &\geq 197580, & K(27) &\geq 201567, & K(38) &\geq 591900, \\ K(43) &\geq 2553792, & K(45) &\geq 7380090, & K(55) &\geq 53301140. \end{aligned}$$

Rather than arising from a single optimization procedure, these improvements were obtained through a sequence of distinct structural discoveries. They include coordinated motions of complete contact layers, labelled liftings that separate direction deletion from reuse, joint support exchanges that exploit shared deletion costs, signed-code replacements generated through Hadamard transfers, compatible directions drawn from a different lattice shell, enlarged lattice-line classes combined through low-overlap isometric images, and spherical-design moment identities and inequalities that determine or bound large lattice sections without enumerating the ambient shell.

A common pattern connects these apparently different constructions. Configurations that appear saturated in one representation retain exploitable freedom in another. In dimension 25, moving an entire 552-point contact layer while preserving its internal Gram matrix creates room for an additional point. In dimension 27, labelled lifting separates direction reuse from deletion; with the first layer fixed, jointly selecting second-layer directions and labels enlarges the final two-layer construction. In dimensions 32–37 and 39, optimizing collective exchanges and individual sign patterns exposes improvements hidden by separate support-insertion scores and complete sign bundles. In dimension 38, changing the search from the minimal Leech shell to a second lattice shell produces 288 additional points. In dimensions 43 and 45, counting millions of minimal lattice vectors reduces to a finite space of projection signatures whose multiplicities satisfy exact spherical-design moments. A single enlarged line class in dimension 48, combined with low-overlap isometric images, similarly generates new constructions throughout dimensions 49–55.

Qiushi Engine carried out the construction searches, mathematical analysis and computational verification. Each discovered mechanism is expressed through explicit geometric or combinatorial conditions. Each reported bound is supported by finite witnesses checked using exact integer and rational arithmetic, lattice and code identities, rigorous interval arithmetic, and finite combinatorial verification. In dimension 25, outward-rounded interval arithmetic certifies the remaining strict inequalities [14]. These mathematical arguments and reproducible certificates make the constructions independently checkable.

These results suggest a role for AI in mathematics that extends beyond accelerating numerical search. For difficult problems with enormous structured search spaces, a central task may be to discover the representation in which progress becomes possible. Autonomous research systems that can alternate between exploration, structural interpretation and rigorous verification may provide a route towards such discoveries at scale.

1.1 The spherical formulation

The kissing number $K(n)$ is the largest number of unit balls that can touch a central unit ball in $\mathbb{R}^n$ without overlapping interiors. Dividing their centres by two gives the equivalent spherical problem

$$K(n) = \max\{|X| : X \subset S^{n-1}, \langle x, y \rangle \leq \frac{1}{2} \quad (x \neq y)\}.$$

Thus the angle between distinct directions is at least 60° .

1.2 Results

Theorem 1.1. *For every dimension n in Table 1, there is a kissing configuration of N_n points, where N_n is the lower bound in the second column. Thus $K(n) \geq N_n$.*

The constructions in this report give the following lower bounds. The comparison column uses the fixed public sources recorded on 2026-09-24 to 2026-09-27.

Table 1 Explicit lower bounds and their public comparison values.

Dimension	Lower bound	Public comparison	Increase	Source
25	197580	197579	1	[15]
27	201567	201566	1	[16]
32	347584	346944	640	[17]
33	363968	362048	1920	[17]
34	384196	381124	3072	[17]
35	409676	409548	128	[17]
36	484760	484568	192	[17]
37	498024	496232	1792	[17]
38	591900	591612	288	[15]
39	763668	756116	7552	[15]
43	2553792	2545056	8736	[18, 19]
45	7380090	7379838	252	[18, 19]
49	52430156	52430140	16	[15]
50	52458468	52458418	50	[15]
51	52500930	52500816	114	[15]
52	52585772	52585516	256	[15]
53	52698696	52698222	474	[15]
54	52923774	52922906	868	[15]
55	53301140	53299730	1410	[15]

The dimension-32 comparison uses Lysenstøen’s 1671-word code listed in Brouwer’s table [17]. The ERS formula gives $2^{17} + 128 \cdot 1671 + 2 \cdot 32 \cdot 31 = 346944$. The 1667-word code of Echols [20] remains the seed used in our search.

1.3 Construction principles

The nineteen bounds arise from seven connected mechanisms. Each changes a specific mathematical object and has a corresponding compatibility or counting argument.

Coordinated motion in dimension 25. A contact layer of 552 points has fixed transverse coordinates and a common latitude. Moving the latitude together with the extra coordinate

preserves every internal product. The horizontal projection remains on chords joining old shell points, so convexity controls the unchanged equator. Two cap repairs complete the construction and increase the count from 197579 to 197580. The gain comes from moving retained points, not from selecting another point in a fixed pool.

Direction reuse and joint labels in dimension 27. A lifted direction contributes several copies but incurs only one equatorial deletion. Tail geometry determines the permitted reuse. The four-triangle construction realizes the capacity of a prescribed twelve-label model. The larger two-layer construction then combines a fixed first layer with a new joint selection of second-layer owners and labels. Selecting 311 owners instead of 310 gives 201567 points. The capacity statement and the final two-layer count concern different stages of the same lifting geometry.

Shared deletion costs in dimensions 32–34, 37 and 39. For a compatible family Y of new supports, the old supports to remove form a union $F(Y)$. The gain is $|Y| - |F(Y)|$. Several insertions may therefore succeed together even when none improves the count alone. The new support families have sizes 1676, 1803, 1960, 2845 and 3383, respectively. Their intersections and the distances of the accompanying dense codes establish the complete layered constructions.

Signed design transfer in dimensions 35–37. Coordinate regions with controlled interaction with the retained layers admit larger signed replacements. Pairing coordinates and applying a Hadamard transform carries transverse weight-four blocks to weight-eight signed vectors, with norms and inner products scaled by the same factor. The resulting codes have 256, 576 and 512 points. The first attains the eleven-coordinate capacity; the second follows directly from four colours of two one-factorizations. In dimension 37, thirteen extra supports contribute 1664 points and the replacement adds 128 more.

Cross-shell completion in dimension 38. The base lifting uses all minimal Leech directions but leaves an empty equator. For a norm-48 vector v and a norm-32 vector u in the same integer Leech lattice, the minimum norm gives $\|v \pm u\|^2 \geq 32$, hence $|v \cdot u| \leq 24$. This removes the entire old shell from the subset-selection constraints. The remaining task is to find mutually compatible new lines. The 144-line construction contributes 288 points, giving 591900.

Image unions in dimensions 49–55. An enlarged class of 7077 minimal lines in P_{48p} produces further compatible classes by isometries. The relevant size is their union: a repeated line is assigned once. With S union lines and t distinct antipodal tail lines, the construction has $52416000 + 2S + 2t$ points. One lifting principle thus supplies seven bounds, with separately specified image families and an explicit decomposition of the gain from class enlargement and image selection.

Target moments in dimensions 43 and 45. The minimal shell of an extremal 48-dimensional lattice is a spherical 11-design. Its moments constrain all admissible projection fibres. For every extremal even unimodular lattice of rank 48 containing $\sqrt{3}E_8$, a rational identity determines exactly 2553792 minimal vectors perpendicular to the specified D_5 child and each of its Weyl conjugates. In dimension 45 the moments, together with the norm-eight

section constraints, give the exact identity

$$f(e_1) + f(e_2) + f(e_3) = 7377408 + 9f(-2, -2, 3).$$

The final fibre is a neighbourhood difference in an anchor graph. Its size is $368 - c$, where c counts common neighbours of the selected nonadjacent pair. The projected configuration therefore has exactly $7380720 - 9c$ points; the supplied anchor attains its maximum at $c = 70$, giving 7380090. In both dimensions, a small set of geometric constraints determines a large section count.

1.4 Mathematical context

The Leech lifting framework of Cohn, Jiao, Kumar and Torquato [7, Theorem 7.5], and its subsequent refinements, combine a mother direction with a low-dimensional tail. PackingStar provides the 496-point blocks used in the four-triangle construction [21, 22]. The 25-dimensional deformation starts from Kravatskiy’s 197579-point configuration; the 27-dimensional two-layer input is Lindow’s 201566-point construction [15, 16]. The empty-equator lifting in dimension 38 and the 7069-line comparison class in dimension 48 also come from [15]. The new objects are the coherent motion and repairs, the second-layer allocation, the added shell lines, and the enlarged class with its selected image families.

The code constructions use the layers of Edel, Rains and Sloane [6], the dense code of Cheng–Sloane [23], and constant-weight codes and tables [17, 20, 24]. Local replacement has a direct precursor in the free-zone method of Takhanov–Yun [25, Section V]. The 51-block weight-four source is the classical Kalbfleisch–Stanton design [26]. Here the new supports, coordinate pairings, signed selections and boundary embeddings produce the larger configurations.

Lattice sections and the antipode construction originate in Conway–Sloane [4, 5]. The public comparisons for 43 and 45 dimensions are the counts of Sun–Wang [18, 19]; the embedded E_8 analysis of Dorofeev, Sun and Wang [27] supplies the parent geometry. Spherical-design theory [28, 29] provides the moment identities. The calculations here select the target statistics and finite witnesses that yield the stated counts.

1.5 Reading the constructions

The report follows the constructions from their governing geometric relations to the explicit finite objects. The final appendices specify coordinates, checking procedures and research development. The accompanying repository contains the bilingual dimension accounts, source data and programs, so that a reader can reconstruct the objects, check the proofs’ finite hypotheses and vary the construction choices.

2 Coordinated shell motion in dimension 25

A coordinated motion of a complete contact layer enlarges the 197579-point configuration supplied in the public [KissingNumbers/dim25-lens-heads](#) package. The construction moves 552 retained equatorial points, adjoins one point, and replaces two cap points. The motion preserves every internal product of the moved layer.

Lemma 2.1 (Coordinated chord motion). *Let n be a unit vector, let $h > 0$, and let $r_i \perp n$ satisfy $\|r_i\|^2 = \rho - h^2$. Put $u_i = r_i - hn$ and $w_i = r_i + hn$. For real c, d with $c^2 + d^2 = h^2$, set $W_i = (r_i + cn, d)$. Then $\|W_i\|^2 = \rho$ and $W_i \cdot W_j = w_i \cdot w_j$. If $z \cdot u_i \leq \rho/2$ and $z \cdot w_i \leq \rho/2$, then $(z, 0) \cdot W_i \leq \rho/2$.*

Proof. The first two claims follow from $r_i \perp n$ and $c^2 + d^2 = h^2$. Since $|c| \leq h$, the horizontal component of W_i is $\frac{h-c}{2h}u_i + \frac{h+c}{2h}w_i$, a convex combination. The last inequality follows by linearity. $\square$

For an axial candidate $Q = (\alpha n, \beta)$ with $\alpha^2 + \beta^2 = \rho$, every moved point has the same product $\alpha c + \beta d$ with Q . Simultaneous contact is possible exactly when $h^2 \geq \rho/4$, at the circle–line intersections

$$(c, d) = \frac{1}{2}(\alpha, \beta) \pm \sqrt{\frac{h^2 - \rho/4}{\rho}}(-\beta, \alpha). \quad (2.1)$$

For a finite nonempty moved layer, compatibility with a fixed point (z, s) is exactly the half-plane constraint

$$c\langle n, z \rangle + ds \leq \rho/2 - \max_i \langle r_i, z \rangle. \quad (2.2)$$

The candidate must also be compatible with the fixed points. In the construction below, two cap replacements resolve the remaining conflicts.

Proposition 2.2 (Finite boundary test). *Let $H_1, \dots, H_m$ be closed half-planes in the (c, d) -plane, and let F consist of $(h, 0)$ together with every intersection of the circle $c^2 + d^2 = h^2$ with their boundary lines. Constant inequalities contribute no boundary points. Then $|F| \leq 2m + 1$, and the circle meets $\bigcap_j H_j$ if and only if some point of F belongs to every H_j . More generally, designate some inequalities as mandatory and assign nonnegative deletion costs to the others. The minimum cost of violated inequalities, when the mandatory constraints admit a circle point, is attained at a feasible point of F .*

Proof. Remove all boundary intersections from the circle. On each remaining open arc, every inequality has constant truth value. At an endpoint, every inequality that held on the arc still holds, because its half-plane is closed. Moving there preserves mandatory inequalities and cannot increase deletion cost. If there are no boundary intersections, all truth values are constant on the circle, and $(h, 0)$ suffices. $\square$

For a nonzero normal $A = (A_1, A_2)$, the boundary $A \cdot (c, d) = B$ contributes points exactly when $B^2 \leq h^2 \|A\|^2$, namely

$$\frac{BA \pm \sqrt{h^2 \|A\|^2 - B^2}}{\|A\|^2} (-A_2, A_1).$$

Rational half-plane data and rational h^2 therefore give an exact test on at most $2m + 1$ quadratic-algebraic candidates. This minimizes deletions; the existence of replacements is a separate requirement.

Normalization and input. All points have squared norm 4; the kissing condition is an inner product at most 2 between distinct points. Write the input configuration as

$$C_0 = \{(z, 0) : z \in L \setminus D\} \cup \{(x_i, \pm 1) : 0 \leq i < 1016\} \cup \{P_0, N, S\},$$

where L is the 196560-vector minimal shell of the Leech lattice, $|D| = 1016$, $P_0 = (P, 0)$, and $N = (0, 2)$, $S = (0, -2)$. The head indices refer to rows of the source `heads_X.npy`; the mathematical values of its 44 rational heads are preserved as exact rational numbers in `baseline-heads.json`.

Let v be the following squared-norm-six lattice vector associated with the first complete block, and put $n = v/\sqrt{6} = -P/2$. An exact integer description of this direction is

$$v = \frac{1}{\sqrt{8}}(-5, -1, -1, -1, -1, 1, -1, 1, -1, -1, -1, 1, 1, -1, -1, -1, 1, 1, -1, 1, -1, -1, 1, -1).$$

The complete block is $U = \{u \in L : \langle u, v \rangle = -3\}$, of size 552. Every $u \in U$ belongs to D . For each such u , the point $w = u + v$ is minimal, lies outside D , and satisfies $\langle u, v \rangle = -3$ and $\langle u, w \rangle = 1$. The corresponding old cap points have horizontal coordinate $u + tv$, with $t = (3 - \sqrt{3})/6$. All these input identities are checked by the shell and owner data.

The modification. For $u \in U$ define

$$r = u + \sqrt{3/2}n = w - \sqrt{3/2}n, \quad a = \frac{\sqrt{3}}{2} + \frac{\sqrt{2}}{4}, \quad b = \frac{\sqrt{6} - 2}{4}.$$

The values (a, b) are the positive choice in (2.1) for $h^2 = 3/2$, $\rho = 4$, and $(\alpha, \beta) = (\sqrt{3}, -1)$. Then $r \perp n$, $\|r\|^2 = 5/2$, and the old cap points over u are $(r - n/\sqrt{2}, \pm 1)$. Replace every $(w, 0)$ by

$$W(r) = (r + an, b),$$

and adjoin

$$Q = (\sqrt{3}n, -1) = (V/4, -1),$$

where $V = \sqrt{8}v$ is the integer vector printed above. In particular, the new point Q is rational and has squared norm $48/16 + 1 = 4$. Finally replace the upper point $(x_{984}, 1)$ and the lower point $(x_{1008}, -1)$ by R_+ and R_- , respectively. The certificate specifies two integer vectors and defines

$$R_+ = \frac{2k}{\sqrt{1007176}}, \quad R_- = \frac{2\ell}{\sqrt{10209}},$$

where k, ℓ are the integer rows in `constructions/d25/repair-points.json`, of squared norms 1007176 and 10209. The opposite old cap points $(x_{984}, -1)$ and $(x_{1008}, 1)$ are retained.

Proposition 2.3 (Two necessary cap changes). *Keep Q fixed and move the 552 points by $W(r) = (r + cn, d)$, with $c^2 + d^2 = 3/2$. Any compatible configuration obtained this way must change at least two original cap points. The specified construction attains this minimum.*

Proof. The exact cap data give $Q \cdot (x_{1008}, -1) > 231/100 > 2$, so this lower cap must change independently of (c, d) . If it were the only changed cap, all first-block cap pairs $(r - n/\sqrt{2}, \pm 1)$ would remain. Their products with their own moved points require $-c/\sqrt{2} + |d| \leq -1/2$. Together with the circle equation this gives $c > 0$ and $|d| \leq 1/3$. On this interval $c = \sqrt{3/2 - d^2}$ and $\sqrt{3}c - d$ is strictly decreasing. Compatibility with Q therefore forces $d \geq b$, since $\sqrt{3}a - b = 2$.

Put $\nu = \langle n, x_{984} \rangle$ and $M = \max_r \langle r, x_{984} \rangle$. The largest moved-layer product with the upper 984 cap is $g(d) = M + \nu\sqrt{3/2 - d^2} + d$. Since $\|x_{984}\| = \sqrt{3}$,

$$g'(d) = 1 - \frac{\nu d}{\sqrt{3/2 - d^2}} \geq 1 - \frac{\sqrt{6}}{5} > 0 \quad (b \leq d \leq 1/3).$$

The rational head and shell data, evaluated with outward-rounded intervals, give $g(b) > 201/100 > 2$. Thus this upper cap must also change, a contradiction. The motion-obstruction

checker reproduces both strict checks. The subsequent pair verification proves that the specified two replacements suffice. $\square$

Pair verification. Lemma 2.1 gives the moved-point norms, internal products and compatibility with the unchanged equator. The identities

$$a^2 + b^2 = \frac{3}{2}, \quad \sqrt{3}a - b = 2, \quad -\frac{a}{\sqrt{2}} + b = -\frac{3}{4}$$

give $Q^2 = 4$ and $\langle W(r), Q \rangle = 2$.

The shell has integral $\langle v, z \rangle \in \{-3, -2, -1, 0, 1, 2, 3\}$. The negative extreme layer is deleted and the positive extreme layer is exactly the moved set. Thus every unchanged equatorial point obeys

$$\langle Q, (z, 0) \rangle = \frac{\langle v, z \rangle}{\sqrt{2}} \leq \sqrt{2} < 2.$$

The shell checker verifies these layer identities in integers. Products of W with (P_0, N, S) are $(-2a, 2b, -2b)$; those of Q are $(-2\sqrt{3}, -2, 2)$. Against its own first-block caps, $W(r)$ has products $7/4$ and $11/4 - \sqrt{6}/2 < 2$; different block indices give smaller products because $\langle r_i, r_j \rangle \leq 1/2$.

The four remaining groups are: moved points against the 2030 unchanged caps; Q against those caps; and each of R_+, R_- against all other final points. Their outward-rounded rational upper bounds are

$$1.999372, \quad 1.965926, \quad 1.994722, \quad 1.987522.$$

The checker uses 192-bit Arb intervals [14] and verifies both repair norms exactly. Unchanged pairs retain the input inequalities.

Theorem 2.4. *The specified configuration consists of 197580 distinct points on the radius-2 sphere in $\mathbb{R}^{25}$ with pairwise inner products at most 2. Consequently the kissing number in dimension 25 is at least 197580.*

Proof. The pair checks above, together with the integer or rational baseline checks, exhaust all pairs. All squared norms are exactly 4. The moved layer and repairs replace equally many old points; only Q increases the count:

$$|C| = 197579 - 552 - 2 + 552 + 2 + 1 = 197580.$$

No duplicate can occur: two equal norm-4 vectors would have inner product 4, contradicting the verified bound 2 for their distinct indices. $\square$

3 Labelled lifts: count the directions that remain

Let $C \subset S^{D-1}$ be a kissing configuration. Split the larger space as $\mathbb{R}^D \oplus \mathbb{R}^k$. Choose pairwise distinct nonzero tails t_i of norm less than one and subsets $A_i \subseteq C$. Write $a_i = \sqrt{1 - \|t_i\|^2}$ and $U = \bigcup_i A_i$. Retain $(x, 0)$ for $x \in C \setminus U$, insert $(a_i x, t_i)$ for $x \in A_i$, and add a pure-tail kissing code $Y \subset S^{k-1}$.

Each original direction is deleted once, giving

$$|X| = |C| - |U| + \sum_i |A_i| + |Y|.$$

For distinct labelled pairs $(x, i) \neq (x', j)$ and pure tails $y \in Y$, the remaining conditions are

$$a_i a_j \langle x, x' \rangle + \langle t_i, t_j \rangle \leq \frac{1}{2}, \quad \langle t_i, y \rangle \leq \frac{1}{2}.$$

3.1 Multiplicity and the four-triangle construction

If every tail has squared norm $h^2 < 1/2$, repeated copies of one direction require

$$\langle t_i, t_j \rangle \leq h^2 - \frac{1}{2}.$$

For m copies, the squared norm of the sum of their tails is at most

$$\frac{m}{2}(1 - m(1 - 2h^2)).$$

Nonnegativity gives the latitude bound. Strictly negative mutual tail products also imply affine independence: an affine dependence would equate positive combinations of disjoint sets whose mutual product is negative. Thus

$$m \leq r = \min\{k + 1, \lfloor (1 - 2h^2)^{-1} \rfloor\}.$$

Both bounds are sharp for one direction, by regular-simplex tails. If label i has capacity α_i , the count is consequently at most $|C| + |Y| + (1 - 1/r) \sum_i \alpha_i$.

More precisely, lifting a nonempty block B at the same $r \geq 2$ tails at some $0 < h^2 < 1/2$ is possible exactly when $k \geq r - 1$ and its distinct products are at most $1/(r + 1)$. Necessity follows from the multiplicity bound and the decreasing same-label threshold $(1/2 - h^2)/(1 - h^2)$. A regular simplex at $h^2 = (r - 1)/(2r)$ attains both bounds, giving $|C| + (r - 1)|B|$ points without pure tails. At $h^2 = 1/3$, this is triple reuse on a zero-sum triangle.

The twelve roots of D_3 , scaled to tail norm $1/\sqrt{3}$, split into four zero-sum triangles. Four disjoint 496-point Leech blocks, each carried on one triangle, give

$$196560 - 4(496) + 12(496) + 12 = 200540.$$

Changing multiplicities from $(3, 3, 2, 2, 2)$ to $(3, 3, 3, 3)$ preserves 496 additional equatorial points with the same lifted count. The blocks are published PackingStar data; their tail assignment changes.

For this fixed twelve-label system with at most α directions per label, the count is at most $196560 + 8\alpha + 12$. Thus the construction attains the allocation capacity at $\alpha = 496$.

3.2 The later 201,567-point construction

The later construction extends a public two-layer configuration. Its first layer, taken unchanged from Lindow's 201,566-point contribution, has 2,258 integer heads of squared norm 192. Of these, 2,090 delete one Leech minimal vector each, and 168 delete none. Each head produces three caps.

Use the integer Leech convention in which minimal vectors have squared norm 32 and the kissing threshold is 16. A first-layer head y produces

$$(y/3, 4v/\sqrt{3}),$$

where v runs through its assigned D_3 triangle. The exact head thresholds are 48 on one triangle and 96 between different triangles.

A second-layer owner u is a Leech minimal vector assigned to one of three coordinate lines. Replace its equatorial point by

$$(\sqrt{3}u/2, \pm\sqrt{8}e_j).$$

The supplied 311-owner witness has line sizes 105, 103, and 103. Every first-layer head satisfies $y \cdot u \leq 32$. Two owners on one line have product at most eight; different lines have product at most sixteen. Owners are distinct and do not overlap the first-layer deletion set.

For example, the first/second-layer product is at most

$$\frac{\sqrt{3}}{6} 32 + \frac{4\sqrt{8}}{\sqrt{3}} = \frac{16\sqrt{3} + 8\sqrt{6}}{3} < 16.$$

The second-layer/pure-tail product is at most $\sqrt{8}\sqrt{32} = 16$, including equality. The two caps of one owner also have product sixteen. These checks give

$$196560 - 2090 - 311 + 3(2258) + 2(311) + 12 = 201567.$$

Replacing the public input's 310 second-layer owners by the new selection of 311 adds one point: each owner replaces one equatorial point by two caps.

4 Layered codes, joint exchanges and signed constructions

Dimensions 32–34 and 39 enlarge the support family of one layered construction; dimensions 35–37 also replace complete sign bundles by different finite sets.

4.1 Compatibility of the layers

The Edel–Rains–Sloane construction [6] places three coordinate patterns on the same unit sphere. Let $n \geq m \geq 32$, let $T \subset \{0, 1\}^m$ be a binary code, and let $\mathcal{B}$ be a family of eight-element subsets of $\{0, \dots, n-1\}$. The dense, middle and root layers are

$$\begin{aligned} X_T &= \left\{ \frac{1}{\sqrt{m}} ((-1)^{c_0}, \dots, (-1)^{c_{m-1}}, 0, \dots, 0) : c \in T \right\}, \\ X_{\mathcal{B}} &= \left\{ \frac{1}{\sqrt{8}} \sum_{i \in B} \varepsilon_i e_i : B \in \mathcal{B}, \varepsilon_i \in \{-1, 1\}, \prod_{i \in B} \varepsilon_i = 1 \right\}, \\ X_R &= \left\{ \frac{\pm e_i \pm e_j}{\sqrt{2}} : 0 \leq i < j < n \right\}. \end{aligned}$$

Each support carries $2^7 = 128$ even-parity sign patterns. The layers have respectively m , eight and two nonzero coordinates, so they are disjoint. Their union contains

$$N = |T| + 128|\mathcal{B}| + 2n(n-1) \tag{4.1}$$

points.

Proposition 4.1 (Three-layer construction [6]). *If T has minimum Hamming distance at least $\lceil m/4 \rceil$ and distinct supports satisfy $|B \cap B'| \leq 4$, then the union is a kissing configuration in dimension n .*

Proof. Two dense-layer points have product $1 - 2d_H(c, c')/m \leq 1/2$. Distinct even-parity patterns on the same support differ in at least two positions, giving at most $1 - 4/8 = 1/2$. Different supports give at most $|B \cap B'|/8 \leq 1/2$. Distinct roots either share at most one coordinate with the same sign, or have the same support and differ in at least one sign; their product is also at most $1/2$.

There are three remaining cross-layer cases. Dense-middle products are at most $8/(\sqrt{m}\sqrt{8}) = \sqrt{8/m} \leq 1/2$; dense-root products are at most $2/(\sqrt{m}\sqrt{2}) = \sqrt{2/m} \leq 1/2$; middle-root products are at most $2/(\sqrt{8}\sqrt{2}) = 1/2$. These cover all distinct pairs. $\square$

The identity $d_H(1_B, 1_{B'}) = 16 - 2|B \cap B'|$ makes the support condition minimum distance eight in a constant-weight code. Each extra support therefore adds 128 spherical points [17, 20, 24].

4.2 Joint exchanges of supports

For a candidate b , put $F(b) = \{B \in \mathcal{B} : |B \cap b| > 4\}$. If Y is disjoint from the old family and its internal intersections are at most four, then

$$\begin{aligned} \mathcal{B}' &= (\mathcal{B} \setminus F(Y)) \cup Y, & F(Y) &= \bigcup_{b \in Y} F(b), \\ |\mathcal{B}'| - |\mathcal{B}| &= |Y| - |F(Y)|. \end{aligned} \tag{4.2}$$

The cost is a union size, not the sum of the individual conflict counts. The stored final exchange in dimension 33 removes nine supports and adds ten; in dimension 34 it removes six and adds seven; an exchange in dimension 37 removes 37 and adds 40. The data contain the parent families and actual deletions and additions, so these unions can be reconstructed.

There is also a direct discrete formulation. For a finite candidate set $\mathcal{C}$, let $z_b \in \{0, 1\}$ select a candidate and let $u_B \in \{0, 1\}$ delete an old support. Impose $z_b + z_{b'} \leq 1$ for incompatible candidates and $z_b \leq u_B$ for $B \in F(b)$, and maximize

$$\sum_{b \in \mathcal{C}} z_b - \sum_{B \in \mathcal{B}} u_B.$$

An optimum deletes only actual blockers, so this is exactly the gain in (4.2). If the entire pool $\mathcal{C}$ is compatible, let H be its bipartite conflict graph with $\mathcal{B}$. Then

$$\max_{Y \subseteq \mathcal{C}} (|Y| - |F(Y)|) = |\mathcal{C}| - \nu(H),$$

where $\nu(H)$ is the maximum matching size. Any matching leaves at least $|Y| - |F(Y)|$ candidates unmatched; alternating paths from the unmatched candidates of a maximum matching attain equality. The saved 33-, 34-, 37-, 38- and 39-dimensional insertion pools have matching sizes 9, 6, 37, 6, 14, certifying optimal subexchange gains 1, 1, 3, 1, 1 within those pools. This does not optimize over all possible supports.

The blocking sets also organize candidate generation. For a chosen integer f , consider eight-subsets outside $\mathcal{B}$ with $|F(b)| \leq f$. If the pool includes all such subsets, it covers every positive

exchange with at most $f + 1$ additions: for $b \in Y$, $|F(b)| \leq |F(Y)| < |Y| \leq f + 1$. The threshold therefore describes a specific exchange neighbourhood rather than a limit on the final code.

Candidate conflicts admit a compact representation. Two eight-subsets intersect in at least five positions exactly when they share a five-subset. Each candidate occupies $\binom{8}{5} = 56$ such subsets, and the constraints are

$$\sum_{b \in \mathcal{C}: Q \subset b} z_b \leq 1 \quad \left(Q \in \binom{\{0, \dots, n-1\}}{5} \right).$$

Add constraints at repeated five-subsets until the selected family is compatible; accept positive gain and recompute blockers. In dimension 34, 10715 candidates with at most three blockers yield the seven-for-six exchange.

The following parameters give direct three-layer constructions. The dimension-37 support family was enlarged from 2843 to 2845 and then modified by a signed replacement, whose final count is given below.

Table 2 Parameters of the direct three-layer constructions.

n	Dense length m	$ T $	$ \mathcal{B} $	Points
32	32	131072	1676	347584
33	32	131072	1803	363968
34	32	131072	1960	384196
39	39	327680	3383	763668

For example, $131072 + 128 \cdot 1676 + 1984 = 347584$ in dimension 32. The gains of 15 and 24 supports in dimensions 33 and 34 give respectively 1920 and 3072 points over the comparison constructions.

The length-32 layer uses the Cheng–Sloane [32, 17, 8] code [23], whose generator expands to $2^{17} = 131072$ words. Dimension 39 uses $T = \bigcup_{a=1}^{10} (H + r_a)$, where H has dimension 15 and minimum weight 12. Enumeration of the supplied kernel and representatives gives

$$\min_{h \in H} \text{wt}(h + r_a + r_b) = 10$$

for all 45 distinct coset pairs. The cosets are disjoint, with $10 \cdot 2^{15} = 327680$ words and minimum distance ten, as required. The kernel is obtained by shortening an extended quadratic-residue code; the programs compare generator row spaces and check the actual distances.

The listed dimension-39 comparison has the same dense-layer size. Increasing its 3324 supports to 3383 adds $128(3383 - 3324) = 7552$ points.

4.3 Hadamard transfer and local replacement

Local replacement changes the signs as well as the supports.

Proposition 4.2 (Paired Hadamard transfer). *Let $\mathcal{A}$ be a family of four-element supports with intersections at most two, transverse to a fixed coordinate pairing. Applying a Hadamard matrix to each pair in all signed versions of $\mathcal{A}$ gives $16|\mathcal{A}|$ weight-eight vectors of squared norm eight and distinct-pair products at most four.*

Proof. Transversality means meeting each coordinate pair at most once. Apply

$$H = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

to each coordinate pair. Both $(\varepsilon, 0)$ and $(0, \varepsilon)$ become two nonzero coordinates, so weight four becomes weight eight. Distinct input vectors have product at most two: different supports satisfy the intersection condition, and distinct sign patterns on the same support differ in at least one position. Since $H^\top H = 2I$, the images have squared norm eight and pairwise products at most four. Invertibility preserves their number. $\square$

After division by $\sqrt{8}$, these vectors form an internally compatible spherical set Q . Compatibility and disjoint insertion require intersections at most four with every retained complete bundle: signs on smaller intersections extend to either parity, and equal supports give either a duplicate or a product above $1/2$. The dense and root cross bounds remain $1/2$, independent of the new signs' parity.

The free-zone replacement principle appears in Takhanov–Yun [25, Section V, Definition 5 and Lemma 4] for signed weight-four codes. Here it is applied to weight-eight regions inside the ERS middle layer. In dimensions 35 and 36, boundary compatibility can be secured before choosing Q . Take a coordinate region W , remove the old supports contained in it, and require $|B \cap W| \leq 4$ for every retained support. Any new weight-eight vector on W is then compatible with the retained middle layer. Only its local internal products remain to be controlled. The regions are $W = \{12, \dots, 22\}$ and $W = \{12, \dots, 23\}$, respectively, with zero-based coordinates. They contain one and three old supports, fixing the deletion costs at 128 and 384 points.

Direct search among signed weight-eight vectors in eleven and twelve coordinates has respectively $2^8 \binom{11}{8} = 42240$ and $2^8 \binom{12}{8} = 126720$ candidates. Weight-four transfer replaces this search with the choice of a combinatorial source and a coordinate pairing, while preserving an algebraic proof of the inner-product bounds. Dimension 35 starts with a Steiner quadruple system $S(3, 4, 10)$: each triple belongs to exactly one block, giving $\binom{10}{3}/4 = 30$ blocks with pairwise intersections at most two. Enumerating the 945 perfect matchings on ten coordinates yields sixteen transverse blocks. For dimension 36 we use the 51-block construction of Kalbfleisch–Stanton [26, Theorem 7], also described in [25, Section V, Construction 1]. Split twelve coordinates into two sets of six and use the same five one-factors of K_6 in each half. For every colour, the nine choices of a left and a right edge give a four-element block, producing 45 crossing blocks. Add, in each half, the complements of the three edges of the first colour to obtain six further blocks.

The resulting 51 blocks still have intersections at most two. Same-colour crossing blocks sharing one edge intersect in two points; different colours share at most one endpoint in each half. Internal blocks in the same half intersect in two points; those in different halves are disjoint. An internal block meets a crossing block in at most two points.

Proposition 4.3 (A 576-point local code). *The paired Hadamard image of 36 transverse blocks from the two one-factorizations is a 576-point weight-eight code in twelve coordinates.*

Proof. Choose the three edges of one colour as the coordinate pairs in each half. Each edge of the other four colours meets every chosen pair at most once. Its union with a same-colour edge from the other half is therefore transverse. There are $4 \cdot 3 \cdot 3 = 36$ such blocks; their intersections are at most two by the preceding argument. Proposition 4.2 gives $36 \cdot 16 = 576$ points. The six internal blocks are unnecessary for this transverse subfamily. $\square$

In the supplied local coordinate order, the chosen pairs are

$$(0, 1), (2, 4), (3, 5), (6, 7), (8, 10), (9, 11).$$

They form the same colour in the two halves. The direct construction with these pairs regenerates exactly the stored 576 vectors. The accompanying program also enumerates all 945 and 10395 matchings of the two source codes, recovering the 256-point and 576-point sets.

The compatible region in dimension 35 has eleven coordinates; ten form five pairs and the remaining coordinate is zero. Dimensions 36 and 37 use twelve coordinates each, with 36 and 32 transverse supports.

Table 3 Signed replacements. The support count precedes the deletion of bundles.

Dimension	Original supports	Bundles removed	Points added	Net gain
35	2157	1	256	128
36	2742	3	576	192
37	2845	3	512	128

Combining the support choices and replacements gives

$$K(35) \geq 131072 + 128(2157 - 1) + 256 + 2(35)(34) = 409676,$$

$$K(36) \geq 131072 + 128(2742 - 3) + 576 + 2(36)(35) = 484760,$$

$$K(37) \geq 131072 + 128(2845 - 3) + 512 + 2(37)(36) = 498024.$$

Relative to the comparison with 2832 supports, dimension 37 gains 1664 points from thirteen extra supports and 128 from the replacement, totalling 1792. Dimensions 35 and 36 keep their original support counts; their gains come entirely from the signed choices.

The local choice in dimension 35 also attains an exact capacity bound. Let $M(11)$ be the largest number of weight-eight vectors in $\{0, \pm 1\}^{11}$ with distinct-pair products at most four. Fix a full sign string $\sigma \in \{\pm 1\}^{11}$. Two vectors agreeing with σ on their supports would intersect in at least five positions and have product at least five. Thus each sign string extends at most one selected vector. Each vector, in turn, has eight extensions on its three zero coordinates. Counting gives

$$8|Q| \leq 2^{11}, \quad M(11) \leq 256.$$

The construction attains equality, so $M(11) = 256$.

The four-colour construction extends to two halves of size $2t$, giving $32t^2(t - 1)$ points for every $t \geq 2$. This is sharp among complete-sign transverse $2 + 2$ transfers: for each edge in one half, its partner edges in the other half must form a matching of size at most t . Since each half has $2t(t - 1)$ available edges, there are at most $2t^2(t - 1)$ source blocks. Equality requires every left and right fibre to be perfect, and synchronized one-factorizations attain it.

For twelve coordinates, fixed-pair boundary restrictions have an exact capacity. A type chooses two coordinate pairs in each half; let $H \subseteq K_{3,3}$ record the allowed types, each allowing all sixteen axis choices. Let D_H have two copies of each edge of H , joined across copies when distinct edges share an endpoint. The maximum number of source blocks is

$$\beta(H) = 8|E(H)| - 4\nu(D_H).$$

A matching in D_H bounds the type counts; a minimum vertex cover constructs an attaining family with zero, four or eight blocks per type. Thus the maximum transferred size is $16\beta(H)$, verified for all 2^9 type relations. Exactly one forbidden type gives 512 points. For the fixed dimension-37 pairing and three deleted bundles, the ten divisions into two halves have capacities 512 (four), 448 (four) and 384 (two). The supplied 512-point witness fits none of these divisions: its ten image types exceed the nine available in a $2 + 2$ division. Neither this capacity nor the 576-point construction bounds arbitrary twelve-coordinate signed codes.

The finite checks use the coordinate order, pairs and four-subsets specified in the data. In dimension 37, a support string numbered from one at the left has coordinate i at integer-mask bit $37-i$. The program regenerates all 512 points from 32 transverse supports and six coordinate pairs, checks equality with the stored replacement, and checks intersections with every retained support.

The next construction instead uses equatorial directions left available by a large lattice lifting.

5 Equatorial completion of a Leech-lattice lifting

The signed replacements balance additions against deletions. Here a large lifting leaves an unused equator. We seek directions that can be inserted directly, compatible both with one another and with the entire lifted configuration. A set of 144 such lines contributes 288 points, increasing the lower bound from 591612 to 591900.

5.1 The lifted configuration and its equatorial constraints

Let $\Sigma \subset \mathbb{Z}^{24}$ be the squared-norm-32 minimal Leech shell, with 196560 vectors. The supplied extended Golay code generates 1104 vectors with two entries ± 4 ; 97152 vectors with entries ± 2 on a weight-eight support and even sign parity; and 98304 vectors obtained from signs $(-1)^{c_i}$ by reversing one chosen sign and multiplying that entry by three. The Golay generator has rank 12 and weight enumerator $1 + 759z^8 + 2576z^{12} + 759z^{16} + z^{24}$. Distinct vectors in this standard shell model have inner product at most 16 [2].

Write $\mathbb{R}^{38} = \mathbb{R}^{24} \oplus \mathbb{R}^{14}$. The public 591,612-point construction uses every minimal Leech direction in a cap layer [15], but leaves the equatorial subspace available for other directions. The new construction exploits this distinction.

Partition the 98,280 antipodal lines of Σ into 644 classes whose representatives satisfy $|u \cdot u'| \leq 8$ within each class. In $\mathbb{R}^{14}$ take 1932 vectors of squared norm 8 and distinct inner products at most 4, partitioned into 644 zero-sum triangles. Each triangle has pairwise products -4 . Pair a head class with a triangle and form six points per head line:

$$\left(\frac{\varepsilon u}{\sqrt{48}}, \frac{d}{\sqrt{24}} \right), \quad \varepsilon \in \{1, -1\},$$

with d ranging over the assigned triangle. These are unit vectors. Equal heads with different tails give product $32/48 - 4/24 = 1/2$; different heads with an equal tail give at most $8/48 + 8/24 = 1/2$. Points from different classes give at most $16/48 + 4/24 = 1/2$, and the other cases give smaller bounds. The cap layer has $6 \cdot 98280 = 589680$ points.

Both signs on a head line are included. Opposite heads contribute $-32/48$, so even with identical tails their total product is only $-1/3$.

Add a pure tail layer $(0, Rd/\sqrt{8})$, using the same 1932 directions and an exact rational rotation $R = Q/D$. Its integer data satisfy

$$QQ^\top = D^2I, \quad |d \cdot Qd'| \leq \sqrt{48}D.$$

The first identity preserves the pure tail products; the second bounds cap–tail products by $1/2$. Integer square roots suffice to check the latter exactly. This gives the base count $589680 + 1932 = 591612$.

5.2 Compatible lines in the second shell

We use the norm-48 shell of the same Leech lattice. If $\mathcal{G}$ is the supplied Golay code, its integer coordinate model is

$$\mathcal{L} = \left\{ x \in \mathbb{Z}^{24} : \begin{array}{l} x_i \equiv m \pmod{2} \text{ for some } m \in \{0, 1\}, \\ ((x_i - m)/2 \bmod 2)_{i=1}^{24} \in \mathcal{G}, \\ \sum_i x_i \equiv 4m \pmod{8} \end{array} \right\}. \quad (5.1)$$

Linearity of $\mathcal{G}$ and its all-one word make this set a lattice. Its minimum squared norm is 32. Indeed, for even coordinates a nonzero Golay residue has at least eight entries of absolute value two. A zero residue makes all coordinates multiples of four, and the sum congruence excludes a single entry of absolute value four. For odd coordinates, a vector of squared norm below 32 would have every entry ± 1 ; the doubly even Golay weight then makes its coordinate sum zero modulo eight, contrary to the last congruence. The minimal vectors described above attain 32.

Lemma 5.1 (Compatibility of the second shell). *If $v \in \mathcal{L}$ has squared norm 48 and $u \in \mathcal{L}$ has squared norm 32, then $|v \cdot u| \leq 24$.*

Proof. The distinct norms imply $v \pm u \neq 0$. Their squared norms are $80 \pm 2v \cdot u \geq 32$, which gives the claimed bound. $\square$

Although all minimal lines are used, equatorial points can come from directions of another length. Squared norm 48 gives the same normalization denominator as the cap heads. The new compatibility conditions consequently become integer-product checks with threshold 24. The 144 supplied vectors satisfy all three congruences in (5.1). It remains to check their mutual products:

$$\|v_i\|^2 = 48, \quad |v_i \cdot v_j| \leq 24 \quad (i \neq j), \quad |v_i \cdot u| \leq 24 \quad (u \in \Sigma).$$

Proposition 5.2 (Equatorial completion). *Adding the 288 points $(\pm v_i/\sqrt{48}, 0)$ to the cap and pure-tail layers gives a 591900-point kissing configuration in dimension 38.*

Proof. The 288 new equatorial points $(\pm v_i/\sqrt{48}, 0)$ are internally compatible. Their products with a cap are $\pm v_i \cdot u/48 \leq 1/2$, and they are orthogonal to the pure tails. No base point lies on the equator, so no deletion is needed. Thus

$$K(38) \geq 591612 + 2 \cdot 144 = 591900.$$

$\square$

Lemma 5.1 leaves a finite independent-set problem: norm-48 lattice lines conflict exactly when $|v \cdot v'| > 24$. The search produced 142 lines and then 144. The data use the base construction's Golay basis, positive first nonzero coordinates and lexicographic order; the verifier also recomputes every product with Σ . Geometrically, the shortened cap heads permit $|z \cdot c| \leq \sqrt{3/8}$ for $c \in \Sigma/\sqrt{32}$, a wider angle range than direct insertion into the unit mother shell.

5.3 Contacts and limits of the shell choice

For minimal lattice heads of squared norm m scaled by $1/\sqrt{q}$, the same minimum-norm argument gives automatic compatibility for equatorial vectors of squared norm $m < b \leq q$. Two added vectors of norms b, c still require

$$|v \cdot w| \leq \sqrt{bc}/2 \iff \min\{\|v - w\|^2, \|v + w\|^2\} \geq b + c - \sqrt{bc}.$$

The lattice minimum alone is smaller than this threshold when $b, c > m$. Here the automatic interval $32 < b \leq 48$ contains only the norm-48 shell, since squared norms in this Leech scaling are multiples of 16.

The next shell is impossible for the unchanged base. Both this obstruction and the norm-48 contact count follow from one calculation. Rescale to minimum four. For $v^2 = s$, the spherical 11-design property of the minimal shell L [29] gives

$$M_{2j}(s) = \sum_{x \in L} (v \cdot x)^{2j} = \frac{196560(4s)^j(2j-1)!!}{24 \cdot 26 \cdots (24+2j-2)} \quad (1 \leq j \leq 5).$$

For $s = 6$ and 8, the products are integers of absolute value at most 3 and 4, respectively. Polynomials vanishing at the smaller values give

$$\frac{M_6(6) - 5M_4(6) + 4M_2(6)}{720} = 552, \quad \frac{M_8(8) - 14M_6(8) + 49M_4(8) - 36M_2(8)}{40320} = 46.$$

Thus each norm-48 added point touches 552 oriented minimal heads, paired by $u \mapsto v - u$, and hence $3 \cdot 552 = 1656$ old caps. Every norm-64 vector instead has 46 heads with $v \cdot u = 32$. Their product with its normalized equatorial point is $32/\sqrt{64 \cdot 48} = 1/\sqrt{3} > 1/2$, giving exactly 138 cap conflicts for either endpoint. The contact checker verifies the first count for all 144 lines; the design argument proves the norm-64 obstruction for the whole shell.

6 Seven dimensions from one enlarged line class

The catalogue lattice P_{48p} is even unimodular of dimension 48 and minimum squared norm six. Its theta series determines a minimal shell of 52,416,000 vectors [30, 31].

6.1 Disjoint blocks from overlapping images

The new finite object is a class of 7,077 minimal lines. Integer representatives c in the catalogue Gram basis satisfy $cGc^t = 6$ and

$$|cGc'^t| \leq 2 \quad (c \neq \pm c').$$

Exact lattice isometries give further classes. Select a family of images and assign every line to its first occurrence. This produces disjoint line subsets with union size S . Taking both signs of every line produces compatible head blocks.

Use the root codes $A_1, A_2, D_3, D_4, D_5, E_6, E_7$ as tails in dimensions one through seven. Their sizes are 2, 6, 12, 24, 40, 72, 126. Assign each head block to a different antipodal tail line. A used mother line loses its two equatorial points and contributes four cap points at head squared norm $3/4$ and tail squared norm $1/4$. The count is

$$K(48 + k) \geq 52416000 - 2S + 4S + \tau_k = 52416000 + 2S + \tau_k.$$

k	Selected images	Union size S	Tail points τ_k
1	1	7,077	2
2	3	21,231	6
3	6	42,459	12
4	12	84,874	24
5	20	141,328	40
6	36	253,851	72
7	63	442,507	126

6.2 Why the lift preserves all angular constraints

Let x be a selected mother vector divided by $\sqrt{6}$, and let p_i represent its assigned unit tail line. The four points replacing $\{\pm x\}$ are

$$\left(\varepsilon \frac{\sqrt{3}}{2} x, \delta \frac{1}{2} p_i \right), \quad \varepsilon, \delta \in \{\pm 1\}.$$

They have unit norm. Within a block, distinct head lines have absolute product at most $1/3$; in the entire minimal shell the corresponding bound is $1/2$. These two facts give all cap inequalities:

$$\begin{aligned} \text{different lines in one block:} & \quad (3/4)(1/3) + 1/4 = 1/2, \\ \text{same head, opposite tails:} & \quad 3/4 - 1/4 = 1/2, \\ \text{opposite heads:} & \quad -3/4 + 1/4 = -1/2, \\ \text{different blocks:} & \quad (3/4)(1/2) + (1/4)(1/2) = 1/2. \end{aligned}$$

The last line uses distinct tail lines. Reusing a tail line for arbitrary different blocks would permit $5/8$. A cap meets a retained equatorial point with product at most $\sqrt{3}/4$, and a pure tail with product at most $1/2$. The retained equator and pure tails already satisfy the kissing condition. Thus the construction and its count are established by the class inequalities and the lattice minimum.

The latitude $1/2$ and disjoint assignment are necessary in this paired model, not only sufficient. Retained pure endpoints bound both heights $h_{\pm} = \sin \theta_{\pm}$ by $1/2$. The paired-head product $\cos(\theta_+ + \theta_-)$ can be at most $1/2$ only when both $\theta_{\pm} = \pi/6$. A head assigned to two tail pairs would then give a product at least $3/4$. Thus, for a fixed pool of compatible classes, maximizing the lifted count is exactly maximizing their union.

This also gives a computable selection guarantee. If a line x occurs in a_x of m candidate classes, a uniform t -subset ($0 \leq t \leq m$) misses it with probability $\binom{m-a_x}{t} / \binom{m}{t}$. Some selection

therefore covers at least

$$\left\lceil \sum_x \left(1 - \frac{\binom{m-a_x}{t}}{\binom{m}{t}} \right) \right\rceil$$

lines. The method of conditional expectations [32] makes the choice deterministic: maximizing the conditional expected final union at each step never decreases this expectation and attains the guarantee. `constructions/p48/coverage.py` implements this with exact integer scores grouped by incidence pattern; the saved bounds above are determined by their explicit image unions.

6.3 Separating the two sources of gain

The gain over the public 7,069-line class [15] has two sources: eight additional mother lines and improved choices of isometric images. Apply the selected images to the original class and call the union size S_0 ; let S_{ref} be the comparison construction’s union size. Then

$$N - N_{\text{ref}} = 2(S - S_0) + 2(S_0 - S_{\text{ref}}).$$

The first term holds the images fixed and measures class growth. The second holds the original class fixed and measures image selection.

Dimension	Class growth	Image selection	Total increase
49	16	0	16
50	48	2	50
51	96	18	114
52	192	64	256
53	320	154	474
54	572	296	868
55	996	414	1410

The three images in dimension 50 are disjoint. In dimension 55, overlap reduces the $8 \cdot 63$ possible new lines to 498 distinct lines; image selection on the original class supplies another 207. This is why enlarging one class and maximizing the union are complementary parts of the construction.

7 Extremal lattice sections and design moments

Dimension 43 counts minimal vectors perpendicular to a five-dimensional section; dimension 45 projects three fibres of a three-dimensional section. Complete projection domains and design moments determine the required statistics without enumerating the ambient shell.

7.1 Mother shells and moment equations

An extremal even unimodular lattice of rank 48 has minimum squared norm six and theta series

$$E_4^6 - 1440E_4^3\Delta + 125280\Delta^2 = 1 + 52416000q^3 + O(q^4).$$

Here $E_4 = 1 + 240q + 2160q^2 + \cdots$ and $\Delta = q - 24q^2 + 252q^3 + \cdots$; the constant term and vanishing first two coefficients determine this weight-24 form [2]. Its minimal shell X is an 11-design [29, Theorem 4.1].

For section vectors t_i with Gram H , let $y_i = x \cdot t_i$ and $f(y) = |\{x \in X : y(x) = y\}|$. Integrality, $y^\top H^{-1}y \leq 6$ and products with short section vectors restrict the signature domain. For $2m \leq 10$, the moments are

$$\sum_{x \in X} \prod_{\ell=1}^{2m} (x \cdot t_{i_\ell}) = \frac{|X|6^m}{48 \cdot 50 \cdots (48 + 2m - 2)} \sum_{\mathcal{P}} \prod_{(a,b) \in \mathcal{P}} H_{i_a i_b}, \quad (7.1)$$

where $\mathcal{P}$ pairs labelled positions; degree zero gives $|X|$. These include even total degrees with odd individual exponents. For a moment system $Av = b$, a rational identity $A^\top w = c$ proves $c^\top v = w^\top b$. Averaging must preserve the domain, moments and target, but need not extend to an ambient lattice automorphism.

7.2 Dimension 43: the fixed child and eight coordinate cuts

In P_{48n} [33], supplied integer rows T embed $\sqrt{3}E_8$: the ambient Gram G satisfies $(TGT^\top)_{ij} = 3\alpha_i \cdot \alpha_j$, where

$$\alpha_1 = \frac{1}{2}(1, -1, -1, -1, -1, -1, -1, 1), \quad \alpha_2 = e_1 + e_2, \quad \alpha_j = e_{j-1} - e_{j-2} \quad (3 \leq j \leq 8).$$

The first five rows specify the $\sqrt{3}D_5$ child U . This embedded-parent approach belongs to the cross-section framework of Dorofeev–Sun–Wang [27].

Proposition 7.1 (E_8 -extension count). *Let L be an extremal even unimodular lattice of rank 48 with an isometric embedding $\iota : \sqrt{3}E_8 \hookrightarrow L$, and let X_L be its squared-norm-six shell. The span U_L of the first five embedded simple roots, and every $W(E_8)$ -conjugate of this child, satisfies*

$$|X_L \cap U_L^\perp| = 2553792.$$

No ambient Weyl action or primitivity assumption is required. The supplied embedding gives $K(43) \geq 2553792$.

The parent projection is $\lambda/\sqrt{3}$ with $\lambda \in E_8$, by integrality and self-duality. Apart from the 240 singleton exceptions $\lambda = 3r$, the minimum gives $\|\lambda\|^2 \leq 18$ and $|\lambda \cdot r| \leq 3$ for every root r . Exact enumeration gives 1, 240, 2160, 6720, 17280 signatures at squared norms 0, 2, 4, 6, 8: 26641 signatures including exceptions. The radial moments force Weyl-averaged multiplicities

$$(a_0, a_2, a_4, a_6, a_8) = (1092000, 116235, 9180, 495, 63/4).$$

The child's perpendicular space meets these orbits in 1, 12, 6, 24, 0 signatures, with twelve exceptions, giving the average count $1092000 + 12(116235) + 6(9180) + 24(495) + 12 = 2553792$.

For the fixed child, the group $\langle W(D_5), W(A_3), -I \rangle$ preserves the target and has 43 domain orbits; A_3 consists of the roots perpendicular to the child. The certificate gives $A^\top w = c$ and $w^\top b = 2553792$ for regenerated moment and fixed-exception rows. This proves the proposition without assuming ambient symmetry. The complete domain is essential: the orthogonality condition belongs in c , not in prior column deletion. Normalizing by $\sqrt{6}$ gives a kissing configuration, since distinct minimal vectors have product at most three. The comparison has 2545056 points from an isometric D_5 : the gain is in its embedding.

Extend ι linearly and put $U_k = \iota(\sqrt{3} \text{span}\{e_1, \dots, e_k\})$, $1 \leq k \leq 8$. With $m = 8 - k$, the eight fixed counts are

$$|X_L \cap U_k^\perp| = 1092000 + 18360m + 464944 \binom{m}{2} + 11880 \binom{m}{3} \\ + 146880 \binom{m}{4} + 2520 \binom{m}{5} + 31680 \binom{m}{6}.$$

These concern the specified cuts and their Weyl conjugates, not arbitrary sections. U_1 is an axis, not a D_1 root system; the other cuts are coordinate D_k spans.

For completeness, in doubled coordinates $u = 2\lambda$, group the nonexceptional domain by permutations within the first k and last $8 - k$ positions and even sign changes. The polynomials of degree at most ten $m_a(u_1^2, \dots, u_k^2) m_b(u_{k+1}^2, \dots, u_8^2)$, $|a| + |b| \leq 5$, use monomial symmetric functions m_a and moments of Gram $12I_8$. The eight orbit matrices have full column ranks 25, 34, 39, 41, 39, 34, 25, 13, certified by nonzero integer minors. Thus actual orbit masses equal the Weyl-averaged masses $|O|a_r$. Counting target signatures gives the formula, whereas radial moments alone would give only an average over cuts. The Weyl map $\frac{1}{2}P_{5,8} \text{diag}(J_4 - 2I_4, J_4 - 2I_4)$ sends the specified D_5 span to the first five coordinate axes. Here $P_{5,8}$ exchanges coordinates five and eight.

k	1	2	3	4
$ X_L \cap U_k^\perp $	16815624	10663920	6688960	4149504
k	5	6	7	8
$ X_L \cap U_k^\perp $	2553792	1593664	1110360	1092000

7.3 Dimension 45: an exact fibre identity

Appendix A proves minimum six for the specified quadratic-residue even neighbour Λ . More generally, the following argument applies to any extremal even unimodular rank-48 lattice with a section of Gram $H = 8I_3 - 2J_3$, $H^{-1} = (I_3 + J_3)/8$, without primitivity.

The section vectors $t_1, t_2, t_3, -t_1 - t_2 - t_3$ and their negatives give eight singleton signatures $E = \{\pm H e_1, \pm H e_2, \pm H e_3, \pm H(1, 1, 1)^\top\}$. Every other signature lies in

$$D_0 = \{y \in \mathbb{Z}^3 : |y_i| \leq 3, |y_1 + y_2 + y_3| \leq 3\}. \quad (7.2)$$

Its 231 signatures have projection norm at most $9/2$. Subtracting E and pairing antipodes gives 161 moment rows and 116 variables, with $A_{a,[y]} = (2 - \mathbf{1}_{y=0})y^a$ and $b_a = M_a - \sum_{y \in E} y^a$.

Proposition 7.2 (Exact fibre identity). *Under these hypotheses,*

$$f(e_1) + f(e_2) + f(e_3) = 7377408 + 9f(-2, -2, 3).$$

Proof. Give c coefficients 1 on $[e_i]$, -9 on $[(-2, -2, 3)]$ and zero elsewhere. The rational multiplier in `dual.json` (Appendix B) satisfies $c - A^\top w \geq 0$, $w^\top b = 7377408$. Since $t_i + t_j$ has squared norm eight, the minimum also gives $|y_i + y_j| \leq 4$. With $y_4 = -\sum_{i=1}^3 y_i$, this removes the 30 permutations of $(3, 3, -3, -3)$ and $(3, 2, -3, -2)$. All seven positive slack entries lie on these excluded orbits. Thus $A_D^\top w = c_D$ on the remaining 101 antipodal orbits, sharpening the original lower certificate to equality. $\square$

Projecting the three fibres leaves squared norm $23/4$ and products at most $11/4$ within one fibre and $23/8$ between fibres. This excludes collisions and gives a kissing configuration after normalization. Thus m distinct witnesses in the last fibre give $K(45) \geq 7377408 + 9m$; its complete size gives an exact cardinality.

7.4 The anchor graph and the certified count

For a norm-eight anchor p , put $A_p = \{x \in X : x \cdot p = 4\}$. The minimum gives $|x \cdot p| \leq 4$. The filter $P(t) = t^2(t^2 - 1)(t^2 - 4)(t^2 - 9)$ has $P(4) = 20160$ and shell sum 90961920. Antipodality therefore gives $|A_p| = 2256$. Pair x with $p - x$. Distinct pairs have endpoint products in $\{1, 2, 3\}$; exchanging an endpoint replaces the product by four minus itself. Products one or three thus define a graph on 1128 pairs.

Filtering the mixed design moments through degree ten gives $\sum_{x \in A_p} (z \cdot x)^2 = 192\|z\|^2$ for $z \perp p$. For $z = r - p/2$, its own pair contributes 32 and each adjacent pair contributes two. Hence $768 = 32 + 2 \deg(\bar{r})$: every such graph is 368-regular.

For nonadjacent vertices represented by r, s , the section $(-s, s - p, p - r)$ has Gram H . Each vertex of $N(\bar{r}) \setminus N(\bar{s})$ has a unique endpoint a with $r \cdot a = 3$, and $w = p - a$ has signature $(-2, -2, 3)$. Conversely, that signature forces $p \cdot w = 4$, $r \cdot w = 1$ and $s \cdot w = 2$, recovering exactly that endpoint. The complete fibre therefore has size $368 - c$, where $c = |N(\bar{r}) \cap N(\bar{s})|$, and the projected count is $7380720 - 9c$. Every ordered section of Gram H has this representation: $p = -t_1 - t_2$, $s = -t_1$, $r = -t_1 - t_2 - t_3$.

Choose one endpoint r_i per pair and set $u_i = r_i - p/2$. The signed matrix S has zero diagonal and entries $S_{ij} = u_i \cdot u_j \in \{0, \pm 1\}$ for $i \neq j$. The filtered second moment gives

$$\sum_i u_i \otimes u_i = 96I_{p^\perp}, \quad S^2 = 88S + 368I, \quad \text{Spec}(S) = \{92^{(47)}, (-4)^{(1081)}\}.$$

Indeed, the Gram $4I + S$ has rank 47 and squares to 96 times itself. Signed common-neighbour counts satisfy $c_{ij}^\pm = (c_{ij} \pm 88S_{ij})/2$; thus codegrees are even and projected counts are divisible by 18. The unsigned adjacency matrix satisfies $A + 16I \succeq 0$ as the Gram of $u_i \otimes u_i$; this does not imply a two-valued unsigned spectrum. Four signed u_i cannot have all mutual products -1 : their sum would be a lattice vector of squared norm four.

The selected anchor has numerator shape $(3^6, 1^{42})$ in coordinates scaled by $1/\sqrt{12}$. Its 2256 certified endpoints are complete. All 428076 nonadjacent pairs have codegrees 70 to 148, with a unique pair attaining 70. Its fibre therefore has exactly 298 points, improving on the earlier 292-witness Pless-lattice construction.

Corollary 7.3 (The 45-dimensional construction). *The specified construction has exactly $7377408 + 9(298) = 7380090$ points, hence $K(45) \geq 7380090$. This is the largest projected count for this fixed anchor and Gram H , not an optimum over other anchors, lattices or section Grams.*

8 Construction principles

The nineteen bounds arise from several distinct ways of enlarging a constrained configuration. Tight contacts do not always prevent motion: the 552-point layer in dimension 25 moves with its internal Gram matrix unchanged. A saturated selection need not exhaust its candidate geometry: recolouring changes the second-layer selection in dimension 27, while shared blockers make a

family of support insertions profitable. A local capacity bound can be attained by importing a different combinatorial object, as the weight-four designs and their Hadamard images show.

Two further mechanisms use information supplied by the ambient lattice. Its minimum makes cross-shell compatibility automatic in dimension 38. Its design moments fix the eight coordinate-section counts, including the dimension-43 count, and express the dimension-45 count through an anchor graph. In the latter case, a complete 298-point fibre determines a configuration of exactly 7380090 points. For dimensions 49–55, a common class supplies seven lifts, and exact union counts distinguish the gain from class growth from the gain from image selection.

Qiushi Engine developed these constructions through sustained autonomous mathematical research, connecting geometric freedom, discrete selection and exact counting. The resulting proofs and finite objects make those connections available for further mathematics: they can be reconstructed, varied and used as starting points for new constructions.

A The minimum of the ambient lattice used in dimension 45

Let C be the extended ternary quadratic-residue $[48, 24, 15]$ code [31], and put $M = \{z \in \mathbb{Z}^{48} : z \bmod 3 \in C\}$ and $L_0 = \{z/\sqrt{3} : z \in M, z^2 \equiv 0 \pmod{6}\}$. The supplied basis determines a characteristic numerator $\chi \in M$ with odd coordinates, $\chi^2 = 264$, and $\chi \cdot z \equiv z^2 \pmod{6}$. Define

$$\Lambda = L_0 \cup (\chi/(2\sqrt{3}) + L_0).$$

The characteristic congruence and $\chi^2/12 = 22$ give integrality and evenness; the two index-two constructions give covolume one. The program reconstructs an integer numerator basis B and verifies $BB^T = 12G$ for the exact Gram matrix used by the 298-vector certificate.

In L_0 , squared norm two or four would require $z^2 \leq 12$. The code minimum weight 15 forces $z \bmod 3 = 0$, whose positive even lattice norms are at least six. In the other coset, $y = \chi + 2z$ has 48 odd coordinates, so $y^2/12 \geq 4$. Equality requires $y_i = \pm 1$.

Let $s_i \in \{\pm 1\}$ represent $\chi_i \bmod 3$. The finite checks give $s \in C$, $\chi \cdot s = 66$, and $\chi_i s_i \equiv 1 \pmod{6}$. Normalizing coordinates by s gives a code containing the all-one word. Its two-symbol enumerator is $1 + 94t^{24} + t^{48}$ [34, equation (20)], so binary-valued codewords have weights 0, 24 or 48. Writing $y = s(1 - 2b)$ gives such a binary word of weight w , but

$$\chi \cdot z = \frac{\chi \cdot s - \chi^2}{2} - \sum_i \chi_i s_i b_i \equiv -99 - w \equiv 3 \pmod{6},$$

contrary to $z/\sqrt{3} \in L_0$. Evenness excludes all other positive norms below six, while $\sqrt{3}(e_1 + e_2)$ has squared norm six. The theta-series and design-moment arguments therefore apply to this specified lattice.

B Construction data and exact verification

This PDF contains the attachment `certificates.zip`. Save it from the PDF viewer's attachment panel, or extract it with `pdfdetach -saveall main.pdf`. It contains the complete finite inputs and the programs used to check them. The English and Chinese editions embed identical archives.

Each construction combines a geometric argument covering all pair types with finite input satisfying its hypotheses. For the code constructions, the input consists of supports, generators,

coset representatives and signed vectors. For the lattice constructions, it consists of integer directions, Gram matrices, embedded sections and rational moment certificates. The catalogue `constructions/catalog/results.json` specifies the defining files, exact count and verification tasks for each dimension. The following paths are relative to `constructions/`.

For dimension 25, `d25/` contains the exact baseline heads, the two integer repair vectors, the complete moving-layer proof and the integer, rational and Arb verifiers. The interval checks require `python-flint`. The motion-obstruction checker proves that at least two cap changes are required for the specified insertion and chord family. For dimension 27, `d27/data/` gives both head layers and their labels; the independent checker regenerates the shell and verifies all deletion and product conditions. The candidate-graph reconstruction is included. For dimensions 49–55, `p48/data/` supplies the catalogue basis, generators, enlarged class and seven image families. The exact verifier rebuilds every image union and checks tail ranks and the lifting inequalities. The search program also verifies the eight-line augmentation and decomposes the union gain relative to the original class.

For dimensions 32–37 and 39, `codes/data/` contains the complete support lists, binary generators, coset representatives and exchange records. The local replacements for dimensions 35 and 36 are in `codes/hadamard/data/`; their generator file and the reconstruction program reproduce the stored signed vectors exactly. Dimension 37 has its own 512-point replacement and coordinate map in `codes/data/`. The solver `codes/hadamard/boundary.py` returns an optimal source family and matching certificate for each complete-type transverse $2+2$ boundary in twelve coordinates.

For dimension 38, `d38/base-data/` specifies the Golay generator, Leech line classes, tail triangles and relative rotation defining the 591,612-point base. The 144 added representatives are in `d38/added-lines.txt`. The checker reconstructs the shell, verifies the base, tests the Leech congruences of every new vector, and checks all new mutual and boundary products in exact arithmetic. The contact checker also derives the 552 endpoint contacts and the 46 positive extreme products excluding norm-64 additions to this base.

For dimension 43, `sections/d43/data/parent-section.txt` gives the ambient Gram matrix and the eight rows embedding the numbered E_8 roots. `sections/d43/data/certificate.json` contains the orbit data, rational moment matrix and multiplier determining the D_5 count. One program reconstructs this system from the geometry; another checks the rational matrix identity independently. The coordinate-family checker reconstructs eight fixed-cut moment systems, certifies full column rank and checks their Weyl correspondence with the specified D_5 section.

For dimension 45, `sections/qr/` reconstructs the extremal neighbour. Within `sections/d45/data/`, `record.json` contains the section and 298 witnesses, `dual.json` the exact multiplier of Proposition 7.2, and `parent-vectors.json` the 2,256-point parent set. The checkers regenerate the moment domain and equations, verify lattice membership and inner products, and recover the same witnesses from the parent graph. The parent checker also verifies the signed Gram identity and determines the minimum common-neighbour count over all 428076 nonadjacent pairs.

After extracting the archive, its `README.md` gives the dependencies and commands. Use a Python environment providing SageMath, NumPy, SymPy and `python-flint`, together with a C++ compiler. In that environment, `python3 tools/reproduce.py --suite all --output ../verification` runs every listed check in a separate directory, without network access. The output records each check and its inputs. The archive’s manifest identifies every distributed file; the full repository also contains the report sources and the recorded verification results.

C Construction methods and research development

The mathematical arguments above arise from choices of finite objects, coordinate models and counting methods. This account connects those choices to the candidate families, searches, intermediate constructions and exact calculations that produced the results.

C.1 Moving the retained shell

The 25-dimensional investigation changes the geometry of retained points, rather than restricting every improvement to the original head pool. A complete block of deleted Leech vectors has retained partners on the opposite latitude. Moving all partners together preserves their internal products. Their horizontal coordinates stay on segments between pairs of original vectors, so convexity settles compatibility with the unchanged equator. Two exceptional caps are then replaced by normalized integer vectors. The analytic motion and finite repairs have separate, short verification arguments.

C.2 From assigning blocks to reusing directions

The first Leech lifts retained a large fixed mother shell and used published 496-point compatible blocks. The useful variable was the assignment of these blocks to tail labels. In dimension 25, the latitude at height one half permitted two poles. The count became 197,058 without enlarging the block.

In dimension 27, twelve labels could be partitioned into four zero-sum triangles. Replacing block multiplicities $(3, 3, 2, 2, 2)$ by $(3, 3, 3, 3)$ preserved one more 496-point block at the equator, giving 200,540. The capacity argument then explained the count: a direction could occur at most three times, and equality required full labels with multiplicity three.

The later 201,567-point result uses a different public input. Lindow's contribution had jointly optimized the first and second layers to 201,566. The new witness keeps that first layer and increases the second layer from 310 to 311 owners. The first layer's 2,090 deleting heads and 168 nondeleting heads remain unchanged.

New heads may contribute two or three points after their own deletion costs, and a second-layer owner contributes one. Optimizing only a layer's gross population can select a worse complete configuration.

C.3 From block libraries to image unions above dimension 48

The initial family used 63 disjoint blocks of a Pless neighbour. Its prefix sizes, combined with root tail codes, produced seven configurations in dimensions 49 through 55. The paper retains the neighbour's construction and minimum-norm proof.

The larger family uses the published catalogue basis of P_{48p} . A public class of 7,069 minimal lines was extended to 7,077. Its automorphism images were then selected jointly, using the size of their union as the objective. Assigning repeated lines to their first occurrence produced disjoint subsets without losing the within-class product bound.

The seven union sizes are

$$7077, 21231, 42459, 84874, 141328, 253851, 442507.$$

Each contributes two points per line, in addition to the mother shell and the root tail code. The dimension-50 witness uses three disjoint images. Larger image families retain some overlaps, whose exact effect is recorded in the union sizes.

Different assigned classes use distinct antipodal tail lines. This gives the cross-class product bound $(3/4)(1/2) + (1/4)(1/2) = 1/2$. An injective tail assignment therefore accompanies the union of the selected head classes.

References

- [1] Oleg R. Musin. The kissing number in four dimensions. *Annals of Mathematics*, 168(1): 1–32, 2008. doi: [10.4007/annals.2008.168.1](https://doi.org/10.4007/annals.2008.168.1).
- [2] John H. Conway and Neil J. A. Sloane. *Sphere Packings, Lattices and Groups*, volume 290 of *Grundlehren der mathematischen Wissenschaften*. Springer, New York, third edition, 1999. doi: [10.1007/978-1-4757-6568-7](https://doi.org/10.1007/978-1-4757-6568-7).
- [3] F. Jessie MacWilliams and Neil J. A. Sloane. *The Theory of Error-Correcting Codes*, volume 16 of *North-Holland Mathematical Library*. North-Holland, Amsterdam, 1977.
- [4] John H. Conway and Neil J. A. Sloane. Laminated lattices. *Annals of Mathematics*, 116(3):593–620, 1982. doi: [10.2307/2007025](https://doi.org/10.2307/2007025).
- [5] John H. Conway and Neil J. A. Sloane. The antipode construction for sphere packings. *Inventiones Mathematicae*, 123(2):309–313, 1996. doi: [10.1007/s002220050028](https://doi.org/10.1007/s002220050028).
- [6] Yves Edel, Eric M. Rains, and Neil J. A. Sloane. On kissing numbers in dimensions 32 to 128. *Electronic Journal of Combinatorics*, 5(1):R22, 1998. doi: [10.37236/1360](https://doi.org/10.37236/1360).
- [7] Henry Cohn, Yang Jiao, Abhinav Kumar, and Salvatore Torquato. Rigidity of spherical codes. *Geometry & Topology*, 15(4):2235–2273, 2011. doi: [10.2140/gt.2011.15.2235](https://doi.org/10.2140/gt.2011.15.2235).
- [8] Kenz Kallal, Tomoka Kan, and Eric Wang. Improved lower bounds for kissing numbers in dimensions 25 through 31. *SIAM Journal on Discrete Mathematics*, 31(3):1895–1908, 2017. doi: [10.1137/16M1095810](https://doi.org/10.1137/16M1095810).
- [9] Alexander Novikov, Ngân Vũ, Marvin Eisenberger, Emilien Dupont, Po-Sen Huang, Adam Zsolt Wagner, Sergey Shirobokov, Borislav Kozlovskii, Francisco J. R. Ruiz, Abbas Mehrabian, M. Pawan Kumar, Abigail See, Swarat Chaudhuri, George Holland, Alex Davies, Sebastian Nowozin, Pushmeet Kohli, and Matej Balog. AlphaEvolve: A coding agent for scientific and algorithmic discovery. [arXiv:2506.13131v1](https://arxiv.org/abs/2506.13131v1), 2025. Version 1, 16 June 2025.
- [10] Chengdong Ma, Théo Tao Zhaowei, Pengyu Li, Minghao Liu, Haojun Chen, Zihao Mao, Yuan Cheng, Yuan Qi, and Yaodong Yang. Finding kissing numbers with game-theoretic reinforcement learning. [arXiv:2511.13391v5](https://arxiv.org/abs/2511.13391v5), 2026. Version 5, 22 September 2026.
- [11] Federico Bianchi, Yongchan Kwon, Aneesh Pappu, and James Zou. Harnessing the collective intelligence of AI agents in the wild for new discoveries. [arXiv:2606.10402v2](https://arxiv.org/abs/2606.10402v2), 2026. Version 2, 26 August 2026.
- [12] Stephen Chung, Wenyu Du, and William J. Wesley. Autonomous mathematical discovery in an open-world multi-agent environment. [arXiv:2608.23691v2](https://arxiv.org/abs/2608.23691v2), 2026. Version 2, 14 September 2026.
- [13] Shuxing Yang, Fujia Chen, Rui Zhao, Junyao Wu, Yize Wang, Haiyao Luo, Ning Han, Qiaolu Chen, Yuze Hu, Wenhao Li, Mingzhu Li, Hongsheng Chen, and Yihao Yang. End-to-end autonomous scientific discovery on a real optical platform. [arXiv:2604.27092v1](https://arxiv.org/abs/2604.27092v1), 2026. Version 1, 29 April 2026.

- [14] Fredrik Johansson. Arb: efficient arbitrary-precision midpoint-radius interval arithmetic. *IEEE Transactions on Computers*, 66(8):1281–1292, 2017. doi: [10.1109/TC.2017.2690633](https://doi.org/10.1109/TC.2017.2690633).
- [15] Alexey Kravatskiy. Kissing numbers: new lower bounds in dimensions 18 through 96, with verification packages. GitHub repository, revision [cf14c5e](#), 2026. Accessed 27 September 2026.
- [16] btindow. Dimension 27: the two-layer configuration jointly re-optimised. GitHub pull request [no. 1](#), 2026. Contribution of a 201566-point configuration; revision [277c297058d58a756e57ad8e3c6c710f4d29f2b6](#).
- [17] Andries E. Brouwer. Bounds for constant weight codes. <https://aeb.win.tue.nl/codes/Andw.html>. Online code table; accessed 27 September 2026.
- [18] Xiaoming Sun and Chengu Wang. Sphere packings and kissing numbers in dimensions 39, 43, and 45 from the antipode construction. [arXiv:2607.20359v3](#), 2026. Version 3, 18 August 2026; version containing the comparison counts in dimensions 43 and 45.
- [19] wcgbg. sphere-packing. GitHub repository, revision [a0bc7f5](#), 2026. Construction data and verification programs; accessed 24 September 2026.
- [20] William Echols. New lower bounds for constant-weight codes via seeded bit-swap tabu search. [arXiv:2608.13906v1](#), 2026. Version 1, 14 August 2026.
- [21] Chengdong Ma, Théo Tao Zhaowei, Pengyu Li, Minghao Liu, Haojun Chen, Zihao Mao, Bo Li, Yuan Cheng, Yuan Qi, and Yaodong Yang. Finding kissing numbers with game-theoretic reinforcement learning. [arXiv:2511.13391v4](#), 2026. Version 4, 2 June 2026; Table 1 and Appendix F.7.
- [22] PackingStar contributors. PackingStar: coordinate data. GitHub coordinate dataset, revision [50ea645](#), 2026. Leech blocks and tail coordinates; accessed 14 September 2026.
- [23] Ying Cheng and Neil J. A. Sloane. Codes from symmetry groups, and a $[32, 17, 8]$ code. *SIAM Journal on Discrete Mathematics*, 2(1):28–37, 1989. doi: [10.1137/0402003](https://doi.org/10.1137/0402003).
- [24] Andries E. Brouwer, James B. Shearer, Neil J. A. Sloane, and Warren D. Smith. A new table of constant weight codes. *IEEE Transactions on Information Theory*, 36(6):1334–1380, 1990. doi: [10.1109/18.59932](https://doi.org/10.1109/18.59932).
- [25] Rustem Takhanov and Stanislav Yun. Classification of independent sets in signed Johnson graphs and applications to kissing arrangements. [arXiv:2606.03299v1](#), 2026. Version 1, 2 June 2026.
- [26] J. G. Kalbfleisch and R. G. Stanton. Maximal and minimal coverings of $(k - 1)$ -tuples by k -tuples. *Pacific Journal of Mathematics*, 26(1):131–140, 1968. doi: [10.2140/pjm.1968.26.131](https://doi.org/10.2140/pjm.1968.26.131).
- [27] Ivan Dorofeev, Xiaoming Sun, and Chengu Wang. Optimal extensions of cross-sections: Sphere packings in dimensions 38 to 43. [arXiv:2607.20359v4](#), 2026. Version 4, 29 August 2026.
- [28] Philippe Delsarte, Jean-Marie Goethals, and Johan J. Seidel. Spherical codes and designs. *Geometriae Dedicata*, 6(3):363–388, 1977. doi: [10.1007/BF03187604](https://doi.org/10.1007/BF03187604).

- [29] Gabriele Nebe. Boris Venkov’s theory of lattices and spherical designs. In *Diophantine Methods, Lattices, and Arithmetic Theory of Quadratic Forms*, volume 587 of *Contemporary Mathematics*, pages 1–19. American Mathematical Society, Providence, RI, 2013. doi: 10.1090/conm/587/11672. Preprint: arXiv:1201.1834v1.
- [30] Gabriele Nebe and Neil J. A. Sloane. Catalogue of Lattices: the lattice P_{48p} . <https://www.math.rwth-aachen.de/~Gabriele.Nebe/LATTICES/P48p.html>, undated. Last modified 18 July 2014; accessed 27 September 2026.
- [31] Gabriele Nebe. On extremal self-dual ternary codes of length 48. *International Journal of Combinatorics*, 2012:1–9, 2012. doi: 10.1155/2012/154281. Article ID 154281.
- [32] Prabhakar Raghavan. Probabilistic construction of deterministic algorithms: Approximating packing integer programs. *Journal of Computer and System Sciences*, 37(2):130–143, 1988. doi: 10.1016/0022-0000(88)90003-7.
- [33] Gabriele Nebe and Neil J. A. Sloane. Catalogue of Lattices: the lattice P_{48n} . *Catalogue entry CQ48a*, undated. Last modified 18 July 2014; accessed 27 September 2026.
- [34] Akihiro Munemasa and Hiroki Tamura. The codes and the lattices of Hadamard matrices. *European Journal of Combinatorics*, 33(4):519–533, 2012. doi: 10.1016/j.ejc.2011.11.007.