

# Efficiently Calculating Jones Polynomials at Lattice Roots of Unity with the ZX-Calculus (Draft)

Alex Townsend-Teague

Konstantinos Meichandzitis

## Abstract

We leverage the ZX-calculus, a diagrammatic language for quantum mechanics, to provide a graphical proof of an established result from pure topology, namely that evaluating the Jones polynomial of arbitrary links at the lattice roots of unity is in  $\mathbf{P}$ . For these cases, it is known that the problem reduces to the evaluation of the partition function  $Z(q)$  of a planar  $q$ -state Potts model. This partition can be represented as a closed tensor network, which can in turn be represented as a ZX-diagram, since the ZX-calculus is sound and complete for tensor networks. The ZX-calculus is based on a set of graphical rewrite rules; we show that there exist polynomially long sequences of such rewrites that reduce this ZX-diagram to an equivalent trivial one, thereby calculating  $Z(q)$  for  $q \in \{1, 2, 3, 4\}$ . This corresponds exactly to evaluating the Jones polynomial at the lattice roots of unity.

## 1 Introduction

[Use Konstantinos' O.G. notes from the start of this project]

## 2 Simplifying Qubit ZX-Diagrams

### 2.1 The Qubit ZX-Calculus

Here we give a quick introduction to the qubit ZX-calculus. For a more detailed treatment, see [Coecke and Kissinger, 2017, Chapter 9.4]. The qubit ZX-calculus is a diagrammatic language for quantum mechanics consisting of *spiders* connected together by *wires*, living within a rectangular plane. Diagrams are read bottom-to-top. Wires that don't end at a spider are *inputs* (if their non-spider end starts at the bottom of the plane) and/or *outputs* (if their non-spider end lies at the top of the plane).

Only the topology matters; spiders can move freely around the plane and wires can be deformed and pass over each other, provided the connections between components remain unchanged. The single exception is the endpoints of input and output wires; these are fixed on the boundary of the plane. Spiders come in two species: green  $Z$ -spiders and red  $X$ -spiders, decorated by a *phase* angle  $\alpha$ . When this phase is zero, we will omit it. The diagrammatic formalism exists in the abstract, but we say that it has a *standard linear algebraic interpretation*.

**Definition 2.1.** Given the usual vectors  $\{|0\rangle, |1\rangle\}$  (the  $Z$ -basis) and  $\{|+\rangle, |-\rangle\}$  (the  $X$ -basis) in  $\mathbb{C}^2$ , the *standard interpretations* of  $Z$ - and  $X$ - spiders respectively are:

$$\left[ \begin{array}{c} \overbrace{\quad}^m \\ \vdots \\ \text{green spider } \alpha \\ \vdots \\ \underbrace{\quad}_n \end{array} \right] = |0\rangle^m \langle 0|^n + e^{i\alpha} |1\rangle^m \langle 1|^n, \quad \left[ \begin{array}{c} \overbrace{\quad}^m \\ \vdots \\ \text{red spider } \alpha \\ \vdots \\ \underbrace{\quad}_n \end{array} \right] = |+\rangle^m \langle +|^n + e^{i\alpha} |-\rangle^m \langle -|^n$$

Using this standard interpretation, and writing  $\simeq$  to mean ‘equal up to a scalar factor’, we can represent common quantum computing concepts as ZX-diagrams.

**Example 2.2.** The standard interpretations of the diagrams below are as follows:


$$\left[ \begin{array}{c} | \\ \text{green spider} \end{array} \right] \simeq |+\rangle, \quad \left[ \begin{array}{c} | \\ \text{red spider } \pi \end{array} \right] \simeq |1\rangle, \quad \left[ \begin{array}{c} | \\ \text{red spider } \alpha \end{array} \right] = X_\alpha, \quad \left[ \begin{array}{cc} | & | \\ \text{green spider} & \text{red spider} \end{array} \right] \simeq \text{CNOT} \quad (1)$$

**Remark 2.3.** In the ZX-calculus, equality of diagrams translates to equality only up to some scalar factor in the standard linear algebraic interpretation. This has often proved to be more convenient in quantum computing, but for calculating Jones polynomials of knots at lattice roots of unity - which are ultimately just complex numbers - this is manifestly insufficient. However, to simply prove that the calculation of such polynomials is efficient, this suffices - provided we also show that keeping track of the ‘missing’ scalars can be done efficiently. See the end of this section for a discussion of this last point.

We introduce a small yellow box to denote the Hadamard gate  $H$ . We will also often omit the yellow box entirely and instead draw a dashed blue line - we call these equivalent diagrams a *Hadamard edge*:

$$\begin{array}{c} | \\ \text{yellow box} \end{array} = \begin{array}{c} | \\ \text{dashed blue line} \end{array} = \begin{array}{c} \text{green spider } \frac{\pi}{2} \\ \text{red spider } \frac{\pi}{2} \\ \text{green spider } \frac{\pi}{2} \end{array} \quad (2)$$

The diagrams that can be formed from such spiders and wires are then subject to the rewrite rules in Figure 1. These rules constitute the definition of the ZX-calculus. Note



**Figure 1:** Rewrite rules for the qubit ZX-calculus. The fusion rule (f) applies to spiders connected by at least one wire.

that all rules hold with the roles of red and green (i.e.  $Z$  and  $X$ ) interchanged.

We are seeking a ZX-diagram whose standard interpretation is equal (up to a scalar) to the matrix  $T_{2\pm}$  in (ToDo: ref). However, via the isomorphism  $\mathbb{C}^4 \cong \mathbb{C}^2 \otimes \mathbb{C}^2$ , we can also find a *qubit* ZX-diagram whose standard interpretation is equal (up to a scalar) to the matrix  $T_{4\pm}$  in (ToDo: ref).

**Proposition 2.4.** Under the standard interpretation as linear maps, the following diagrams give (up to a scalar) the required matrices:

$$\left[ \begin{array}{c} \text{red spider with } \pm \frac{\pi}{2} \end{array} \right] \simeq \left[ \begin{array}{c} \text{white box with } \pm \end{array} \right]_{q=2}, \quad \left[ \begin{array}{c} \text{green spider with } \pi \text{ and red spider with } \pi \end{array} \right] \simeq \left[ \begin{array}{c} \text{white box with } \pm \end{array} \right]_{q=4}. \quad (3)$$

*Proof.* See Appendix [ToDo]. □

The *stabilizer fragment* of the calculus consists of all diagrams in which all phases are integer multiples of  $\frac{\pi}{2}$ . We note that both diagrams above fall into this stabilizer fragment. Now, in [Duncan et al., 2020, Theorem 5.4] the authors give an efficient algorithm for


simplifying any qubit stabilizer ZX-diagram to an equivalent diagram with fewer spiders. In particular, given a stabilizer diagram that is *closed* (has neither inputs nor outputs), the algorithm efficiently simplifies it until it contains at most one spider. If it exists, this spider will be legless with phase in  $\{0, \pi\}$ . We briefly outline the process here, referring the reader to Duncan et al. [2020] for a full discussion. First it is shown that every diagram is equivalent to a *graph-like* diagram.

**Definition 2.5.** [Duncan et al., 2020, Definition 3.1] A qubit ZX-diagram is *graph-like* when:


1. Every spider is a Z-spider.
2. Spiders are only connected by Hadamard edges.
3. There are no parallel Hadamard edges or self-loops.
4. Every input and output is connected to a spider.
5. Every spider is connected to at most one input or output.

Then the following two results are used to eliminate spiders. For more details on *local complementation* and *pivoting*, see [Duncan et al., 2020, Section 4].

**Theorem 2.6.** The following equation holds in the qubit ZX-calculus:



This says we can remove any spider  $i$  with phase  $\pm\frac{\pi}{2}$ , at the cost of performing a local complementation at  $i$ . Similarly, the following equation holds in the qubit ZX-calculus:



This says we can remove any pair  $i, j$  of spiders with phases in  $\{0, \pi\}$  connected by a Hadamard edge, at the cost of performing a local pivot along the edge  $uv$ .

*Proof.* These are Lemmas 5.2 and 5.3 in Duncan et al. [2020] □

After each application of (4) or (5), the following lemma is used to remove any parallel Hadamard edges, and ensure the diagram remains graph-like.

**Lemma 2.7.** The following result holds in the qubit ZX-calculus:

$$(6)$$

*Proof.* This is Equation (3) in Duncan et al. [2020].  $\square$

Since any ZX-diagram derived from a knot in the manner described in (ToDo: ref) will be closed, this algorithm suffices to prove that the calculation of the Jones polynomial of any knot at the lattice roots of unity  $\pm 1$  and  $\pm i$  is efficient. This is because reading off the scalar at the end is trivial; either we have the empty diagram, which has standard interpretation 1, or a single legless  $Z$ -spider with phase  $k\pi$ :

$$\llbracket \textcircled{k\pi} \rrbracket = \left\llbracket \begin{array}{c} \circ \\ | \\ \textcircled{k\pi} \end{array} \right\rrbracket = (\langle 0| + \langle 1|)(|0\rangle + e^{ik\pi} |1\rangle) = 1 + e^{ik\pi} \quad (7)$$

Furthermore, keeping track of the scalar factors introduced with each application of Theorem 2.6 or Lemma 2.7 can be done efficiently. [ToDo: proof, if not explicitly doing all this in a scalar-exact fashion. Reference Backens]

### 3 Simplifying Qutrit ZX-Diagrams

For the remaining case  $q = 3$ , we will show that we can generalise all the ideas of the previous section to the qutrit ZX-calculus, which we must first define.

#### 3.1 The Qutrit ZX-Calculus

[ToDo: make it more bitesize - theorems, definitions, etc?]

As before, this calculus is based around spiders and wires, but there are key differences, some subtler than others. To get a feel for the qutrit calculus, we outline these differences briefly here before giving a formal definition. First of all, the qubit  $Z$ -basis consisted of two vectors  $|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ , whereas the qutrit  $Z$ -basis consists of three vectors:

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad |1\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad |2\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (8)$$

Similarly, letting  $\omega = e^{\frac{4}{3}\pi i}$ , the qutrit  $X$ -basis consists of the three vectors:

$$|+\rangle = \frac{1}{\sqrt{3}} (|0\rangle + |1\rangle + |2\rangle) \quad (9)$$

$$|\omega\rangle = \frac{1}{\sqrt{3}} (|0\rangle + \omega |1\rangle + \bar{\omega} |2\rangle) \quad (10)$$

$$|\bar{\omega}\rangle = \frac{1}{\sqrt{3}} (|0\rangle + \bar{\omega} |1\rangle + \omega |2\rangle) \quad (11)$$

Spiders thus carry *two* phases  $\alpha$  and  $\beta$ , and have the following standard interpretation as linear maps:

$$\left[ \begin{array}{c} \overbrace{\quad}^m \\ \vdots \\ \text{green circle with } \alpha \text{ and } \beta \\ \vdots \\ \underbrace{\quad}_n \end{array} \right] = |0\rangle^m \langle 0|^n + e^{i\alpha} |1\rangle^m \langle 1|^n + e^{i\beta} |2\rangle^m \langle 2|^n \quad (12)$$

$$\left[ \begin{array}{c} \overbrace{\quad}^m \\ \vdots \\ \text{red circle with } \alpha \text{ and } \beta \\ \vdots \\ \underbrace{\quad}_n \end{array} \right] = |+\rangle^m \langle +|^n + e^{i\alpha} |\omega\rangle^m \langle \omega|^n + e^{i\beta} |\bar{\omega}\rangle^m \langle \bar{\omega}|^n \quad (13)$$

When  $\alpha = \beta = 0$  we will again omit the angles entirely, and just draw a small green or red dot.

**Warning 3.1.** Throughout, we slightly abuse notation by letting 1 and 2 denote the commonly used angles  $\frac{2\pi}{3}$  and  $\frac{4\pi}{3}$  respectively. This shouldn't ever cause confusion.

Hadamard boxes change too: they are now neither self-conjugate nor self-adjoint, though they remain self-transpose. Therefore in keeping with the diagrammatic paradigm we ought to stop representing them as boxes, and instead pick something that has neither vertical nor horizontal symmetry, but is invariant under rotation by  $\pi$ . A parallelogram fits the bill.

**Definition 3.2.** In the qutrit ZX-calculus, the *Hadamard edge* ( $H$ -edge) and its adjoint ( $H^\dagger$ -edge) are respectively the two diagrams below:

$$\text{yellow parallelogram} = \begin{array}{c} \text{green circle with } \frac{2}{2} \\ \text{red circle with } \frac{2}{2} \\ \text{green circle with } \frac{2}{2} \end{array} \quad (14)$$

Now comes the most important difference: in the qutrit ZX-calculus there is no ‘plain’ cap or cup. That is, we have the following two results:

$$\text{cup}_{\text{green}} \neq \text{cup}_{\text{red}} \quad , \quad \text{cap}_{\text{green}} \neq \text{cap}_{\text{red}} \quad . \quad (15)$$

This has several consequences. Firstly, the maxim that ‘only topology matters’ no longer applies. That is, it is now important to make clear the distinction between a spider’s inputs and outputs, unlike in the qubit case where we could freely interchange the two. Diagram components can still be isotoped around the plane but only so long as this input/output distinction is respected. This gives the qutrit calculus a slightly more rigid flavour than its qubit counterpart. That said, this rigidity is loosened by the following two results: firstly, this distinction is irrelevant for  $H$ - and  $H^\dagger$ -edges.

**Lemma 3.3.** In the qutrit ZX-calculus, the following results hold:

(16)

*Proof.* These are Lemmas 3.2 and 3.3 in Gong and Wang [2017].  $\square$

Secondly, the ‘snake equations’ continue to hold in some form; given a cap and cup of different colours, we can still straighten out the wire between them as before, only now it comes at the cost of adding two Hadamard boxes to the wire. For a cup and cap of the same colour, we can straighten out the wire as usual.

**Lemma 3.4.** In the qutrit ZX-calculus, the following results hold. Moreover, they hold with the roles of green and red interchanged.

$$\text{cup}_{\text{green}} \text{cap}_{\text{green}} = \text{straight wire} \quad , \quad \text{cup}_{\text{green}} \text{cap}_{\text{red}} = \text{straight wire with two Hadamard boxes} \quad . \quad (17)$$


*Proof.* ToDo  $\square$

We now give a full set of rules defining the qutrit ZX-calculus. Our presentation aims for clarity and accessibility; for a more rigorous description, see [Wang, 2018].

**Definition 3.5.** The *qutrit ZX-calculus* is a graphical calculus generated by the following diagrams, where  $\alpha, \beta \in [0, 2\pi]$ :

$$\begin{array}{c} \dots \\ \text{green spider } \alpha, \beta \\ \dots \end{array} \quad , \quad \begin{array}{c} \dots \\ \text{red spider } \alpha, \beta \\ \dots \end{array} \quad , \quad \text{X} \quad , \quad \text{Hadamard box} \quad (18)$$

and their adjoints  $(-)^{\dagger}$ . The rightmost two generators are self-adjoint, whereas spiders’ adjoints are found by vertical reflection and negation of angles:



**Figure 2:** Rewrite rules for the qutrit ZX-calculus.



$$\left( \begin{array}{c} \overbrace{\quad \quad \quad}^m \\ \vdots \\ \text{green circle with } \alpha \text{ over } \beta \\ \vdots \\ \underbrace{\quad \quad \quad}_n \end{array} \right)^\dagger = \begin{array}{c} \overbrace{\quad \quad \quad}^n \\ \vdots \\ \text{green circle with } -\alpha \text{ over } -\beta \\ \vdots \\ \underbrace{\quad \quad \quad}_m \end{array}, \quad \left( \begin{array}{c} \overbrace{\quad \quad \quad}^m \\ \vdots \\ \text{red circle with } \alpha \text{ over } \beta \\ \vdots \\ \underbrace{\quad \quad \quad}_n \end{array} \right)^\dagger = \begin{array}{c} \overbrace{\quad \quad \quad}^n \\ \vdots \\ \text{red circle with } -\alpha \text{ over } -\beta \\ \vdots \\ \underbrace{\quad \quad \quad}_m \end{array} \quad (19)$$

These generators can be composed in parallel ( $\otimes$ ) and sequentially ( $\circ$ ), and the resulting diagrams are governed by the rewrite rules in Figure 2, wherein addition is modulo  $2\pi$ . The fusion rule (f) applies to spiders of the same colour connected by at least one wire. Importantly, all the rules hold under taking adjoints, where for diagrams  $D$  and  $E$  we have:

$$(D \otimes E)^\dagger = D^\dagger \otimes E^\dagger \quad (20)$$

$$(D \circ E)^\dagger = D^\dagger \circ E^\dagger \quad (21)$$

Furthermore, all but the commutation equations (**cm**<sub>1,2</sub>) and (**cm**<sub>2,1</sub>) and the colour change equations (**cc**<sub>a</sub>) and (**cc**<sub>b</sub>) continue to hold when the roles of green and red (i.e.  $Z$  and  $X$ ) are interchanged. For these four exceptions, however, analogous equations can be derived from the existing ones; for example, the corresponding colour change equations will be relevant for us later.

**Proposition 3.6.** The following equations are derivable in the qutrit ZX-calculus:

$$\begin{array}{c} \vdots \\ \text{red circle with } \alpha \text{ over } \beta \\ \vdots \end{array} = \begin{array}{c} \text{yellow trapezoid} \\ \vdots \\ \text{green circle with } \alpha \text{ over } \beta \\ \vdots \\ \text{yellow trapezoid} \end{array}, \quad \begin{array}{c} \vdots \\ \text{red circle with } \alpha \text{ over } \beta \\ \vdots \end{array} = \begin{array}{c} \text{yellow trapezoid} \\ \vdots \\ \text{green circle with } \alpha \text{ over } \beta \\ \vdots \\ \text{yellow trapezoid} \end{array} \quad (22)$$

*Proof.* Add  $H$ - and  $H^\dagger$ -boxes to both sides of the original colour change equations in such a way that we can then cancel Hadamards on the legs of the red spiders via (**H**).  $\square$

Having now defined the qutrit ZX-calculus we turn our attention back to our tensor network for the Jones polynomial of a knot (ToDo: ref). We are seeking a diagram in the qutrit ZX-calculus that equals (up to a scalar) the matrix  $T_{3\pm}$  from (ToDo: ref):

**Proposition 3.7.** Under the standard interpretation as a linear map, the following diagram gives (up to a scalar) the required matrix:

$$\left[ \begin{array}{c} \text{green circle with } \frac{2}{1} \\ \text{red circle with } \frac{\pm 2}{0} \\ \text{green circle with } \frac{1}{2} \end{array} \right] \simeq \left[ \begin{array}{c} \boxed{\pm} \end{array} \right]_{q=3} = \begin{pmatrix} e^{\mp i \frac{\pi}{3}} & 1 & 1 \\ 1 & e^{\mp i \frac{\pi}{3}} & 1 \\ 1 & 1 & e^{\mp i \frac{\pi}{3}} \end{pmatrix} \quad (23)$$

*Proof.* ToDo: prove in appendix.  $\square$

Crucially, the ZX-diagram in Proposition 3.7 above is a *stabilizer diagram* in the qutrit ZX-calculus - that is, all angles are integer multiples of  $\frac{2\pi}{3}$ . Therefore if we can find an algorithm analogous to [Duncan et al., 2020, Theorem 5.4] that efficiently reduces any stabilizer diagram to a trivial one, then we will have shown that the Jones polynomial of any knot at the lattice roots of unity  $\pm e^{i\frac{\pi}{3}}$  is efficiently computable. [ToDo: justify the  $\pm$ ]. In the next subsection, we will do exactly that.

### 3.2 Graph-Like Qutrit ZX Diagrams

[Make all of the following scalar-exact (?)]

We first define a graph-like diagram in the qutrit ZX calculus.

**Definition 3.8.** A qutrit ZX-diagram is *graph-like* when:

1. Every spider is a Z-spider.
2. Spiders are only connected by Hadamard edges ( $H$ -edges) or their adjoints ( $H^\dagger$ -edges).
3. Every pair of spiders is connected by at most one  $H$ -edge or  $H^\dagger$ -edge.
4. Every input and output is connected to a spider.
5. Every spider is connected to at most one input or output.

For our specific needs, the last two items above will not be relevant, since ZX-diagrams arising from knots will be closed, but we include them to keep the definition consistent with the qubit case. However, note the difference compared to the qubit case: we need not worry about self-loops because the qutrit ZX calculus doesn't define a 'plain' cap or cup. But this comes at a cost: spiders in the qutrit case fuse more fussily. Specifically, when two spiders of the same colour are connected by at least one plain edge and at least one  $H$ - or  $H^\dagger$ -edge, fusion is not possible. The following equation, which holds with the roles of  $H$  and  $H^\dagger$  reversed, helps us get around this:

$$\text{Diagram 1} \stackrel{(\mathbf{H})}{=} \text{Diagram 2} \stackrel{(\mathbf{id})}{=} \text{Diagram 3} \quad (24)$$

We will shortly show that every qutrit ZX-diagram is equivalent to a graph-like one, making use of the following lemmas:

**Lemma 3.9.** The following equation holds in the qutrit ZX-calculus. Moreover, it holds with the roles of  $H$  and  $H^\dagger$  interchanged:

$$(25)$$

*Proof.* It is shown in [Gong and Wang, 2017, Lemma 2.8] that the qutrit ZX-calculus satisfies the following ‘Hopf law’:

$$(26)$$

Therefore we can argue as follows:

$$(27)$$

□

**Lemma 3.10.** The following two equations hold in the qutrit ZX-calculus. Moreover, they hold with the roles of  $H$  and  $H^\dagger$  interchanged:

$$(28)$$

*Proof.* This is Lemma 3.4 in Gong and Wang [2017].

□

As we will formalise later, the lemmas above say that we can think of Hadamard edges as 1-weighted edges and their adjoints as 2-weighted edges, then work modulo 3, since every triple of parallel edges disappears. This motivates defining ‘parametrised’ Hadamard gates, which will come in use later:

$$\begin{array}{ccc}
 \begin{array}{c} | \\ \text{yellow triangle} \end{array} = \begin{array}{c} \text{dashed box} \end{array} & 
 \begin{array}{c} | \\ \text{yellow triangle} \end{array} = \begin{array}{c} \text{yellow triangle} \end{array} & 
 \begin{array}{c} | \\ \text{yellow triangle with 2} \end{array} = \begin{array}{c} | \\ \text{yellow triangle} \end{array}
 \end{array} \quad (29)$$

The first equation above just says that a ‘0-Hadamard edge’ is in fact the empty diagram, and not an edge at all. Where the previous lemmas relate single  $H$ - and  $H^\dagger$ -boxes across multiple edges, the next relates multiple  $H$ - and  $H^\dagger$ -boxes on single edges.

**Lemma 3.11.** The following three equations hold in the qutrit ZX calculus. Moreover, they hold with the roles of  $H$  and  $H^\dagger$  interchanged:

$$\begin{array}{ccc}
 \begin{array}{c} \text{yellow triangle} \\ \text{yellow triangle} \\ \text{yellow triangle} \end{array} = \begin{array}{c} | \end{array} & 
 \begin{array}{c} \text{yellow triangle} \\ \text{yellow triangle} \end{array} = \begin{array}{c} | \\ \text{yellow triangle} \end{array} & 
 \begin{array}{c} \text{yellow triangle} \\ \text{yellow triangle} \end{array} = \begin{array}{c} | \\ \text{green circle} \end{array}
 \end{array} \quad (30)$$

*Proof.* For the first equation, turn the top two  $H$ -boxes into  $H^\dagger$ -boxes via (s), then cancel twice via (H). Similarly, for the second equation turn the top two  $H$ -boxes into  $H^\dagger$ -boxes via (s), then cancel once via (H), leaving a single  $H^\dagger$ -box. The third equation is just a simple application of (id). Analogous reasoning applies when the roles of  $H$  and  $H^\dagger$  are swapped.  $\square$

Again, intuitively we can think of Hadamard boxes of having value 1 and their adjoints  $-1$  and then work modulo 4.

**Corollary 3.12.** Every qutrit ZX diagram is equivalent to one that is graph-like.

*Proof.* First use the colour change rule to turn all X-spiders into Z-spiders. Then use Lemma 3.11 to remove excess  $H$ - and  $H^\dagger$ -boxes, inserting a spider between any remaining consecutive pair of such boxes, so that all spiders are connected only by plain edges,  $H$ -edges or  $H^\dagger$ -edges. Fuse together as many as possible, and apply (24) where fusion is not possible, so that no plain edge connects two spiders. Apply Lemmas 3.9 and 3.10 to all connected pairs of spiders until at most one  $H$ - or  $H^\dagger$ -edge remains between them. Finally, to ensure every input and output is connected to a spider and every spider is connected to at most one input or output, we can use (H) and (id) to add a few spiders,  $H$ - and  $H^\dagger$ -boxes as needed:

$$\begin{array}{ccc}
 \begin{array}{c} | \\ \text{green circle} \\ \text{yellow triangle} \\ \text{green circle} \\ \text{yellow triangle} \end{array} = \begin{array}{c} | \\ \text{green circle} \\ \text{yellow triangle} \\ \text{green circle} \\ \text{yellow triangle} \end{array} & 
 \begin{array}{c} | \\ \text{yellow triangle} \end{array} = \begin{array}{c} | \\ \text{green circle} \end{array} & 
 \begin{array}{c} \text{green circle with } \alpha, \beta \end{array} = \begin{array}{c} \text{green circle with } \alpha, \beta \\ \text{yellow triangle} \\ \text{green circle} \\ \text{yellow triangle} \end{array}
 \end{array} \quad (31)$$

$\square$

**Definition 3.13.** A graph-like qutrit ZX diagram is a *graph state* when every spider has zero phase (top and bottom) and is connected to an output.

A graph state is described fully by its underlying multigraph, or equivalently by an adjacency matrix, where edges take weights in  $\mathbb{Z}_3$  [Wang, 2018, Lemma 4.2]. Nodes correspond to phaseless green spiders, edges of weight 1 correspond to Hadamard edges, and edges of weight 2 correspond to  $H^\dagger$  edges. As in the qubit case, graph states admit a local complementation operation [Wang, 2018, Definition 2.6], though the effect is now slightly more complicated. We'll give the intuition after the formal definition:

**Definition 3.14.** Given  $a \in \mathbb{Z}_3$  and a graph state  $G$  with adjacency matrix  $W = (w_{i,j})$ , the  $a$ -local complementation at node  $k$  is the new graph state  $G *_a k$ , whose adjacency matrix  $W' = (w'_{i,j})$  given by:

$$w'_{i,j} = w_{i,j} + aw_{i,k}w_{j,k} \quad (32)$$

So only those edges between neighbours of node  $k$  are affected, but rather than just having their weight increased by 1 (modulo 2) as in the qubit case, the increase in weight also depends on the weights of the edges from  $i$  and  $j$  to  $k$ . As always, this is best seen graphically. We reintroduce the blue dashed line notation for Hadamard edges, and now also use purple dashed lines for  $H^\dagger$ -edges:

$$\begin{array}{c} | \\ \text{---} \end{array} = \begin{array}{c} | \\ \text{---} \end{array} \quad \begin{array}{c} | \\ \text{---} \end{array} = \begin{array}{c} | \\ \text{---} \end{array} \quad (33)$$

For two nodes  $i$  and  $j$  both connected to  $k$  by the same colour edge,  $a$ -local complementation at  $k$  increases weight  $w_{i,j}$  by  $a$ . If instead  $i$  and  $j$  are connected to  $k$  by edges of different colour,  $a$ -local complementation at  $k$  decreases  $w_{i,j}$  by  $a$ . We show a fragment of a ZX-diagram below under the effect of this operation:

$$\begin{array}{ccc} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & \mapsto & \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \end{array} \quad \begin{array}{ccc} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & \mapsto & \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \end{array} \quad (34)$$

But the fragment above doesn't give the full picture. As in the qubit case, local complementation gives an equality up to introducing some single qubit phase gates on the outputs.

**Theorem 3.15.** Given  $a \in \mathbb{Z}_3$  and a graph state  $(G, W)$  containing a node  $k$ , let  $N(k)$  denote the neighbours of  $k$  - that is, nodes  $i$  with weight  $w_{i,k} \in \{1, 2\}$ . Then the following equality holds:

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

*Proof.* This is Theorem 4.4 and Corollary 4.5 in Wang [2018].  $\square$

Composing local complementations gives a local pivot operation.


**Definition 3.16.** Given  $a, b, c \in \mathbb{Z}_3$  and a graph state  $G$  containing nodes  $i$  and  $j$ , the  $(a, b, c)$ -local pivot along  $ij$  is the new graph state  $G \wedge_{(a,b,c)} ij := ((G *_{a,i}) *_{b,j}) *_{c,i}$ .

This again results in an equality, up to introducing some extra gates on outputs. Here we shall only consider an  $(a, -a, a)$ -local pivot along an edge  $ij$  of non-zero weight, for  $a \in \{1, 2\}$ . We will call this a *proper  $a$ -local pivot* along  $ij$ , and denote it  $G \wedge_a ij$ .

**Theorem 3.17.** Given  $a \in \mathbb{Z}_3$  and a graph state  $(G, W)$  containing connected nodes  $i$  and  $j$ , define the following:

- $N_=(i, j) := \{k \in N(i) \cap N(j) \mid w_{k,i} = w_{k,j}\}$
- $N_\neq(i, j) := \{k \in N(i) \cap N(j) \mid w_{k,i} \neq w_{k,j}\}$

Then the following equation relates  $G$  and its proper  $a$ -local pivot along  $ij$ :




*Proof.* See A.1 in the Appendix.  $\square$

These two operations are again the drivers behind the simplification procedure. We classify spiders into three families:

$$\mathcal{M} = \left\{ \begin{pmatrix} \cdots \\ 0 \\ 0 \\ \cdots \end{pmatrix}, \begin{pmatrix} \cdots \\ 1 \\ 2 \\ \cdots \end{pmatrix}, \begin{pmatrix} \cdots \\ 2 \\ 1 \\ \cdots \end{pmatrix} \right\}, \quad \mathcal{N} = \left\{ \begin{pmatrix} \cdots \\ 0 \\ 1 \\ \cdots \end{pmatrix}, \begin{pmatrix} \cdots \\ 1 \\ 0 \\ \cdots \end{pmatrix}, \begin{pmatrix} \cdots \\ 0 \\ 2 \\ \cdots \end{pmatrix}, \begin{pmatrix} \cdots \\ 2 \\ 0 \\ \cdots \end{pmatrix} \right\}, \quad \mathcal{P} = \left\{ \begin{pmatrix} \cdots \\ 1 \\ 1 \\ \cdots \end{pmatrix}, \begin{pmatrix} \cdots \\ 2 \\ 2 \\ \cdots \end{pmatrix} \right\}.$$

exactly as in [Wang, 2018, Theorem 3.1]. We call a spider in a graph-like ZX-diagram *interior* if it isn't connected to an input or output (so for our use case all spiders are interior). Given any graph-like ZX-diagram, we will show that we can eliminate pairs of connected interior  $\mathcal{M}$ -spiders by local pivoting, and standalone interior  $\mathcal{N}$ - and  $\mathcal{P}$ -spiders by local complementation.


**Theorem 3.18.** Given any graph-like ZX-diagram containing two interior  $\mathcal{M}$ -spiders  $i$  and  $j$  connected by edge  $ij$  of weight  $w_{i,j} =: w \in \{1, 2\}$ , suppose we perform a proper  $\pm w$ -local pivot along  $ij$ . Then the new ZX-diagram is related to the old one by the equality:




where all changes to weights of edges where neither endpoint is  $i$  or  $j$  are omitted. Furthermore, in order to save space, each node with phase  $\begin{pmatrix} a_x^y \\ b_x^y \end{pmatrix}$  is representative of *all* nodes connected to  $i$  by an  $x$ -weighted edge and to  $j$  by a  $y$ -weighted edge.

*Proof.* See A.2 in the Appendix.  $\square$

**Theorem 3.19.** Given any graph-like ZX-diagram containing an interior  $\mathcal{N}$ -spider  $k$  with phase  $\begin{pmatrix} 0 \\ n \end{pmatrix}$  for  $n \in \{1, 2\}$ , suppose we perform a  $(-n)$ -local complementation at  $k$ . Then the new ZX-diagram is related to the old one by the equality:




where all changes to weights of edges where neither endpoint is  $k$  are omitted. If instead  $k$  has phase  $\begin{pmatrix} n \\ 0 \end{pmatrix}$  for  $n \in \{1, 2\}$ , suppose we perform the same  $(-n)$ -local complementation at  $k$ . Then the equality relating the new and old diagrams becomes:



*Proof.* See A.3 in the Appendix.  $\square$

**Theorem 3.20.** Given any graph-like ZX-diagram containing an interior  $\mathcal{P}$ -spider  $k$  with phase  $\begin{pmatrix} p \\ p \end{pmatrix}$  for  $p \in \{1, 2\}$ , suppose we perform a  $p$ -local complementation at  $k$ . Then the new ZX-diagram is related to the old one by the equality:



where all changes to weights of edges where neither endpoint is  $k$  are omitted.

*Proof.* See A.4 in the Appendix.  $\square$

We can now combine these three elimination theorems into an algorithm for efficiently simplifying a closed graph-like ZX-diagram. First note that after applying any one of the three elimination theorems to such a diagram we may end up with a state that is no longer graph-like. Fortunately the only way in which this can happen is if two spiders end up being connected by multiple  $H$ - or  $H^\dagger$ -edges, and we have shown via Lemmas 3.9 and 3.10 that these can always be reduced to just one edge.

**Theorem 3.21.** Given any closed graph-like ZX-diagram, the following algorithm will always terminate after a finite number of steps, returning an equivalent graph-like ZX-diagram with no  $\mathcal{N}$ -spiders,  $\mathcal{P}$ -spiders, or adjacent pairs of  $\mathcal{M}$ -spiders. Repeat the steps below until no rule matches. After each step, apply Lemmas 3.9 and 3.10 as needed until the resulting diagram is graph-like:

1. Eliminate an  $\mathcal{N}$ -spider via Theorem 3.19.
2. Eliminate a  $\mathcal{P}$ -spider via Theorem 3.20.
3. Eliminate two adjacent  $\mathcal{M}$ -spiders via Theorem 3.18.

*Proof.* At every step the total number of spiders decreases by at least one, so since we start with a finite diagram the algorithm terminates after a finite number of steps. By construction, when it does so we are left with an equivalent graph-like ZX-diagram with no  $\mathcal{N}$ -spiders,  $\mathcal{P}$ -spiders, or adjacent pairs of  $\mathcal{M}$ -spiders.  $\square$

**Corollary 3.22.** In particular, if we start with a stabilizer diagram, we can eliminate all but perhaps one  $\mathcal{M}$ -spider, depending on whether the initial number of  $\mathcal{M}$ -spiders was odd or even.

*Proof.* No step introduces any non-stabilizer phases.  $\square$

The algorithm above could be extended to graph-like diagrams with inputs or outputs as in [Duncan et al., 2020, Theorem 5.4], but since for our purposes we don't need to do so, we have not gone to the trouble.



## References

- B. Coecke and A. Kissinger. *Picturing Quantum Processes*. 2017.
- R. Duncan, A. Kissinger, S. Perdrix, and J. van de Wetering. Graph-theoretic simplification of quantum circuits with the zx-calculus. 2020.
- X. Gong and Q. Wang. Equivalence of local complementation and euler decomposition in the qutrit zx-calculus. 2017.
- Q. Wang. Qutrit zx-calculus is complete for stabilizer quantum mechanics. 2018.


## A Appendix

We begin with a proof of the local pivot equality from Theorem 3.17.

**Theorem A.1. / Theorem 3.17.** Given  $a \in \mathbb{Z}_3$  and a graph state  $(G, W)$  containing connected nodes  $i$  and  $j$ , define the following:

- $N_=(i, j) := \{k \in N(i) \cap N(j) \mid w_{k,i} = w_{k,j}\}$
- $N_\neq(i, j) := \{k \in N(i) \cap N(j) \mid w_{k,i} \neq w_{k,j}\}$


Then the following equation relates  $G$  and its proper  $a$ -local pivot along  $ij$ :




*Proof.* Annoyingly, proving this in full generality in one go - i.e. for a proper  $a$ -local pivot along an edge  $ij$  of weight  $b$  - becomes a bit tricky diagrammatically, because it becomes hard to keep track of all the variable edge weights. Fortunately the four cases  $(a, b \in \{1, 2\})$  split into two pairs of symmetric cases:  $a = b$  and  $a \neq b$ .


Now, it suffices to only draw a fragment of a graph state. Certainly we consider nodes  $i$  and  $j$  and the edge  $ij$  between them. Then define  $N_x^y$  to be the set  $\{k \mid w_{k,i} = x, w_{k,j} = y\}$ , for  $x, y \in \mathbb{Z}_3$ . We will consider a representative node  $k_x^y$  from each  $N_x^y \neq N_0^0$ , as well as its edges  $ik_x^y$  and  $jk_x^y$ . All other nodes and edges are irrelevant; this is because we are only interested in nodes and edges that *affect* the three local complementation operations on  $i$  and  $j$  - we aren't concerned with those that are only *affected by* the operations.

So for the case  $a = b$ , we show the proper 1-local pivot along  $ij$  of weight 1:



The story is similar for the case  $a \neq b$ ; here we show the proper 1-local pivot along  $ij$  of weight 2:






The case  $a = b = 2$  is the same as  $a = b = 1$ , except with the roles of purple and blue edges interchanged, so by symmetry of the diagram the only difference is in the phase gates added to the outputs. Namely, on each phase gate we replace all instances of 1 with a 2. Thus the roles of  $H$  and  $H^\dagger$  are swapped too. Likewise for the case  $(a, b) = (2, 1)$  with respect to the case  $(a, b) = (1, 2)$ .

□

Next we prove the three elimination theorems for  $\mathcal{M}$ -,  $\mathcal{N}$ - and  $\mathcal{P}$ -spiders.


**Theorem A.2. / Theorem 3.18.** Given any graph-like ZX-diagram containing two interior  $\mathcal{M}$ -spiders  $i$  and  $j$  connected by edge  $ij$  of weight  $w_{i,j} =: w \in \{1, 2\}$ , suppose we perform a proper  $\pm w$ -local pivot along  $ij$ . Then the new ZX-diagram is related to the old one by the equality:




$$\begin{aligned}
&= \begin{array}{c} \text{red circle } \begin{array}{c} m_i \\ -m_i \end{array} \\ \text{yellow triangle } w \\ \text{red circle } \begin{array}{c} -m_j \\ m_j \end{array} \end{array} \quad \begin{array}{c} \left( \begin{array}{c} \dots \\ a_1^0 + m_j \\ b_1^0 - m_j \end{array} \right) \left( \begin{array}{c} \dots \\ a_2^0 - m_j \\ b_2^0 + m_j \end{array} \right) \dots \left( \begin{array}{c} \dots \\ a_0^1 + m_i \\ b_0^1 - m_i \end{array} \right) \left( \begin{array}{c} \dots \\ a_0^2 - m_i \\ b_0^2 + m_i \end{array} \right) \end{array} \\
&\quad \begin{array}{c} \left( \begin{array}{c} \dots \\ a_1^1 \mp w + m_j + m_i \\ b_1^1 \mp w - m_j - m_i \end{array} \right) \left( \begin{array}{c} \dots \\ a_2^1 \pm w + m_j - m_i \\ b_2^1 \pm w - m_j + m_i \end{array} \right) \left( \begin{array}{c} \dots \\ a_2^1 \pm w - m_j + m_i \\ b_2^1 \pm w + m_j - m_i \end{array} \right) \left( \begin{array}{c} \dots \\ a_2^2 \mp w - m_j - m_i \\ b_2^2 \mp w + m_j + m_i \end{array} \right) \end{array}
\end{aligned}$$

where all changes to weights of edges where neither endpoint is  $i$  or  $j$  are omitted. Furthermore, in order to save space, each node with phase  $\frac{a_x^y}{b_x^y}$  is representative of *all* nodes connected to  $i$  by an  $x$ -weighted edge and to  $j$  by a  $y$ -weighted edge.

*Proof.* We show the case where  $w_{ij} =: w = 1$ , with the case  $w = 2$  being completely analogous. We can choose either a proper 1-local pivot or a proper 2-local pivot; both give the same result. Here we only show the former:





$$\begin{aligned}
&= \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \\
&= \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}
\end{aligned}$$

The diagrams show a sequence of transformations. The first diagram shows a red node with phase  $\frac{m_i}{-m_i}$  connected to a yellow node with phase  $\frac{-m_j}{m_j}$ , which is then connected to a complex network of green nodes. The second diagram shows a similar structure but with different phases and connections, illustrating the result of a  $(-n)$ -local complementation.

□

**Theorem A.3. / Theorem 3.19.** Given any graph-like ZX-diagram containing an interior  $\mathcal{N}$ -spider  $k$  with phase  $\frac{0}{n}$  for  $n \in \{1, 2\}$ , suppose we perform a  $(-n)$ -local complementation at  $k$ . Then the new ZX-diagram is related to the old one by the equality:

$$\begin{array}{c} \text{Diagram 1} \end{array} = \begin{array}{c} \text{Diagram 2} \end{array}$$

The diagram shows a central node  $\frac{0}{n}$  connected to several nodes  $\frac{a_i}{b_i}$  and  $\frac{c_i}{d_i}$ . The second diagram shows the result of the  $(-n)$ -local complementation, where the weights of the edges are updated according to the theorem.


where all changes to weights of edges where neither endpoint is  $k$  are omitted. If instead  $k$  has phase  $\frac{n}{0}$  for  $n \in \{1, 2\}$ , suppose we perform the same  $(-n)$ -local complementation at  $k$ . Then the equality relating the new and old diagrams becomes:

$$\begin{array}{c} \text{Diagram 1} \end{array} = \begin{array}{c} \text{Diagram 2} \end{array}$$

The diagram shows a central node  $\frac{n}{0}$  connected to several nodes  $\frac{a_i}{b_i}$  and  $\frac{c_i}{d_i}$ . The second diagram shows the result of the  $(-n)$ -local complementation, where the weights of the edges are updated according to the theorem.


*Proof.* We prove the case where  $k$  has phase  $\frac{0}{n}$  for  $n \in \{1, 2\}$ , the other case being near

identical.



□

**Theorem A.4. / Theorem 3.20.** Given any graph-like ZX-diagram containing an interior  $\mathcal{P}$ -spider  $k$  with phase  $\frac{p}{p}$  for  $p \in \{1, 2\}$ , suppose we perform a  $p$ -local complementation at  $k$ . Then the new ZX-diagram is related to the old one by the equality:



where all changes to weights of edges where neither endpoint is  $k$  are omitted.

*Proof.*

